

Review article

Genistein Mediated Molecular Pharmacology, Cell-Specific Anti-Breast Cancer Mechanism with Synergistic Effect and *in silico* Safety Measurement

Md Sohel¹, Partha Biswas^{2,3}, Md. Al Amin⁴, Dipta Dey⁵, Habiba Sultana⁴, Suraiya Aktar⁶,
Md. Sharif Khan⁴, Md. Arju Hossain⁴, Priyanka Paul⁵, Nurul Islam⁷, Md. Aatur
Rahman^{8,9,10*}, Bonglee Kim^{9,10*}, Abdullah Al Mamun^{1*}

¹Department of Biochemistry and Molecular Biology, Mawlana Bhashani Science and Technology University, Santosh, Tangail-1902, Bangladesh.

²Department of Genetic Engineering and Biotechnology, Faculty of Biological Science and Technology, Jashore University of Science and Technology (JUST), Jashore-7408, Bangladesh.

³ABEx Bio-Research Center, East Azampur, Dhaka-1230, Bangladesh.

⁴Department of Biotechnology and Genetic Engineering, Faculty of life science, Mawlana Bhashani Science and Technology University, Santosh, Tangail-1902, Bangladesh.

⁵Department of Biochemistry and Molecular Biology at Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj-8100, Bangladesh.

⁶Department of Biochemistry and Molecular Biology, Rajshahi University, Rajshahi, Bangladesh.

⁷Department of Pharmacy, Faculty of Life Science, Mawlana Bhashani Science and Technology University, Santosh, Tangail-1902, Bangladesh.

⁸Global Biotechnology & Biomedical Research Network (GBBRN), Department of Biotechnology and Genetic Engineering, Faculty of Biological Sciences, Islamic University, Kushtia-7003, Bangladesh.

⁹Department of Pathology, College of Korean Medicine, Kyung Hee University, Seoul-02447, South Korea.

¹⁰Korean Medicine-Based Drug Repositioning Cancer Research Center, College of Korean Medicine, Kyung Hee University, Seoul-02447, Korea.

*Corresponding Author:

Md. Aatur Rahman, Email: ataur1981rahman@hotmail.com

Bonglee Kim, Phone: E-mail: bongleekim@khu.ac.kr

Abdullah Al Mamun, Email: mamunbmb@mbstu.ac.bd

Abstract

Breast cancer (BC) is the most common type of cancer in both men and women alike, but it is more prevalent in women. Natural compounds that can modulate the oncogenic process can be considered a significant anti-cancer agent for treating BC. These natural compounds are more effective than synthetic drugs, which have profound side effects on the normal cell and resistance to cancer cells. Genistein is a type of dietary phytoestrogen included in the flavonoid group with a similar structure of estrogen that might provide a strong alternative and complementary medicine to existing chemotherapeutic drugs. Several research studies demonstrated that it can target the estrogen receptor (ER), Human epidermal growth factor receptor-2 (HER2), and Breast cancer gene-1 (BRCA-1) in multiple BC cell lines, as well as sensitize cancer cell lines to this compound when used at an optimal inhibitory concentration. Genistein effectively showed anti-cancer activities through apoptosis induction, arresting cell cycle, inhibiting angiogenesis with metastasis, reducing inflammation, mammosphere formation, tumor growth, up-regulating tumor suppressor gene, and downregulating oncogene in suppressing cancer progression in vitro and animal model study. In addition, research studies have also suggested that these phytochemicals synergistically reverse the resistance mechanism of chemotherapeutic drugs, increasing the efficacy of some chemoinformatics drugs. Our review article aims to unbox and validate the molecular pharmacology in breast tissue, cell-specific anti-cancer mechanism with synergistic activity, and possible pharmacokinetic parameters of Genistein as a potential alternative therapeutic option for the treatment of BC.

Keywords: Genistein; Estrogen receptor; BRCA-1; Cell cycle; Tumor suppressor gene; Molecular pharmacology; Synergistic activity.

1. Introduction:

Breast cancer (BC) is considered a major public health concern among women in health science. BC is responsible for 30% of all cancers in females and is the world's second leading cause of mortality [1]. Although they're diverse risk factors, which can increase the possibility of developing BC including aging, sex, gene mutations, family history, and unhealthy lifestyle, [2], abnormal hormones namely estrogen play an effective role in BC progression [3]. There are currently no established treatments are available, and having some important side effects including-reduction of blood cells, [4], sore throat, hair loss, ulcers, fatigue, nausea, change in taste, appetite loss, constipation, diarrhea, change in skin color, and changes of several hormonal levels [5] and some limitations of existing treatment are high cost, less effective, allergic reaction [6]. Multidrug-resistant (MDR) tumors are the major limitation of conventional treatment, leading to increased cancer-related deaths. Numerous drug molecules likely- anthracyclines (daunorubicin, doxorubicin, mitoxantrone, and epirubicin), taxanes (docetaxel, paclitaxel), and capecitabine were previously used successfully, but now these drugs are becoming resistant to cancer patients [7]. So, the modern medical emphasis is on preventing and treating breast cancer with natural nutritional components [8] [9] [10]. Under the pharmaceutical compounds- phytoestrogens are a natural dietary component with having potent anti-cancer activity against multiple cancer; most importantly, ovarian, prostate, as well as breast [11][12].

One of the major soy phytoestrogens is Genistein (Gen), which has been described as a breast cancer preventive factor [13]. Soy-based foods, for example-soy cheese, and soy drinks i.e., soy milk with soy-based beverages, are the plentiful sources of Gen; of late, the amount of Gen content within mature soybeans has shown from 5.6 to 276 mg/100 g, but 81 mg/100 g is an average content [14], where regular daily intake of this isoflavones among adults in China, and Japan is about 25–50 mg which is several-folds higher than the consumption of these compounds by women in the western countries which accounted for less than 3 mg [15]. Gen has a structural similarity to 17β -estradiol and it binds to estrogen receptor ER- β compare to ER- α [16][17]. Diverse studies suggested the Gen can directly inhibit the cell cycle [18], induce cellular apoptosis process [19][20], suppressed metastasis [21] angiogenesis [22], inhibit inflammation [23], reduce oxidative stress, inhibit tumor growth, and mammosphere formation [24] for both estrogen-responsive, as well as non-responsive BC cells *in vitro* and several animal models.

Furthermore, this phytoestrogen has several synergistic effects, where Gen can enhance the efficacy of breast cancer therapy along with reducing chemotherapy drug resistance [25]. Major Synergistic effect of Gen with Doxorubicin, Tamoxifen, Trastuzumab, and so on; of later, their activity rely on dose concentration[26][27][28] . Consequently, Gen effects on markers of cancer risk, which were observed from the numerous clinical studies in humans both *in vitro* and *in vivo* clinical investigation respectively[29][30][31][32][33].Genistein safety measurement and future predictions were discovered by using *in silico* assessment. It possesses a better bioavailability score of 0.55. Similarly, Gen is highly hydrophobic and absorbed by human intestinal, with permeable to Caco-2 cell. In addition, its capacity to be well distributed in the various tissues, also excrete little amount after the metabolism of Gen. Finally, Gen has no severe toxic effect on the human body which was analyzed by the online server, and therefore, it's maintained the drug-like characteristics without any violating of Lipinski, Ghose Veber, Muegge, and Egan rules.

Therefore, in this current investigation, molecular anti-cancer mechanisms of Genistein in multiple BC cells were analyzed, mechanism of resistance drug recovery with synergistic effects, clinical trials, and pharmacokinetics was predicted using Schrodinger's QuickPro modules and online accessible server admetSAR, SwissADME, and pKCSM for the better understanding of Genistein and breast cancer treatment.

2. Molecular pharmacology of Genistein in breast tissue

Genistein is a natural phytochemical that belong to phytoestrogen and possesses a similar structure to estrogen and most interestingly has both mimic and antagonize estrogen effects, at the same time it inhibits the BC cell proliferation at high concentrations [34]. Estrogen receptor-mediated growth of BC cells by Gen is concentration-dependent. T. T. Y. Wang et al. summarized that Genistein stimulated growth at lower concentrations (10^{-8} ~ 10^{-6} M), but Gen inhibited cancer cell growth at higher concentrations ($>10^{-5}$ M) [35]. There are two types of estrogen receptors [36]. Genistein has a similar structure both of these types of receptors but binds to estrogen receptor- β (ER- β) with higher affinity compared to estrogen receptor- α (ER- α)[16][17].

In the case of estrogen receptor- α , Genistein acts as an antagonist. So, Genistein-mediated anti-cancer activity is involved in suppressing the expression and activity of estrogen receptor- α . E. J. Choi summarised that Genistein regulates cell proliferation with apoptosis via ER- α dependent pathway in MCF-7 breast cancer cells through the underline mechanism of downregulating Cyclin D1 and up-regulating Bcl-2/Bax (full elaboration add here) ratio at the dose of 50 μ M [37]. But in ER- β , Gen plays as an agonist activity and increases the receptor activity. Therefore, ER- β dependent anti-cancer activity of Genistein mediated by activating the receptor and potentiate chemotherapy efficacy to treat cancer treatment [38]. H. Jiang state that ER- β 1 increased the anti-cancer efficacy of Gen in MDA-MB-231, MCF-7, cells, and in BALB/c mice by inhibiting cell proliferation through arresting cells in G2/M and G0/G1 phase, which led to cell cycle blockade at the dose of 10^{-6} - 10^{-4} mol/l [39]. It has also been found to bind with the estrogen-responsive G protein-coupled receptor-30 (GPR-30) or G protein-coupled estrogen receptor-1 (GPER-1) [40], and inhibit cell proliferation [41]. HER-2 is an essential biomarker in breast cancer overexpressed in around 20–30% of breast cancer tumors [42]. So, regulating the HER-2 is a significant factor in BC treatment. M. S. Sakla et al. summarised that Genistein at $>$ or $=1$ microM inhibited proto-oncogene of HER-2, and subsequently followed the HER-2 protein expression, phosphorylation, and promoter activity through an ER-independent mechanism in BC cells as delaying tumors onset in transgenic mice [43]. Therefore, GPR-30 is a potent alternate to ER that causes BC and can be a target in breast cancer cell treatment [44]. Kim GY et al. summarized that Genistein suppresses the GPR-30 activation in BRCA-1 mutated BC cells, resulting in G2/M phase arrest mediated by suppressing Akt phosphorylation [45]. Furthermore, Gen is a naturally occurring protein tyrosine kinase (PTK) inhibitor that is hypothesized to be responsible for the lower rate of BC observed in Asian women consuming soy. T. Akiyama et al. reported that Gen scarcely inhibited the enzyme activities of threonine- and serine-specific protein kinases like cAMP-dependent protein kinase, Ca^{2+} /phospholipid-dependent enzyme protein kinase C, and phosphorylase kinase, and this mechanism are mediated through phosphorylation of the EGF receptor [46].

3. Cell-specific molecular mechanism of Genistein mediated anti-breast cancer activity *in vitro*

Cancerous cell lines derived from humans are critical models for cancer biology research in laboratories and determine the therapeutics advantage of anti-cancer agents [47]. Anti-cancer activity of phytochemicals is cell-specific, where one phytochemical is effective in only one or more than one cell line. This may be the difference in the cell components system. In this section, cell line-specific anti-cancer activity of Genistein is summarized in **Fig 01**.




Figure 01. Graphical overview of the anti-cancer mechanisms of Genistein: Genistein induces apoptosis through mitochondrial mediated classical caspase-dependent pathway with modulating Bcl-2 family proteins. It induces cell cycle arrest by modulating the cycle regulatory proteins. It inactivates signaling pathways, namely PI3K/AKT pathway, as well as MAPK (ERK1/2) pathways. Formononetin also modulates several miRNA expressions as well as suppresses cell migration, invasion, and angiogenesis and regulate epigenetics control.

3.1. Genistein effect on MCF-7 BC cell

According to R. F. Prietsch *et al.*, Gen (0.01-100 μ m) promoted apoptosis via mediating the autophagy-dependent mechanism along with increasing the ratio of Bax/Bcl-2 and inhibit the oxidative stress of cancer progression through changing the expression of antioxidant enzymes. For example, Gen down-regulates the thioredoxin reductase enzyme (TrxR) and up-regulates the expression of Glutathione peroxidase (GPx) [48]. X. Liu et al. summarized that Genistein (5-20 μ m) induces apoptosis through the mitochondrial-dependent pathway by decreasing the Bcl-2/Bax ratio, increasing tumor suppressor gene P⁷³ expression and ATM phosphorylation with G2/M phase arrest permanently [49]. Similarly, Genistein (50-200 μ m) hinders the cellular growth and induces the apoptosis mediated pathway by following the downregulation of Bcl-2 protein, as well as upregulation of the Bax, together with decreasing the cyclin D1 expression in MCF-7 breast cancer cell [37]. At low concentration Gen (1 μ M) stimulate cell proliferation but at higher concentration (25 μ M) induces apoptosis mechanism by up-regulating of the CDKN1A and p53 responsive gene and down-regulating the CCNG1 GADD45A, B NF-kappa B, BCL-2, and TNFR, ESR1, NCOA2, and NCOA3 [50]. Another research investigated that Gen (50 μ M) highlights apoptosis by up-regulating Poly-(ADP-ribose)-polymerase, p53, and down-regulating Bcl-2/Bax protein [26]. *In vitro* study by M. L. De Lemos investigated that Gen (10 μ M) induces apoptosis by breaking plasma membrane breakdown, nuclear membrane, and up-regulating pS2 expression [27]. A late study reported that Gen (100 μ M) induces programmed cell death and suppresses cell growth by up-regulating Caspase 7, apoptosis signalling kinase-1, ADP ribose, and p38 dependent mitogen protein kinase [51].

Inhibition of metastasis is common in breast cancer treatment. *In vitro* study demonstrated that Gen (3.125-12.5 μ M) decrease metastasis with increasing GSTP1 and RAR β 2 gene expression and activity [52]. Shon et al. concluded that Gen (50 μ M) suppressed angiogenesis by downward COX, Topoisomerase II enzyme and TPA, EROD protein [53], while Gen (1-10 μ g/ml) inhibit angiogenesis and cell mutation by decreasing Tyrosine kinase, Ribosomal S6 kinases, and DNA topoisomerases I, II [54]. *In vitro* study, Genistein (10nm-10 μ m) lowered cell proliferation via mitochondrial-dependent pathway by reducing Fis1 (mitochondrial fission), and Opa1 (mitochondrial fusion) mRNA expression [55], while (4-10mol/l) of Gen inhibits cell proliferation by downregulating the cyclin D1 and arresting the cell cycle G0/G1 phase, resulting

in blockage of the cell survival by H. Jiang et al. [39]. J. Chen et al. reported that Gen (5-100 μ m) inhibited the proliferating cells by inducing apoptosis through IGF-1R-PI3 K/Akt-mediated pathway inactivation and up-regulating Bax/Bcl-2 ratio [56]. Furthermore, it has been shown that Genistein (5-30 μ m) inhibits BC cell growth, proliferation and promotes apoptosis by following the downregulation of the Hedgehog–Gli1 signaling pathway by decreasing the mRNA level of Smo and Gli1[57]. Marik et al. also find similar result, Gen (0.1 μ M) at low concentration stimulate Cancer progression, but Gen (20 μ M) at high concentration inhibits cell proliferation by down lifting mRNA expression of ER- α protein and arrest the cell cycle at the G2/M phase [34]. Chinni et al. examined that Gen (100 μ M) inhibits cell proliferation by downregulating Akt mediated signaling pathways, telomere length, and overexpression of cyclin-dependent kinase inhibitor p21WAF1 [58].

An early study demonstrated that Gen (50 μ M) inhibits tumor growth with apoptosis inductions by increasing Ca²⁺dependent proapoptotic proteases, μ -calpain, and caspase-12 [59]. On the other hand, Liao et al. showed that Gen (100 μ M) inhibits cell growth alongside decreasing Paclitaxel-induced tubulin Polymerization, Bcl-2, Cyclin B1, and CDK2 kinase, leading to cell cycle arrest at G2/M phase [60]. Chen et al. showed that Gen (50-100 μ M) inhibits cell division through uplifting HSP protein activity and down lifting SRF mRNA, RAG-1, DOC 2 expression [61]. In breast cancer stem cells, Genistein (40nm -2 μ m) inhibits mammosphere formation by suppressing PI3K/Akt signaling with up-regulating the PTEN expression[24]. Similar result found by Y. Liu et al. confirmed that Genistein (40nm-2 μ m) inhibits mammosphere formation and induces differentiation by activating PI3K/Akt and MEK/ERK signaling in a paracrine manner increases E-cadherin mRNA expression by reducing the ratio of CD44+/CD24-/ESA in MCF-7 breast cancer cell [62]. Genistein (1 μ m) induces an anti-cancer effect through up-regulating pro-inflammatory genes, i.e pS2 and COX2, and downregulating anti-inflammatory gene expression i.e., TFG β and PPAR γ in MCF-7 breast cancer cell [63]. Furthermore, Kazi et al. appeared that Gen (50-200 μ M) hold back cancer progression by up-regulating I κ B- α , p27Kip1 level, and downregulating proteasomal chymotrypsin-like activity, CDKs [64]. Epigenetics regulation by Gen (60-100 μ m) is mediated by diminishing DNA methylation levels, DNMT1 expression, DNA methyltransferase enzyme activity. However, this reduction of DNA methylation occurs in the promoter region of multiple Tumor suppressor genes (TSGs) such as

Adenomatous polyposis coil (APC), Ataxia telangiectasia mutated (ATM), phosphatase- and tensin homolog (PTEN), mammary serpin peptidase inhibitor (SERPINB5) [65].

3.2. Genistein effect on MDA-MB-231 BC cell

Recently an experimental conducted by X. Liu et al. Gen (5-20 μ m) induces apoptosis through the mitochondrial-dependent pathway by reducing the Bcl-2/Bax ratio and growth inhibition with decreasing expression of mutant p53 and increasing the expression of p73, leading to the activation of G2/M phase arrest and ATM/Cdc25C/Chk2/Cdc2 checkpoint pathway [49]. Genistein induces the apoptosis process and directly inhibits the growth of cells through the prevention of NF- κ B activity by Notch-1 pathway and with downregulating cyclin B1, Bcl-xL, and Bcl-2 expression, resulting in the arrest of the cell cycle at G2/M phase at 5-20 μ m [66], while at 5-50 μ m, this phytochemicals induces apoptosis by targeting the endogenous copper ion, reducing Cu(II) to Cu(I) through the production of reactive oxygen species (ROS) [67]. Before that, *in vitro* study by K. Dampier et reported that Gen (10 μ M) induces apoptosis and inhibits the rapidly proliferating cells, cell cycle arrest at G2 phase with degrading proto-oncogene c-fos, prohibited protein-1 (AP-1), and also ERK activity [68]. Another research by X. Yang et al. demonstrated that Gen (50 μ M) highlights apoptosis by up-regulating Poly-(ADP-ribose)-polymerase, activating p53, and downregulating Bcl-2/ Bax protein [26].

In the case of angiogenesis, Mukund et al. explained that Genistein (100 μ m) reduced angiogenesis by blocking the transactivation of downstream HIF-1 α effectors, e.g., VEGF, leading to the reduction of hypoxia-inducible factor-1 α expression in MDA-MB-231 BC cell [22]. Furthermore, 1-10 μ g/ml of Gen suppressed angiogenesis and cell mutation by decreasing tyrosine kinase, Ribosomal S6 kinases, and DNA topoisomerases I, II [54], while at 50 μ M concentration decreased angiogenesis with inhibiting cell division through the underline mechanism of downregulating COX, Topoisomerase II enzyme, and TPA, EROD protein activity [53]. Followed by angiogenesis, Gen (15-30 μ M) [21], (5-20 μ M) [20] obstructs cancer cell migration and invasion respectively through down lifting levels of CDKs, Tyrosine kinase, DNA topoisomerase II, and Paracrine stimulation and decreasing MEK5, ERK5, Phospho-ERK5, NF- κ B/p65, and Bcl-2/ Bax. Another study done by O. C. Kousidou et al. reported that Gen (35-100 μ M) progresses slowdown invasiveness by decreasing MMP gene expression, PTKs activity, and glucose uptake rate leading to phagocytosis cancer cell [69]. Apart from this,

inhibition of cell viability through decreasing the DNA methyltransferase activity, DNMT1 expression and affecting the expression of TSGs likely- APC, ATM, PTEN, SERPINB5 by at 60-100 μ m of Genistein [65]. Another recent study by D. G. Pons *et al.* summarized that Genistein (1 μ m) causes a considerable decrease in cell viability through the mitogen-dependent protein kinase pathway that can promote apoptosis [63].

In MDA-MB-231, BC cell growth controlling is a significant target for Genistein. Gong et al. state that Gen (5-50 μ M) inhibits cell growth by partly inducing apoptosis via downregulation of the Akt, NF- κ B cascade pathways *in vitro* study [70]. Another *in vitro* analysis displayed the cell growth inhibitory activity was evident by Genistein (2.5-400 μ m) through the up-regulation of two crucial TSGs e.g. p21WAF1 (p21), p16INK4a (p16), and downregulation of two tumor-promoting genes like- c-MYC, and, BMI1, ultimately inhibit cancer progression [71]. Y. Fang et al. concluded that Genistein (40 μ m) inhibits cellular growth via following the activation of DNA-dependent damage response and ATR signaling pathway with activating the BRCA-1 complex, inhibiting cohesion complex and increasing phosphatide, which is distributed among CDK1, CDK2, and CDK3[72]. Recently it was established that Genistein (1000ppm) suppressed the tumor growth by cell cycle regulating via maintaining the expression level of the cyclin D1 protein, leading to G0/G1 phase arrest, which causes cell cycle blockage[39]. Subsequently, T. Rajah et al. summarised that Gen (10-100 μ M) inhibits tumor growth by down-regulating MEK5, pERK5, and NF- κ B proteins [73].

In the case of cell proliferation, low dose Genistein (10 μ m) slightly inhibited cell proliferation by reducing the level of P-STAT3/STAT-5 ratio[55]. In comparison, at a higher dose of 20-40 μ m, it significantly prevents cell proliferation by inducing apoptosis by suppressing Skp2 expression by up-regulating the tumor suppressor genes, i.e. p21, p27, resulting in G2/M phase arrest [74]. Li et al. investigated that Gen (5-20 μ M) inhibits differentiation with the arresting at G2/M phase by decreasing Cdk1, CyclinB1, Cdc25C, c-Jun, and c-Fos level [18]. Gen can also play a role in MDA-MB-231 specific by inhibiting mammosphere formation. A lower dose of Genistein (2 μ m) prevents the mammosphere formation through PI3K/Akt signaling by increasing the PTEN expression[24], while at a higher dose Genistein (40nm-2 μ m) prevents the formation of mammosphere cells and promotes differentiation through the PI3K/Akt and MEK/ERK signaling pathway by reducing the CD44+/CD24-/ESA ratio and increasing E-cadherin mRNA

expression [62]. Finally, Gen (50 μ M) impeded primary tumor formation by down-regulate the chelator neocuproine, Bcl-2/ Bax, at the same time up-regulate the caspase-3 pathway [67].

3.3 Effect of Genistein on T-47D breast cancer cell:

V. Mukund *et al.* summarize that Genistein (50 μ m) lowers angiogenesis by preventing the transactivation of downstream HIF-1 α effectors such as VEGF, resulting in the lessening the expression of hypoxia-inducible factor-1 α in T-47D type of breast cancer cell lines [22]. Cell proliferation efficacy was evident by Genistein (10nm) with apoptosis induction through the mitochondrial-dependent pathway via up-regulating the cytochrome c oxidase activity and downregulating the ATP synthase/cytochrome c oxidase ratio [55]. Genistein at the 1nm-100 μ m inhibits cell proliferation through ERK1/2 mediated signaling by the downregulation of phosphorylated p90RSK [75], while 10 μ M of Gen induces apoptosis and inhibits cell proliferation through degrading proto-oncogene c-fos levels and prohibited Protein 1 (AP-1) and ERK expression [68]. Another *in vitro* study by T. T. Rajah revealed that Gen (10-100 μ M) inhibits Cell proliferation and Tumor growth by downregulating MEK5, pERK5, and NF- κ B proteins [73]. Additionally, Marik et al. demonstrated that while a low concentration of Gen (0.1 M) promotes cancer progression, a high concentration of Gen (20 M) inhibits cell proliferation by reducing ER-messenger RNA transcription and arresting the cell cycle at the G2/M phase [34]. According to A. M. Sotoca et al. summarized that Gene (500nm) inhibits cell growth and induces apoptosis by activating cytoskeleton restructuring that results in interaction among integrins, focalized adhesion kinase and CDC42 that leads to cell cycles arrest in T-47D breast cancer cell line [76]. while according to D. G. Pons et al., Gen (1 μ m) causes a significant decrease in cell viability by increasing Sirt1, TGF β , PRAR γ and decreasing IL-1 β expression in T-47D breast cancer cell [63].

3.4 The Actions of Genistein on HCC1395 breast cancer cell:

W. Y. Lee et al. demonstrated that Genistein (1-200 μ m) inhibits invasion and metastasis through the upregulation of TFPI-2, ATF3, DNMT1, MTCBP-1 genes expression and downregulation of MMP-2, MMP-7, CXCL12 genes, leading to arrest G2/M phase and therefore, reduces cell viability [77].

3.5 The Actions of Genistein on HCC38 breast cancer cell:

M. G. Donovan stated that Genistein (4-10ppm) inhibits cell growth by increasing the BRCA1 protein level and reducing the CpG methylation, consequently decreases the AHR binding at BRCA1 [78].

3.6 The Actions of Genistein on Hs578t breast cancer cell:

According to C. De La Parra et al., Genistein (1-50 μ m) inhibits cell viability and induces apoptosis through the downregulation of mir-155, resulting in upregulation of casein kinase, FOXO3a, p27, PTEN expression and reducing the level of β -catenin [79].

3.7 The Actions of Genistein on DD-762 Cell and Sm-MT C breast cancer cell:

Nakagawa et al. appraised that Gen (7-274.2 μ M) inhibits cell proliferation by up-regulating caspase-3 protein activity [80].

3.8 The Actions of Genistein on BT-474 breast cancer cell

Genistein low concentration (1 μ M) can promote cancer but at high concentration (50 μ M) inhibits the cell division with downregulating Tyrosine kinase, HER2 activation, and MAPK pathway [43]. A similar study approached the same result on the BT-474 cell line. Gen (3.125-25 M) inhibits cell replication and arrests the cell cycle in the G2/M phase, as well as inhibiting the expression of EGFR, HER2, and ER-alpha [81].

3.9 The Actions of Genistein on BT20 breast cancer cell

Cappelletti et al. revealed that Gen (15-30 μ M) inhibits metastasis through down lifting levels of CDKs, Tyrosine kinase, DNA topoisomerase II, and Paracrine stimulation BT20 Cell Line [21].

3.10 The Actions of Genistein on 21PT breast cancer cell

Marik et al. demonstrated that while Gen (0.1 M) stimulates cancer progression, Gen (20 M) inhibits cell proliferation by decreasing ER-messenger RNA expression and arresting the cell cycle at the G2/M phase in the 21PT Cell Line [34].

3.11 The Actions of Genistein on 184-B5/HER breast cancer cell

Katdare et al. showed that Gen (2.5-10 μM) impedes the cell cycle by inducing apoptosis by increasing the P16INK4a gene and decreasing HER-2/neu, Tyrosine kinase [82].

3.12 The Actions of Genistein on MCF-10A, MCF-ANeoT, MCF-T63B breast cancer cell

An early study showed that Gen (1-10 $\mu\text{g/ml}$) obstructs angiogenesis and cell mutation by decreasing the expression of DNA topoisomerases I and II, Ribosomal S6 kinases and Tyrosine kinase [54]. Overview of anti-cancer activities were tabulated in **Table 01**.

Table 01. tabular representation of *in vitro* anti-breast cancer activity Genistein

Target	Pharmacological interaction	Type of study (<i>in vitro</i> and <i>in vivo</i>)	Dose	Molecular mechanism	Molecular target	Ref
		<i>In vitro</i> (MCF-7)	5-100 μM	\uparrow Apoptosis \downarrow Cell proliferation	\downarrow IGF-1R-PI3 K/Akt pathway \downarrow Bcl-2/Bax mRNA	[56]
ER- α	Antagonist	<i>In vitro</i> (MDA-MB-231 and T-47D)	50-100 μM	\downarrow Angiogenesis	\downarrow VEGF \downarrow HIF-1 α expression	[22]
ER- α ER- β	Agonist Antagonist	<i>In vitro</i> (MCF-7, T47D, and MDA-MB-231)	10-10 μM	\downarrow Cell proliferation \downarrow Mitochondrial activity	\downarrow Opa1, Fis1 \uparrow cytochrome c oxidase \downarrow ATP synthase/cytochrome c ratio \downarrow P-STAT3/STAT-3 ratio	[55]
		<i>In vitro</i> (MCF-7)	5-30 μM /20-50 mg/kg	\downarrow Cell growth and proliferation	\downarrow Hedgehog–Gli1 signaling pathway	[57]
		<i>In vitro</i> (MDA-MB-231, and MDA-MB-468)	5-50 μM	\uparrow Apoptosis	\downarrow Endogenous copper ion \uparrow generation of reactive oxygen species (ROS)	[67]
		<i>In vitro</i> (MCF-7 and MDA-MB-231)	60-100 μM	\downarrow Cell viability	\downarrow mRNA expression of DNMT1 \downarrow DNA methylation in tumor suppressor Genes	[65]
		<i>In vitro</i> (MCF-7)	.01-100 μM	\uparrow Apoptosis	\uparrow BAX/Bcl-2 ratio \downarrow Thioredoxin reductase (TrxR), \uparrow Glutathione peroxidase (GPx)	[48]
		<i>In vitro</i> (MDA-MB-231)	5-20 μM	\uparrow Apoptosis \uparrow Cell cycle arrest	\uparrow Arrest G2/M phase \downarrow NF- κ B, Notch-1 pathway \downarrow cyclin B1, \downarrow Bcl-2 and Bcl-xL	[66]
		<i>In vitro</i> (MDA-MB-231 and SKBR3)	20-40 μM	\downarrow Cell proliferation \uparrow Apoptosis \downarrow Metastasis	\downarrow Skp2 expression \uparrow Arrest G2/M phase \uparrow p21, p27	[74]
		<i>In vitro</i> (MCF-7)	75-200 μM	\downarrow Cell growth	\uparrow miR-23b \uparrow Target PAK2 gene	[83]

		<i>In vitro</i> (MDA-MB-231)	2.5-400µm/250mg/kg	↓Cell growth	↑ p21WAF1 (p21), ↑p16INK4a (p16) ↓ BMI1, c-MYC	[71]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> (MCF-7 and MDA-MB-231)	5-20µm	↑Apoptosis ↑DNA damage	↑Arrest G2/M ↑ATM, Bax, P ⁷³ ↓Bcl-2, ↓ Bcl-2/Bax rate, mutant P ⁵³	[49]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> (MCF-7, MDA-MB-435, and Hs578t)	1-50µm	↓Cell viability ↑Apoptosis	↓ miR-155 ↓β-catenin ↑Casein kinase, ↑ FOXO3a, p27, PTEN, CK1α	[79]
		<i>In vitro</i> (MDA-MB-231)	40µm	↓Cell growth	↑ phosphopeptide ↓Cohesin complex ↑DNA damage response pathway ↑ BRCA1 ↑ ATR signaling pathway	[72]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> (MCF-7 and MDA-MB-231)	40 - 2µm	↓ Mammosphere formation	↑PTEN expression ↓PI3K/Akt signaling	[24]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> (MCF-7, T47D and MDA-MB-231)	1µm	↓Cell viability ↓ Cell proliferation	↑ROS, pS2, Sirt1, COX2 ↓ IL-1β, TFGβ, PPARγ	[63]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> (T47D)	500nm	↓Cell growth ↑Apoptosis	↑Cytoskeleton remodeling ↑integrins, focal adhesion kinase, CDC42 ↑Arrest cell cycle	[76]
ER-α	Antagonist	<i>In vitro</i> (T47D)	1-100µm	↓Cell proliferation	↓ ERK1/2, p90RSK	[75]
ER-α	Antagonist	<i>In vitro</i> (MCF7, UACC3199, and HCC38)	4-10ppm	↓Cell growth	↑Activate BARCA-1 ↓ CpG methylation ↓AHR activity	[78]
ER-β	Agonist	<i>In vitro</i> (MCF-7 and MDA-MB-231)	10 ⁻⁶ mol/l- 10 ⁻⁴ mol/l /100-1000ppm	↓Cell proliferation	↓Cyclin D1 ↑Arrest G0/G1 phase	[39]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> (MCF-7 and MDA-MB-231)	40nm-2µm	↓Mammosphere formation ↑Differentiation	↓CD44+/CD24-/ESA ratio ↑PI3K/Akt, MEK/ERK signaling ↑E-cadherin expression	[62]
ER-α	Antagonist	<i>In vitro</i> (MCF-7 and UACC-3199)	0.5-20µm	↓Cell growth	↑ BRCA-1 ↓CpG methylation, ↓Cyclin D1, ↓ DNMT-1 ↑AhR ↑CYP1A1	[84]

ER- α	Antagonist	<i>In vitro</i> (HCC1395)	1-200 μ m	↓Cell viability ↓Invasion ↓Metastasis	↑TFPI-2, ATF3, DNMT1, MTCBP-1 ↓MMP-2, ↓MMP-7, CXCL12 ↑Arrest G2/M phase	[77]
ER- β	Agonist	<i>In vitro</i> (SUM1315MO2)	1-100 μ m	↓Cell proliferation	↑ER- β expression ↑Restore BRCA1 function	[85]
ER- α	Antagonist	<i>In vitro</i> (MCF-7)	50-200 μ m	↓Cell growth ↑Apoptosis	↓Bcl-2, ↑Bax ↓Cyclin D1 ↓Bcl-2/Bax ratio	[37]
ER-α	Agonist & Antagonist	<i>In vitro</i> MCF-7	1-25 μ m	↑Apoptosis ↓Cell proliferation	↑CDKN1A, TNF- α p53 responsive gene ↓CCNG1 and GADD45A ↓BCL-2, Bcl-3 and NF-kappa B and TNFR ↓NCOA2 and NCOA3	[50]
ER-α	Antagonist	<i>In vitro</i> MDA-MB-435	750 μ g /g	↑Apoptosis ↓Metastasis	↓Tyrosine phosphorylation ↑Matrix degrading enzymes	[86]
ER-α	Antagonist	<i>In vitro</i> MCF-7, ZR-75.1, T47-D, MDA-MB 468, MDA-MB 231 and HBL	1-10 μ m	↑Apoptosis ↓Cell proliferation	↓c-fos levels, Protein-1 (AP-1) activity, (ERK) activity ↑Arrest at G2 phase	[68]
ER-α	Antagonist	<i>In vitro</i> MCF-7 and MDA-MB-231	50 μ m	↑Apoptosis ↓Cell cycle kinetics	↑Poly-(ADP-ribose)-polymerase, p53 ↓Anti-apoptotic protein	[26]
ER-α ER-β	Antagonist	<i>In vitro</i> MCF-7 MDA-MB-231	50 μ m	↑Apoptosis ↓Cell division ↓Angiogenesis	↓EROD, TPA ↓Cyclooxygenase, Topoisomerase II, Tyrosine Kinase.	[53]
ER-α ER-β	Antagonist	<i>In vitro</i> MDA-MB-468	25-100 μ m	↓Cell cycle kinetics ↑Apoptosis ↓Cell proliferation	↑Arrest at G2/M phase ↑Nuclear membrane breakdown during G2/M transition ↓DNA synthesis	[87]
ER-β	Antagonist	<i>In vitro</i> MCF-7	0.001-10 μ m	↑Cells underwent apoptosis	↑Plasma membrane breakdown ↑Nuclear membrane breakdown ↑pS2 expression	[27]
ER-β	Antagonist	<i>In vitro</i> MCF-7 and MDA-MB-231	10-100 μ m	↑Apoptosis ↓Cell division	↓PTK, Akt, FAK, ErbB-2, and Bcl-2 ↓Topoisomerase II, Tyrosine kinase ↓Osteoclast activity	[46] , [69] , [88] – [95]
ER-α ER-β	Agonist Antagonist	<i>In vitro</i> BT-474	1-50 μ m	↑Apoptosis ↓Cell division	↓Tyrosine kinase, HER2 activation, MAPK pathway	[43]
ER-α ER-β	Antagonist	<i>In vitro</i> MDA-MB-231	0.5-15 μ m	↓Cell cycle kinetics ↑Apoptosis	↑Anti-growth signals protein ↑Connexin phosphorylation blocks the homeostatic regulators	[96]

ER-α	Antagonist	<i>In vitro</i> MDA-MB-231, BT20, T47D, and ZR75.1	15-30 μ m	↑Cell underwent apoptosis ↓ Migration	↑ Arrest at G2/M phase ↓ Tyrosine kinase ↓ DNA topoisomerase II ↓ Paracrine stimulation	[21]
ER-α ER-β	Agonist & Antagonist	<i>In vitro</i> MDA-MB-231 & T47D	10-100 μ m	↓Decrease cell proliferation ↓ Tumor growth	↓ MEK5, pERK5, NF- κ B proteins	[73]
ER-α ER-β	Antagonist	<i>In vitro</i> AS-4, NEO & BG-1	25-150 μ m	↑ Inhibit cell proliferation ↑Induce apoptosis	↓ Cytotoxic effect in AS4, Tyrosine kinase, Topoisomerase II ↑ Ubiquitin E3 ligase	[97]
ER-α	Antagonist	<i>In vitro</i> MDA-MB-231	5-20 μ m	↑ Trigger apoptosis ↓ Invasion	↓ MEK5, ERK5, NF- κ B/p65 ↓ Bcl-2/ Bax	[20]
ER-α	Antagonist	<i>In vitro</i> MDA-MB-231	5-20 μ m	↑ Cell depletion ↓Differentiation	↑ Trigger G2/M cell cycle arrest ↓ Cdk1, CyclinB1 and Cdc25C ↓ c-Jun and c-Fos	[18]
ER-α	Antagonist	<i>In vitro</i> MCF-7	50 μ m	↓ Tumor growth ↑ Induces apoptosis	↑ Ca ²⁺ dependent proapoptotic proteases ↑ μ -calpain, and caspase-12	[59]
ER-β	Agonist & Antagonist	<i>In vitro</i> MDA-MB-231	20-80 μ m & 750 μ g/g	↓ Cell growth ↑ Tumor formation in vivo	↑ Cell cycle blocked at G2/M	[98]
ER-α	Antagonist	<i>In vitro</i> MCF-7 and MDA-MB- 231	100 μ m	↑ Induced apoptosis ↓ Cell proliferation	↑ Arrest at G2/M phase ↓ Paclitaxel-induce tubulin Polymerization ↓ Bcl-2 phosphorylation ↓Cyclin B1 and CDC2 kinase	[60]
ER-α	Antagonist	<i>In vitro</i> BT-474	3.125-25 μ m	↑ Induction of apoptosis ↓ Duplication	↓ HER2 and ER-alpha ↑Cell cycle arrest at S and G2/M ↓ Expression of surviving, EGFR	[81]
ER-α ER-β	Antagonist	<i>In vitro</i> MDA-MB- 231, MCF-7, and MCF- 12A	35-100 μ m	↓ Cell invasiveness ↓ Cell cycle ↑ Phagocytosis	↓ Protein tyrosine kinase pathway ↑ Arrest at G2/M phase ↓ MMP genes ↓ Glucose uptake	[69]
ER-α	Antagonist	<i>In vitro</i> MCF-7	100 μ m	↑ Induce Apoptosis ↓ Cell Growth	↑ Caspase 7 and poly (ADP ribose) polymerase ↑ Apoptosis signalling kinase 1 ↑ p38 MPK	[51]
ER-α	Antagonist	<i>In vitro</i> MDA-MB-231	5-50 μ m	↑ Induce Apoptosis ↓ Cell cycle	↑Apoptosis related genes ↓ Akt and ↓ NF- κ B ↓ EGF	[70]
ER-α	Antagonist	<i>In vitro</i> MCF-7	50-200 μ m	↓ Cell invasion ↑ Apoptosis ↓ Cancer progression	↓Proteasomal chymotrypsin-like activity ↓CDKs inhibit by p27Kip1 ↑ I κ B- α level	[64]
ER-α	Antagonist	<i>In vitro</i> MCF-7	50-100 μ m	↓ Cell division ↓ Cancer progression	↓SRF, RAG-1, DOC 2 ↑ Arrest at G2/M phase ↑ Heat shock protein 105 (HSP) mRNA	[61]

ER-α	Agonist Antagonist	<i>In vitro</i> MCF-7 and MDA-MB-468	3.125-12.5 μ m	↓ primary tumor ↓ Metastasis	↑ GSTP1 gene ↑ RAR β 2 gene	[52]
ER-α	Antagonist	<i>In vitro</i> MCF-7	1-100 μ m	↑ Apoptosis, Cell cycle arrest	↓ CDKs, Akt activity ↓ Telomere length	[58]
ER-α	Agonist Antagonist	<i>In vitro</i> MCF-7, 21PT, and T47D	0.1-20 μ m	↓ Cell proliferation ↓ Cancer progression	↓ ER- α messenger RNA expression ↑ Er- β ↑ Cell cycle arrest at the G2-M phase	[34]
	Antagonist	<i>In vitro</i> MDA-MB-231, and MDAMB-468	50 μ m	↓ Primary tumor ↑ Apoptosis	↓ Chelator neocuproine ↓ Bcl-2 ↑ Bax and caspase-3	[67]
ER-α	Antagonist	<i>In vitro</i> 184-B5/HER	2.5-10 μ m	↑ Cellular apoptosis ↓ Cell cycle regulators	↑ P16 ^{INK4a} gene ↓ HER-2/neu ↓ Tyrosine kinase ↑ Arrest at S + G2/M phase	[82]
ER-α	Antagonist	<i>In vitro</i> MCF-10A1, MCF-ANeoT, MCF-T63B, MCF-7, MDA-231, MDA-435	1-10 μ g/ml	↓ Angiogenesis ↓ Cell proliferation ↓ Cell mutation	↓ Dysplastic and malignant ↓ Tyrosine kinase ↓ DNA topoisomerases I and II ↓ Ribosomal S6 kinases	[54]

4. Mechanism of synergistic activities of Genistein with Chemotherapeutics Resistance Breast Cancer Treatment breast cancer drugs

Except for the strong anti-cancer activity alone, Genistein possesses the synergistic property with many other anti-cancer drugs, which assist it in overcoming the chemopreventive resistance mechanism in breast cancer treatment. The synergistic activity of Genistein can be carried by many anti-cancer drugs such as Doxorubicin, Trastuzumab, Tamoxifen, Trichostatin A, Cisplatin, Capsaicin, Paclitaxel, and vincristine. Tumor cells frequently appear to have a decreased capacity for cytoprotective drug accumulation as a result of the drugs efflux protein (MDR) serving as the pharmacological background for multidrug resistance, these efflux proteins, namely as the BCRP, MRP 1, and P-glycoprotein [99] [100] and the changes in beta-tubulin [100]; It is significant to mention that the multidrug resistance protein (MRP) family has nine members (MRP1–9) [101].

Research study conducted by C.H.M. and colleagues has shown that Genistein inhibited drug accumulation in various types of multiresistant tumor cell lines mediated by non-P-glycoprotein (HT1080/DR4, HL60/ADR, MCF7/Mitox, GLC4/ADR, and SW-1583/2R120) [102]. According

to another research study, genistein, in the form of phytoestrogens, can effectively suppress the breast cancer resistance protein, also known as ABCG2, which is responsible for multidrug resistance activity [103]. Moreover, others research studies have analyzed that the Genistein's ability to autophosphorylate protein tyrosine kinases (PTK), particularly EGF-R, by contesting ATP than other the protein substratum. The genistein antiproliferative effect was also attributable to interfering with the mitogen-activated tyrosine kinase cascade [46].

4.1 Synergistic Properties of Genistein in the treatment of breast cancer

4.1.1 Synergistic Properties of Genistein with Doxorubicin in MCF-7/Adr cell: Doxorubicin is an antibiotic that exhibits no inhibitory effects on Adriamycin resistance breast cancer cell. But the combination of Genistein at 30 μ mol/l and increasing doses of Doxorubicin possess synergistic effects on MCF-7/Adr cell. Genistein enhances the cytotoxic effect of Doxorubicin and decreases the MCF-7/Adr breast cancer cell's chemoresistance. In addition, Genistein and Doxorubicin induce apoptosis synergistically by decreased expression of Her2/new mRNA and c-erbB2, resulting in cell cycle arrests in MCF-7/Adr breast cancer cell in G2/M phase [25]. Another study by X. Yang et al. reported that Gen (50 μ m) with a combination of Doxorubicin slightly induces apoptosis by destroying the plasma membrane of cells and increase Poly (ADP-ribose) polymerase cleavage in MDA-MB-231 and MCF-7 breast cancer cells [26].

4.1.2 Synergistic effect of Genistein with Trastuzumab: Genistein and Trastuzumab synergistically develop the antitumor activity in BT-474 breast cancer cells. C. Lattrich et al. stated that the combination of Genistein (10 μ mol/l) and Trastuzumab (1/10 μ g/ml) enhances the growth-inhibitory effect and reduces viable cell numbers by increasing the ER- β 2 expression, which causes an antiestrogenic effect, leading to reducing cell proliferation in ER- α/β -positive, and HER2-overexpressing BT-474 breast cancer cell. Furthermore, both Genistein and trastuzumab reduce cyclin A2 mRNA expression, c-fos, HER2, and induce cyclin D1 expression which suppresses the cell proliferation of BT-474 breast cancer cell [104].

4.1.3 Synergistic Properties of Genistein with Tamoxifen: Tamoxifen is a well-established medicine for treating breast cancer, although its efficacy has been hampered by the development of gemcitabine resistance. In this regard, Gen can improve the efficacy of Tamoxifen. Y. Li et al. described that Genistein enhanced the anti-cancer capacity of Tamoxifen at the dose of 25 μ m

through the reactivation of ER- α through epigenetic pathways i.g., histone modification, resulting in the reduction of HDAC1 & DNMT1 expression both *in vitro* (ER- α negative MDA-MB-231 breast cancer cell) & *in vivo* leading to inhibits the cell growth and cell viability [28]. On the other hand, D. G. Pons et al., concluded that 1 μ M Genistein and 10 μ M Tamoxifen declines the ROS production in T47D, and MCF-7 breast cancer cell and up-regulate the autophagic vacuole formation & PARP protein level and also reduce cell viability, resulting in autophagic cell death only in T47D breast cancer cell [105]. Another early study reported that Tamoxifen, with Gen (1-10 μ g/ml), impede angiogenesis and cell mutation by downregulating DNA topoisomerases I and II, Ribosomal S6 kinases, tyrosine kinase, and Cell cycle regulators [54]. Gen also shows a prohibitory effect with Tamoxifen which induces apoptosis by destroying nuclear membrane [27], arrest cell cycle by decreasing expression of HER2 in a dose-dependent manner [81].

4.1.4 Synergistic effect of Genistein with Trichostatin A: Genistein and Trichostatin A act synergistically to inhibit PGR (progesterone receptors) expression (elaboration required), resulting in a significant change in cell growth in ER-positive and ER-negative MCF-7 cells. Breast cancer cell line MDA-MB-231. According to Y. Li et al., the association of Genistein (25 μ m) and Trichostatin A (100ng/ml) synergistically induces platinum synergistically decreases the ROS production and enhances antioxidant enzyme i.e the Mn-SOD (Manganese-Superoxide dismutase) & catalase, in MCF-7 & T47D breast cancer cell. Furthermore, Genistein and Cisplatin also synergistically arrest the cell cycle at S phase and cause a drop in subG0/G1 phase, resulting in MCF-7 cells that are apoptotic at (25-50mol/L) concentrations [105].

4.1.5 Synergistic effect of Genistein with capsaicin: The combination of genistein and capsaicin exerted anti-inflammatory and anticarcinogenic effects in MCF-7 cells as well as *in vivo* 48 weeks female Sprague-Dawley rats by modulating the mitogen-activated protein kinase (AMPK) and COX-2, as well as possibly other mitogen-activated protein kinases [23].

4.1.6 Synergistic effect of Genistein with Paclitaxel and vincristine: Paclitaxel and vincristine both are chemotherapy drugs, also called plant alkaloids. Together with Gen (100 μ m), they can suppress cell growth and viability through inhibiting CDC2 and Cyclin B1 kinase, and inhibit microtubule polymerization in human MDA-MB-231 and MCF-7 and breast cancer cell line, ultimately cell death by inducing apoptosis via decreasing Bcl-2 phosphorylation without

changing p21, p53, and Bax protein expression [60]. The **Table 02** contains a summary of the effects of phytoestrogens in combination with anti-cancer therapies that have been previously described.

Table 02. Summary of the described effects of phytoestrogens in combination with anti-cancer therapies.

Treatment option	Study type	Effective mechanism	Ref
Doxorubicin	<i>In vitro</i> (MCF-7/Adr)	↓Chemoresistance of tumor cells ↑Cell cycle Arrest G ₂ /M phase ↑Apoptosis ↓Her2/neu mRNA expression ↓c-erbB2 expression	[25]
	<i>In vitro</i> MCF-7 and MDA-MB-231	↑ Destroyed the plasma membrane of cells ↑ Apoptosis in most of the individual MCF-7 cells ↑ Poly (ADP-ribose) polymerase cleavage	[26]
Trastuzumab	<i>In vitro</i> (SK-BR-3, BT-474, and MDA-MB-231)	↓Cell proliferation ↑ Cell cycle arrest ↓c-fos, ↓ HER2	[104]
Tamoxifen (TAM)	<i>In vitro and vivo</i> (MDA-MB-231) female immunodeficiency nude mice)	↓Cell growth and Cell viability ↓HDAC1, ↓ DNMT1	[28]
	<i>In vitro</i> MCF-7	↑ Destroyed the plasma membrane of cells ↑ Nuclear membrane breakdown ↑ pS2 expression	[27]
	<i>In vitro</i> BT-474	↓ Expression of HER2 and ER- α ↓ Expression of factor and EGFR	[81]
	<i>In vitro</i> Dysplastic,malignant cells	↓ Cell growth, proliferation	[54]
Trichostatin A (TSA)	<i>In vitro</i> (MCF-7, MDA-MB-231 and MDA-MB-157)	↑ER- α reactivation, ↑Histone remodeling ↓HDACs, DNMTs, PGR expression	[28]
Cisplatin (CDDP)	<i>In vitro</i> (MCF-7 and T47D)	↓ROS production ↑MnSOD, Catalase activity ↑Arrest S and G ₀ /G ₁ phase	[105]
Tamoxifen (TAM)	<i>In vitro</i> (MCF-7 and T47D)	↓ROS, LC3-II/LC3-I ratio ↓cell viability ↑PARP	[105]
Ornithine decarboxylase	<i>In vitro</i> MCF-7 MDA-MB-231	↑ Putrescine, Spermidine and Spermine ↓ 12-Otetradecanoylphorbol- 13-acetate (TPA) ↓ Cyclooxygenase (COX)	[53]
paclitaxel and vincristine	<i>In vitro</i> MCF-7 and MDA-MB-231	↓ Bcl-2 phosphorylation ↓Cyclin B1 and CDC2 kinase ↓ Cell viability ↓ Microtubule polymerization	[60]
Lycopene	<i>In vivo</i>	↓ Serum MDA	[106]

	<i>Female Wistar rats</i>	↓ 8-isoprostane and 8-OhdG levels ↓ Bcl-2, caspase 3, and caspase 9 ↓ Expression of Bax	
--	---------------------------	---	--

5. Clinical trials

Human clinical trials have confirmed the in vitro and in vivo research findings; in some cases, when consumed at a consistent dose, pure Genistein had no estrogenic effect on breast tissue [29], [30], although others dietary soy supplementation had proestrogenic properties on breast tissue [31]–[33]. Several secondary endpoints were evaluated in a recently published clinical study to determine whether purified Genistein has an effect on the endometrial thickness, vaginal cytology, and breast density (Table 3) [29][93],[94]. Following the implementation of these safety measures, it was possible to identify the potential estrogenic effects of 54 mg/day of purified Genistein as indicators of breast cancer risk in the research participants.

Indeed, while the placebo group maintained a constant endometrial thickness, the Genistein group demonstrated a time-dependent reduction that reached statistical significance during the 36-month follow-up (approximately 12% reduction, $P < 0.01$). Moreover, levels of gene expression of BRCA-1, and 2, breast tumor suppressor genes [109], [110] have been preserved for the three years in the genistein-administered group, while levels of both BRCA-1, and 2 have decreased in the placebo group (Table 3) [29][30]. Genistein also significantly reduced sister chromatid exchanges, implying that it may prevent genotoxicity and subsequent mutagenesis (**Table 03**) [30].

In this regard, based on the use of Genistein in breast cancer, 2 clinical trials, a phase II study entitled “Gemcitabine Hydrochloride and Genistein in Treating Women with Stage IV Breast Cancer,” and a phase I study entitled “Genistein in Preventing Breast or Endometrial Cancer in Healthy Postmenopausal Women,” have been completed, but the results are not yet published ([NCT00244933](#), [NCT00099008](#)). Effects of Genistein on markers of cancer risk observed from human clinical studies was summarized in

Table 03. Table 04: Effects of Genistein on markers of cancer risk observed from human clinical studies (post-menopausal women)

Subjects (n)	Number/intervention (mg/day)	Study length (months)	Results	Ref
57	54g/day genistein (n = 30) or placebo (n = 27)	12	Protect genomic damage	[30]
90	HRT (n = 30; 1 mg 17 beta-estradiol combined with 0.5 mg norethisterone acetate), Gen (n = 30; 54 mg), or placebo (n = 30)	12	↑ BMD in the femoral neck and lumbar spine; ↑ serum B-ALP and BGP;	[108]
389	Placebo (n = 191; 1000 mg calcium and 800 mg vitamin D) or Gen (n = 198; 54 mg Gen, 1000 mg calcium, and 800 mg vitamin D)	24	↑ BMD at the lumbar spine and the femoral neck; ↑ B-ALP and IGF-1;	[107]
220	Placebo (n = 111) or isoflavone (n = 109; ~50 mg isoflavones)	24	↑ breast area ↓ breast density; neither was significant when compared to placebo	[111]
34	Placebo (n = 17) or soy (n = 17; 100 mg isoflavone, ~76 mg aglycones)	12	↓ Breast area no change in dense areas or percent breast density No significant changes in any hormone levels	[112], [113]
84	Placebo (n = 23) or soy (n = 28; 60 mg, ~45 mg isoflavones)	14 days	↓ Nipple aspirate levels of apolipoprotein D; ↑ pS2	[32][31]
389	54 mg of genistein aglycone daily (n = 71) or placebo (n = 67)	36	unchanged breast density, BRCA1 and BRCA2; ↓ sister chromatid exchange. ↓ pyridinoline, NF-B receptor activator ↑ alkaline phosphatase, IGF-I, and osteoprotegerin	[29]

6. Safety measurement of Genistein and future prediction

ADMET (absorption, distribution, metabolism, elimination, and toxicity) testing evaluates prospective pharmacological substances, identifying both those with promise and those with significant limitations. ADME/Tox profile of Genistein was predicted using online accessible SwissADME [114](drug likeness properties), pkCSM [115](absorption, distribution, excretion) [115]and server admetSAR [116] (metabolism, toxicity) in silico tools and tabulated in **Table 04**.

With a good bioavailability score of 0.55, Genistein maintained drug-like characteristics without any violating of Lipinski, Ghose Veber, Muegge, and Egan rules. In drug development, absorption is an important parameter. Our *in silico* predicted result showed that Genistein is highly soluble in water (-3.376) and absorbed by human intestine, with permeable to Caco-2 cell. After absorption, consequentially comes distribution, another principal descriptor for drug development, which depends numerous factors i.e., steady-state volume of distribution (VD_{ss}), blood-brain barrier (BBB) and central nervous system(CNS) permeability, binding to plasma protein, and many more [117]. Our analysis suggested that Gen has the potentiality to have a poor VD_{ss} value, but is able to cross the BBB and central CNS properly. Human cytochrome P450 (CYP) isoforms are involved in drug metabolism in the liver. Its inhibition could lead to drug toxicity, drug-drug interactions, and other adverse effects. The predicted metabolic result reported that Genistein is the inhibitor of CYP1A2, CYP2C19, CYP2C9, and CYP3A4 but a non-inhibitor of CYP2D6. Excretion property based on total renal clearance parameter was predicted, where total clearance (logCL_{tot}) of Genistein was 0.151 ml/min/kg. The toxicity profile of Genistein has been predicted based on eye corrosion, hepa-toxicity, Skin sanitization, OATP2B1 inhibitor, OATP1B3 inhibitor, OCT2 inhibitor, carcinogenicity, AMES toxicity, hERG I inhibitors, and BSEP inhibitors. Results outlined that Genistein showed toxicity only in the liver and OATP2B1 inhibitor, whereas positive to eye corrosion, hepatotoxicity, AMES toxicity, hERG potassium channel inhibition, and carcinogenicity. Anti-cancer activities were determined based on P-glycoprotein and aromatase enzyme inhibition and estrogen receptors modulating affinity. Our analysis revealed that Genistein has a potent affinity to estrogen receptors, aromatase enzyme, but its non- inhibitor P-glycoprotein.

To achieve more information for better bioavailability and drug-likeness of Genistein, detailed information is summarized in **Fig 02**. The results of the bioavailability radar have been depicted by the lipophilicity: XLOGP3 between -0.7 and +5.0, size: MW between 150 and 500 g/mol, polarity: TPSA between 20 and 130 Å², solubility: log S not higher than 6, saturation: fraction of carbons in the sp³ hybridization not less than 0.25 and flexibility: no more than 9 rotatable bonds with the colored zone defined the desired physicochemical space for good oral bioavailability indicating that they possess good drug-likeness properties. After the validation of bioavailability, potent drug likeness properties in cancer research models, it can be estimated that Genistein can

be used as an effective anti-cancer natural agent with its synergistic use with other chemotherapeutics for the treatment of breast cancer.

Table 04. *In silico* pharmacokinetics study of Genistein

Pharmacokinetics parameter		Predicted remarks	Unit
Drug likeness	Lipinski	Yes	Yes/No
	Ghose	Yes	Yes/No
	Egan	Yes	Yes/No
	Veber	Yes	Yes/No
	Muegge	Yes	Yes/No
	Bioavailability Score	0.55	N/A
Absorption	Water solubility	-3.595	log mol/L
	Caco2 permeability	0.9	log Papp in 10 ⁻⁶ cm/s
	Human intestinal absorption	93.387	% Absorbed
	Skin Permeability	-2.735	log Kp
	P-glycoprotein substrate	Yes	Yes/No
	P-glycoprotein inhibitor	No	Yes/No
Distribution	VDss (human)	0.094	log L/kg
	BBB permeability	-0.71	logBB
	CNS permeability	-2.048	logPS
Metabolism	CYP1A2 inhibitor	Yes	Yes/No
	CYP2C19 inhibitor	Yes	Yes/No
	CYP2C9 inhibitor	Yes	Yes/No
	CYP2D6 inhibitor	No	Yes/No
	CYP3A4 inhibitor	Yes	Yes/No
Excretion	Total clearance	0.151	log ml/min/kg
Toxicity	Eye corrosion	No	Yes/No
	Hepa-toxicity	Yes	Yes/No
	Acute oral toxicity	1.842	Kg/mol
	OATP2B1 inhibitor	Yes	Yes/No
	OATP1B3 inhibitor	No	Yes/No
	OCT2 inhibitor	No	Yes/No
	AMES toxicity	No	Yes/No
	hERG I inhibitor	No	Yes/No
	Carcinogenicity	No	Yes/No
BSEP inhibitors	No	Yes/No	
Anti-cancer affects	P-GP Inhibitor	No	Yes/No
	Aromatase	Yes	Yes/No
	ER binding	Yes	Yes/No




Figure 02. Radar graph of Genistein

7. Concluding remarks:

Nowadays, breast cancer is the major leading cause of death due to have diverse pernicious side effects of proposed synthetic drugs, multi drug resistance issues and many more. But natural food products are given more attention to the people in combating diseases such as- cancer insurgencies, mainly, breast cancer. For this reason, this current investigation was conducted to explore the better pharmacological activity of natural phytochemicals namely-Genistein that offer therapeutic activities in tissues of breast cancer patients, cell specific anti-breast cancer mechanism, and most importantly, possess potential synergistic activity with multidrug-resistant tumors. Here, the safety assessment of the Genistein conducted by the online-based server and predict their pharmacokinetics profiling as a safe therapeutic compound but more wet-lab based research is badly needed to discover the cell-specific breast cancer controlling, toxicological mechanism and their pharmacokinetics properties.

Author's contribution:

M.S., P.B., and D.D. have equally contributed for the conceptualization and study designing; M.S., P.B., M.A.A., D.D., H.S., S.A., M.S.K., M.A.H., P.P., and N.I. equally participated in data collection, writing, and draft preparation; M.A.R., and A.A.M. review and editing; Visualization

and supervision by M.A.R., A.A.M., and B.K.; The Project Funded by B.K. All authors have revised and agreed to the published version of the manuscript.

Acknowledgement

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2020R111A2066868), the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2020R1A5A2019413), a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number : HF20C0116), and a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number : HF20C0038).

References

- [1] R. L. Siegel, K. D. Miller, and A. Jemal, "Cancer statistics, 2020," *CA. Cancer J. Clin.*, vol. 70, no. 1, pp. 7–30, 2020, doi: 10.3322/caac.21590.
- [2] Y. S. Sun *et al.*, "Risk factors and preventions of breast cancer," *Int. J. Biol. Sci.*, vol. 13, no. 11, pp. 1387–1397, 2017, doi: 10.7150/ijbs.21635.
- [3] W. Yue *et al.*, "Effects of estrogen on breast cancer development: Role of estrogen receptor independent mechanisms," *Int. J. Cancer*, vol. 127, no. 8, pp. 1748–1757, 2010, doi: 10.1002/ijc.25207.
- [4] E. Bryer and D. Henry, "Chemotherapy-induced anemia: etiology, pathophysiology, and implications for contemporary practice," *Int. J. Clin. Transfus. Med.*, vol. Volume 6, pp. 21–31, 2018, doi: 10.2147/ijctm.s187569.
- [5] R. R. Love, H. Leventhal, D. V. Easterling, and D. R. Nerenz, "Side effects and emotional distress during cancer chemotherapy," *Cancer*, vol. 63, no. 3, pp. 604–612, 1989, doi: 10.1002/1097-0142(19890201)63:3<604::AID-CNCR2820630334>3.0.CO;2-2.
- [6] S. Crawford, "Is it time for a new paradigm for systemic cancer treatment? Lessons from a century of cancer chemotherapy," *Front. Pharmacol.*, vol. 4 JUN, 2013, doi: 10.3389/fphar.2013.00068.
- [7] P. Fumoleau *et al.*, "Multicentre, phase II study evaluating capecitabine monotherapy in patients with anthracycline- and taxane-pretreated metastatic breast cancer," *Eur. J. Cancer*, vol. 40, no. 4, pp. 536–542, 2004, doi: 10.1016/j.ejca.2003.11.007.
- [8] S. Mitra and R. Dash, "Natural Products for the Management and Prevention of Breast Cancer," *Evidence-based Complementary and Alternative Medicine*, vol. 2018. 2018, doi: 10.1155/2018/8324696.
- [9] N. S. Wind and I. Holen, "Multidrug Resistance in Breast Cancer: From In Vitro Models to Clinical Studies," *Int. J. Breast Cancer*, vol. 2011, pp. 1–12, 2011, doi: 10.4061/2011/967419.

- [10] A. Luque-Bolivar, E. Pérez-Mora, V. E. Villegas, and M. Rondón-Lagos, “Resistance and overcoming resistance in breast cancer,” *Breast Cancer Targets Ther.*, vol. 12, pp. 211–229, 2020, doi: 10.2147/BCTT.S270799.
- [11] I. Bilal, A. Chowdhury, J. Davidson, and S. Whitehead, “Phytoestrogens and prevention of breast cancer: The contentious debate,” *World Journal of Clinical Oncology*, vol. 5, no. 4, pp. 705–712, 2014, doi: 10.5306/wjco.v5.i4.705.
- [12] M. Torrens-Mas and P. Roca, “Phytoestrogens for cancer prevention and treatment,” *Biology*, vol. 9, no. 12, pp. 1–19, 2020, doi: 10.3390/biology9120427.
- [13] A. H. Wu, M. C. Yu, C. C. Tseng, and M. C. Pike, “Epidemiology of soy exposures and breast cancer risk,” *British Journal of Cancer*, vol. 98, no. 1, pp. 9–14, 2008, doi: 10.1038/sj.bjc.6604145.
- [14] S. Bhagwat, D. B. Haytowitz, and J. M. Holden, “USDA Database for the Isoflavone Content of Selected Foods,” *U.S. Dep. Agriculture*, pp. 1–156, 2011.
- [15] K. Sak, “Current epidemiological knowledge about the role of flavonoids in prostate carcinogenesis,” *Experimental Oncology*, vol. 39, no. 2, pp. 98–105, 2017, doi: 10.31768/2312-8852.2017.39(2):98-105.
- [16] G. G. J. M. Kuiper *et al.*, “Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor β ,” *Endocrinology*, vol. 139, no. 10, pp. 4252–4263, 1998, doi: 10.1210/endo.139.10.6216.
- [17] K. Morito *et al.*, “Interaction of phytoestrogens with estrogen receptors alpha and beta (II).,” *Biol. Pharm. Bull.*, vol. 25, no. 1, pp. 48–52, 2002, doi: 10.1248/bpb.25.48.
- [18] Z. Li *et al.*, “Genistein induces G2/M cell cycle arrest via stable activation of ERK1/2 pathway in MDA-MB-231 breast cancer cells,” *Cell Biol. Toxicol.*, vol. 24, no. 5, pp. 401–409, 2008, doi: 10.1007/s10565-008-9054-1.
- [19] Y. Li, S. Upadhyay, M. Bhuiyan, and F. H. Sarkar, “Induction of apoptosis in breast cancer cells MDA-MB-231 by genistein,” *Oncogene*, vol. 18, no. 20, pp. 3166–3172, 1999, doi: 10.1038/sj.onc.1202650.

- [20] Z. Li *et al.*, “Genistein induces cell apoptosis in MDA-MB-231 breast cancer cells via the mitogen-activated protein kinase pathway,” *Toxicol. Vitro.*, vol. 22, no. 7, pp. 1749–1753, 2008, doi: 10.1016/j.tiv.2008.08.001.
- [21] V. Cappelletti, L. Fioravanti, P. Miodini, and G. Di Fronzo, “Genistein blocks breast cancer cells in the G2M phase of the cell cycle,” *J. Cell. Biochem.*, vol. 79, no. 4, pp. 594–600, 2000, doi: 10.1002/1097-4644(20001215)79:4<594::AID-JCB80>3.0.CO;2-4.
- [22] V. Mukund, M. S. Saddala, B. Farran, M. Mannavarapu, A. Alam, and G. P. Nagaraju, “Molecular docking studies of angiogenesis target protein HIF-1 α and genistein in breast cancer,” *Gene*, vol. 701, no. January, pp. 169–172, 2019, doi: 10.1016/j.gene.2019.03.062.
- [23] J. T. Hwang, Y. K. Lee, J. I. Shin, and O. J. Park, “Anti-inflammatory and anticarcinogenic effect of genistein alone or in combination with capsaicin in TPA-treated rat mammary glands or mammary cancer cell line,” in *Annals of the New York Academy of Sciences*, 2009, vol. 1171, pp. 415–420, doi: 10.1111/j.1749-6632.2009.04696.x.
- [24] M. T. E. Montales *et al.*, “Repression of mammosphere formation of human breast cancer cells by soy isoflavone genistein and blueberry polyphenolic acids suggests diet-mediated targeting of cancer stem-like/progenitor cells,” *Carcinogenesis*, vol. 33, no. 3, pp. 652–660, Mar. 2012, doi: 10.1093/carcin/bgr317.
- [25] J. P. Xue, G. Wang, Z. Bin Zhao, Q. Wang, and Y. Shi, “Synergistic cytotoxic effect of genistein and doxorubicin on drug-resistant human breast cancer MCF-7/Adr cells,” *Oncol. Rep.*, vol. 32, no. 4, pp. 1647–1653, 2014, doi: 10.3892/or.2014.3365.
- [26] X. Yang, L. Taylor, J. Yu, M. J. Fenton, and P. Polgar, “Different effects of genistein on molecular markers related to apoptosis in two phenotypically dissimilar breast cancer cell lines,” *J. Cell. Biochem.*, vol. 82, no. 1, pp. 78–88, 2001, doi: 10.1002/jcb.1147.
- [27] M. L. De Lemos, “Effects of soy phytoestrogens genistein and daidzein on breast cancer growth,” *Ann. Pharmacother.*, vol. 35, no. 9, pp. 1118–1121, 2001, doi: 10.1345/aph.10257.
- [28] Y. Li, S. M. Meeran, S. N. Patel, H. Chen, T. M. Hardy, and T. O. Tollefsbol, “Epigenetic

- reactivation of estrogen receptor- α (ER α) by genistein enhances hormonal therapy sensitivity in ER α -negative breast cancer,” *Mol. Cancer*, vol. 12, no. 1, pp. 1–17, 2013, doi: 10.1186/1476-4598-12-9.
- [29] H. Marini *et al.*, “Breast safety and efficacy of genistein aglycone for postmenopausal bone loss: A follow-up study,” *J. Clin. Endocrinol. Metab.*, vol. 93, no. 12, 2008, doi: 10.1210/jc.2008-1087.
- [30] M. Atteritano *et al.*, “Effects of phytoestrogen genistein on cytogenetic biomarkers in postmenopausal women: 1 year randomized, placebo-controlled study,” *Eur. J. Pharmacol.*, vol. 589, no. 1–3, 2008, doi: 10.1016/j.ejphar.2008.04.049.
- [31] D. F. McMichael-Phillips *et al.*, “Effects of soy-protein supplementation on epithelial proliferation in the histologically normal human breast,” in *American Journal of Clinical Nutrition*, 1998, vol. 68, no. 6 SUPPL., doi: 10.1093/ajcn/68.6.1431S.
- [32] D. F. Hargreaves *et al.*, “Two-week dietary soy supplementation has an estrogenic effect on normal premenopausal breast,” *J. Clin. Endocrinol. Metab.*, vol. 84, no. 11, 1999, doi: 10.1210/jc.84.11.4017.
- [33] S. A. Khan *et al.*, “Soy isoflavone supplementation for breast cancer risk reduction: A randomized phase ii trial,” *Cancer Prev. Res.*, vol. 5, no. 2, 2012, doi: 10.1158/1940-6207.CAPR-11-0251.
- [34] R. Marik, M. Allu, R. Anchoori, V. Stearns, C. B. Umbricht, and S. Khan, “Potent genistein derivatives as inhibitors of estrogen receptor alpha-positive breast cancer,” *Cancer Biol. Ther.*, vol. 11, no. 10, pp. 883–892, 2011, doi: 10.4161/cbt.11.10.15184.
- [35] T. T. Y. Wang, N. Sathyamoorthy, and J. M. Phang, “Molecular effects of genistein on estrogen receptor mediated pathways,” *Carcinogenesis*, vol. 17, no. 2, pp. 271–275, 1996, doi: 10.1093/carcin/17.2.271.
- [36] I. Paterni, C. Granchi, J. A. Katzenellenbogen, and F. Minutolo, “Estrogen receptors alpha (ER α) and beta (ER β): Subtype-selective ligands and clinical potential,” *Steroids*, vol. 90, pp. 13–29, 2014, doi: 10.1016/j.steroids.2014.06.012.

- [37] E. J. Choi, J. Y. Jung, and G. H. Kim, “Genistein inhibits the proliferation and differentiation of MCF-7 and 3T3-L1 cells via the regulation of ER α expression and induction of apoptosis,” *Exp. Ther. Med.*, vol. 8, no. 2, pp. 454–458, 2014, doi: 10.3892/etm.2014.1771.
- [38] G. Pinton, S. Nilsson, and L. Moro, “Targeting estrogen receptor beta (ER β) for treatment of ovarian cancer: Importance of KDM6B and SIRT1 for ER β expression and functionality,” *Oncogenesis*, vol. 7, no. 2, 2018, doi: 10.1038/s41389-018-0027-9.
- [39] H. Jiang, J. Fan, L. Cheng, P. Hu, and R. Liu, “The anticancer activity of genistein is increased in estrogen receptor beta 1-positive breast cancer cells,” *Onco. Targets. Ther.*, vol. 11, pp. 8153–8163, 2018, doi: 10.2147/OTT.S182239.
- [40] P. Thomas and J. Dong, “Binding and activation of the seven-transmembrane estrogen receptor GPR30 by environmental estrogens: A potential novel mechanism of endocrine disruption,” *J. Steroid Biochem. Mol. Biol.*, vol. 102, no. 1-5 SPEC. ISS., pp. 175–179, 2006, doi: 10.1016/j.jsbmb.2006.09.017.
- [41] W. Wei *et al.*, “The activation of G protein-coupled receptor 30 (GPR30) inhibits proliferation of estrogen receptornegative breast cancer cells in vitro and in vivo,” *Cell Death Dis.*, vol. 5, no. 10, 2014, doi: 10.1038/cddis.2014.398.
- [42] Z. Mitri, T. Constantine, and R. O’Regan, “The HER2 Receptor in Breast Cancer: Pathophysiology, Clinical Use, and New Advances in Therapy,” *Chemother. Res. Pract.*, vol. 2012, pp. 1–7, 2012, doi: 10.1155/2012/743193.
- [43] M. S. Sakla, N. S. Shenouda, P. J. Ansell, R. S. MacDonald, and D. B. Lubahn, “Genistein affects HER2 protein concentration, activation, and promoter regulation in BT-474 human breast cancer cells,” *Endocrine*, vol. 32, no. 1, pp. 69–78, 2007, doi: 10.1007/s12020-007-9006-1.
- [44] O. Treeck, S. Schüler-Toprak, and O. Ortmann, “Estrogen Actions in Triple-Negative Breast Cancer,” *Cells*, vol. 9, no. 11, 2020, doi: 10.3390/cells9112358.
- [45] G. Y. Kim *et al.*, “Genistein Inhibits Proliferation of BRCA1 Mutated Breast Cancer

- Cells: The GPR30-Akt Axis as a Potential Target ,” *J. Cancer Prev.*, vol. 24, no. 4, pp. 197–207, 2019, doi: 10.15430/jcp.2019.24.4.197.
- [46] T. Akiyama *et al.*, “Genistein, a specific inhibitor of tyrosine-specific protein kinases.,” *J. Biol. Chem.*, vol. 262, no. 12, pp. 5592–5595, 1987, doi: 10.1016/s0021-9258(18)45614-1.
- [47] S. V. Sharma, D. A. Haber, and J. Settleman, “Cell line-based platforms to evaluate the therapeutic efficacy of candidate anticancer agents,” *Nature Reviews Cancer*, vol. 10, no. 4, pp. 241–253, 2010, doi: 10.1038/nrc2820.
- [48] R. F. Prietsch *et al.*, “Genistein induces apoptosis and autophagy in human breast MCF-7 cells by modulating the expression of proapoptotic factors and oxidative stress enzymes,” *Mol. Cell. Biochem.*, vol. 390, no. 1–2, pp. 235–242, 2014, doi: 10.1007/s11010-014-1974-x.
- [49] X. Liu *et al.*, “Genistein enhances the radiosensitivity of breast cancer cells via G2/M cell cycle arrest and apoptosis,” *Molecules*, vol. 18, no. 11, pp. 13200–13217, 2013, doi: 10.3390/molecules181113200.
- [50] J. A. Lavigne *et al.*, “Concentration-dependent effects of genistein on global gene expression in MCF-7 breast cancer cells: An oligo microarray study,” *Breast Cancer Res. Treat.*, vol. 110, no. 1, pp. 85–98, 2008, doi: 10.1007/s10549-007-9705-6.
- [51] H. Y. Shim, J. H. Park, H. D. Paik, S. Y. Nah, D. S. H. L. Kim, and Y. S. Han, “Genistein-induced apoptosis of human breast cancer MCF-7 cells involves calpain-caspase and apoptosis signaling kinase 1-p38 mitogen-activated protein kinase activation cascades,” *Anticancer. Drugs*, vol. 18, no. 6, pp. 649–657, 2007, doi: 10.1097/CAD.0b013e3280825573.
- [52] A. King-Batoon, J. M. Leszczynska, and C. B. Klein, “Modulation of gene methylation by genistein or lycopene in breast cancer cells,” *Environ. Mol. Mutagen.*, vol. 49, no. 1, pp. 36–45, 2008, doi: 10.1002/em.20363.
- [53] Y. H. Shon, S. D. Park, and K. S. Nam, “Effective chemopreventive activity of genistein against human breast cancer cells,” *J. Biochem. Mol. Biol.*, vol. 39, no. 4, pp. 448–451,

- 2006, doi: 10.5483/bmbrep.2006.39.4.448.
- [54] V. Tanos, A. Brzezinski, O. Drize, N. Strauss, and T. Peretz, “Synergistic inhibitory effects of genistein and tamoxifen on human dysplastic and malignant epithelial breast cells in vitro,” *Eur. J. Obstet. Gynecol. Reprod. Biol.*, vol. 102, no. 2, pp. 188–194, 2002, doi: 10.1016/S0301-2115(01)00582-6.
- [55] D. G. Pons, M. Nadal-Serrano, M. M. Blanquer-Rossello, J. Sastre-Serra, J. Oliver, and P. Roca, “Genistein modulates proliferation and mitochondrial functionality in breast cancer cells depending on ERalpha/ERbeta ratio,” *J. Cell. Biochem.*, vol. 115, no. 5, pp. 949–958, 2014, doi: 10.1002/jcb.24737.
- [56] J. Chen, Y. Duan, X. Zhang, Y. Ye, B. Ge, and J. Chen, “Genistein induces apoptosis by the inactivation of the IGF-1R/p-Akt signaling pathway in MCF-7 human breast cancer cells,” *Food Funct.*, vol. 6, no. 3, pp. 995–1000, 2015, doi: 10.1039/c4fo01141d.
- [57] P. Fan *et al.*, “Genistein decreases the breast cancer stem-like cell population through Hedgehog pathway,” *Stem Cell Res. Ther.*, vol. 4, no. 6, 2013, doi: 10.1186/scrt357.
- [58] S. R. Chinni, S. A. Alhasan, A. S. Multani, S. Pathak, and F. H. Sarkar, “Pleotropic effects of genistein on MCF-7 breast cancer cells,” *Int. J. Mol. Med.*, vol. 12, no. 1, pp. 29–34, 2003, doi: 10.3892/ijmm.12.1.29.
- [59] I. N. Sergeev, “Genistein induces Ca²⁺-mediated, calpain/caspase-12-dependent apoptosis in breast cancer cells,” *Biochem. Biophys. Res. Commun.*, vol. 321, no. 2, pp. 462–467, 2004, doi: 10.1016/j.bbrc.2004.06.173.
- [60] C. H. Liao, S. L. Pan, J. H. Guh, and C. M. Teng, “Genistein inversely affects tubulin-binding agent-induced apoptosis in human breast cancer cells,” *Biochem. Pharmacol.*, vol. 67, no. 11, pp. 2031–2038, 2004, doi: 10.1016/j.bcp.2004.02.010.
- [61] W. F. Chen, M. H. Huang, C. H. Tzang, M. Yang, and M. S. Wong, “Inhibitory actions of genistein in human breast cancer (MCF-7) cells,” *Biochim. Biophys. Acta - Mol. Basis Dis.*, vol. 1638, no. 2, pp. 187–196, 2003, doi: 10.1016/S0925-4439(03)00082-6.
- [62] Y. Liu *et al.*, “Genistein-induced differentiation of breast cancer stem/progenitor cells

- through a paracrine mechanism,” *Int. J. Oncol.*, vol. 48, no. 3, pp. 1063–1072, 2016, doi: 10.3892/ijo.2016.3351.
- [63] D. G. Pons *et al.*, “The phytoestrogen genistein affects inflammatory-related genes expression depending on the ER α /ER β ratio in breast cancer cells,” *Int. J. Food Sci. Nutr.*, vol. 70, no. 8, pp. 941–949, 2019, doi: 10.1080/09637486.2019.1597025.
- [64] A. Kazi, K. G. Daniel, D. M. Smith, N. B. Kumar, and Q. P. Dou, “Inhibition of the proteasome activity, a novel mechanism associated with the tumor cell apoptosis-inducing ability of genistein,” *Biochem. Pharmacol.*, vol. 66, no. 6, pp. 965–976, 2003, doi: 10.1016/S0006-2952(03)00414-3.
- [65] Q. Xie *et al.*, “Genistein inhibits DNA methylation and increases expression of tumor suppressor genes in human breast cancer cells,” *Genes Chromosom. Cancer*, vol. 53, no. 5, pp. 422–431, 2014, doi: 10.1002/gcc.22154.
- [66] H. Pan *et al.*, “Genistein inhibits MDA-MB-231 triple-negative breast cancer cell growth by inhibiting NF- κ B activity via the Notch-1 pathway,” *Int. J. Mol. Med.*, vol. 30, no. 2, pp. 337–343, 2012, doi: 10.3892/ijmm.2012.990.
- [67] M. F. Ullah *et al.*, “Soy isoflavone genistein induces cell death in breast cancer cells through mobilization of endogenous copper ions and generation of reactive oxygen species,” *Mol. Nutr. Food Res.*, vol. 55, no. 4, pp. 553–559, 2011, doi: 10.1002/mnfr.201000329.
- [68] K. Dampier, E. A. Hudson, L. M. Howells, M. M. Manson, R. A. Walker, and A. Gescher, “Differences between human breast cell lines in susceptibility towards growth inhibition by genistein,” *Br. J. Cancer*, vol. 85, no. 4, pp. 618–624, 2001, doi: 10.1054/bjoc.2001.1980.
- [69] O. C. Kousidou, T. N. Mitropoulou, A. E. Roussidis, D. Kletsas, A. D. Theocharis, and N. K. Karamanos, “Genistein suppresses the invasive potential of human breast cancer cells through transcriptional regulation of metalloproteinases and their tissue inhibitors,” *Int. J. Oncol.*, vol. 26, no. 4, pp. 1101–1109, 2005, doi: 10.3892/ijo.26.4.1101.

- [70] L. Gong, Y. Li, A. Nedeljkovic-Kurepa, and F. H. Sarkar, "Inactivation of NF- κ B by genistein is mediated via Akt signaling pathway in breast cancer cells," *Oncogene*, vol. 22, no. 30, pp. 4702–4709, 2003, doi: 10.1038/sj.onc.1206583.
- [71] Y. Li, H. Chen, T. M. Hardy, and T. O. Tollefsbol, "Epigenetic Regulation of Multiple Tumor-Related Genes Leads to Suppression of Breast Tumorigenesis by Dietary Genistein," *PLoS One*, vol. 8, no. 1, 2013, doi: 10.1371/journal.pone.0054369.
- [72] Y. Fang *et al.*, "Quantitative phosphoproteomics reveals genistein as a modulator of cell cycle and DNA damage response pathways in triple-negative breast cancer cells," *Int. J. Oncol.*, vol. 48, no. 3, pp. 1016–1028, 2016, doi: 10.3892/ijo.2016.3327.
- [73] T. T. Rajah, N. Du, N. Drews, and R. Cohn, "Genistein in the presence of 17 β -estradiol inhibits proliferation of ER β breast cancer cells," *Pharmacology*, vol. 84, no. 2, pp. 68–73, 2009, doi: 10.1159/000226123.
- [74] D. Ye, Z. Li, and C. Wei, "Genistein inhibits the S-phase kinase-associated protein 2 expression in breast cancer cells," *Exp. Ther. Med.*, vol. 15, no. 1, pp. 1069–1075, 2018, doi: 10.3892/etm.2017.5489.
- [75] J. Gwin, N. Drews, S. Ali, J. Stamschror, M. Sorenson, and T. T. Rajah, "Effect of genistein on p90RSK phosphorylation and cell proliferation in T47D breast cancer cells," *Anticancer Res.*, vol. 31, no. 1, pp. 209–214, 2011.
- [76] A. M. Sotoca *et al.*, "Quantitative Proteomics and Transcriptomics Addressing the Estrogen Receptor Subtype-mediated Effects in T47D Breast Cancer Cells Exposed to the Phytoestrogen Genistein," *Mol. Cell. Proteomics*, vol. 10, no. 1, p. M110.002170, 2011, doi: 10.1074/mcp.m110.002170.
- [77] W. Y. Lee, S. C. Huang, C. C. Tzeng, T. L. Chang, and K. F. Hsu, "Alterations of metastasis-related genes identified using an oligonucleotide microarray of genistein-treated HCC1395 breast cancer cells," *Nutr. Cancer*, vol. 58, no. 2, pp. 239–246, 2007, doi: 10.1080/01635580701328636.
- [78] M. G. Donovan, O. I. Selmin, T. C. Doetschman, and D. F. Romagnolo, "Epigenetic

- activation of BRCA1 by genistein in vivo and triple negative breast cancer cells linked to antagonism toward aryl hydrocarbon receptor,” *Nutrients*, vol. 11, no. 11, 2019, doi: 10.3390/nu11112559.
- [79] C. De La Parra *et al.*, “Soy isoflavone genistein-mediated downregulation of miR-155 contributes to the anticancer effects of genistein,” *Nutr. Cancer*, vol. 68, no. 1, pp. 154–164, 2016, doi: 10.1080/01635581.2016.1115104.
- [80] G. Voelcker, B. Pfeiffer, A. Schnee, and H. J. Hohorst, “Increased antitumour activity of mesyl-I-aldophosphamide- perhydrothiazine, in vivo but not in vitro, compared to I-aldophosphamide- perhydrothiazine,” *J. Cancer Res. Clin. Oncol.*, vol. 126, no. 2, pp. 74–78, 2000, doi: 10.1007/s004320050012.
- [81] Z. Mai, G. L. Blackburn, and J. R. Zhou, “Genistein sensitizes inhibitory effect of tamoxifen on the growth of estrogen receptor-positive and HER2-overexpressing human breast cancer cells,” *Mol. Carcinog.*, vol. 46, no. 7, pp. 534–542, 2007, doi: 10.1002/mc.20300.
- [82] M. Katdare, M. Osborne, and N. T. Telang, “Soy isoflavone genistein modulates cell cycle progression and induces apoptosis in HER-2/neu oncogene expressing human breast epithelial cells,” *Int. J. Oncol.*, vol. 21, no. 4, pp. 809–815, 2002, doi: 10.3892/ijo.21.4.809.
- [83] C. B. Avci *et al.*, “Genistein-induced mir-23b expression inhibits the growth of breast cancer cells,” *Wspolczesna Onkol.*, vol. 19, no. 1, pp. 32–35, 2015, doi: 10.5114/wo.2014.44121.
- [84] D. F. Romagnolo, M. G. Donovan, A. J. Papoutsis, T. C. Doetschman, and O. I. Selmin, “Genistein prevents BRCA1 CpG methylation and proliferation in human breast cancer cells with activated aromatic hydrocarbon receptor,” *Curr. Dev. Nutr.*, vol. 1, no. 6, pp. 1–11, 2017, doi: 10.3945/cdn.117.000562.
- [85] M. Privat, C. Aubel, S. Arnould, Y. Communal, M. Ferrara, and Y. J. Bignon, “Breast cancer cell response to genistein is conditioned by BRCA1 mutations,” *Biochem. Biophys. Res. Commun.*, vol. 379, no. 3, pp. 785–789, 2009, doi: 10.1016/j.bbrc.2008.12.151.

- [86] S. A. Vantighem, S. M. Wilson, C. O. Postenka, W. Al-Katib, A. B. Tuck, and A. F. Chambers, "Dietary genistein reduces metastasis in a postsurgical orthotopic breast cancer model," *Cancer Res.*, vol. 65, no. 8, pp. 3396–3403, 2005, doi: 10.1158/0008-5472.CAN-04-4109.
- [87] S. Balabhadrapathruni, T. J. Thomas, E. J. Yurkow, P. S. Amenta, and T. Thomas, "Effects of genistein and structurally related phytoestrogens on cell cycle kinetics and apoptosis in MDA-MB-468 human breast cancer cells," *Oncol. Rep.*, vol. 7, no. 1, pp. 3–12, 2000, doi: 10.3892/or.7.1.3.
- [88] T. N. Mitropoulou, G. N. Tzanakakis, D. Nikitovic, A. Tsatsakis, and N. K. Karamanos, "In vitro effects of genistein on the synthesis and distribution of glycosaminoglycans / proteoglycans by estrogen receptor-positive and -negative human breast cancer epithelial cells," *Anticancer Res.*, vol. 22, no. 5, pp. 2841–2846, 2002.
- [89] L. Fioravanti, V. Cappelletti, P. Miodini, E. Ronchi, M. Brivio, and G. Di Fronzo, "Genistein in the control of breast cancer cell growth: Insights into the mechanism of action in vitro," *Cancer Lett.*, vol. 130, no. 1–2, pp. 143–152, 1998, doi: 10.1016/S0304-3835(98)00130-X.
- [90] D. T. Zava and G. Duwe, "Estrogenic and antiproliferative properties of genistein and other flavonoids in human breast cancer cells in vitro," *Nutr. Cancer*, vol. 27, no. 1, pp. 31–40, 1997, doi: 10.1080/01635589709514498.
- [91] L. A. Fitzpatrick, "Soy isoflavones: Hope or hype?," *Maturitas*, vol. 44, no. SUPPL. 2003, doi: 10.1016/S0378-5122(02)00345-6.
- [92] K. D. R. Setchell and E. Lydeking-Olsen, "Dietary phytoestrogens and their effect on bone: Evidence from in vitro and in vivo, human observational, and dietary intervention studies," in *American Journal of Clinical Nutrition*, 2003, vol. 78, no. 3 SUPPL., doi: 10.1093/ajcn/78.3.593s.
- [93] J. Markovits *et al.*, "Inhibitory Effects of the Tyrosine Kinase Inhibitor Genistein on Mammalian DNA Topoisomerase II," *Cancer Res.*, vol. 49, no. 18, pp. 5111–5117, 1989.

- [94] A. Okura, H. Arakawa, H. Oka, T. Yoshinari, and Y. Monden, “Effect of genistein on topoisomerase activity and on the growth of [VAL 12]Ha-ras-transformed NIH 3T3 cells,” *Biochem. Biophys. Res. Commun.*, vol. 157, no. 1, pp. 183–189, 1988, doi: 10.1016/S0006-291X(88)80030-5.
- [95] O. Kousidou, G. Tzanakakis, and N. Karamanos, “Effects of the Natural Isoflavonoid Genistein on Growth, Signaling Pathways and Gene Expression of Matrix Macromolecules by Breast Cancer Cells,” *Mini-Reviews Med. Chem.*, vol. 6, no. 3, pp. 331–337, 2006, doi: 10.2174/138955706776073420.
- [96] C. M. J. Conklin, J. F. Bechberger, D. MacFabe, N. Guthrie, E. M. Kurowska, and C. C. Naus, “Genistein and quercetin increase connexin43 and suppress growth of breast cancer cells,” *Carcinogenesis*, vol. 28, no. 1, pp. 93–100, 2007, doi: 10.1093/carcin/bgl106.
- [97] K. A. A. Thasni *et al.*, “Genistein induces apoptosis in ovarian cancer cells via different molecular pathways depending on Breast Cancer Susceptibility gene-1 (BRCA1) status,” *Eur. J. Pharmacol.*, vol. 588, no. 2–3, pp. 158–164, 2008, doi: 10.1016/j.ejphar.2008.04.041.
- [98] R. C. Santell, N. Kieu, and W. G. Helferich, “Genistein inhibits growth of estrogen-independent human breast cancer cells in culture but not in athymic mice,” *J. Nutr.*, vol. 130, no. 7, pp. 1665–1669, 2000, doi: 10.1093/jn/130.7.1665.
- [99] Z. E. Sauna, M. M. Smith, M. Muller, K. M. Kerr, and S. V. Ambudkar, “The mechanism of action of multidrug-resistance-linked P-glycoprotein,” *J. Bioenerg. Biomembr.*, vol. 33, no. 6, pp. 481–491, 2001, doi: 10.1023/A:1012875105006.
- [100] E. Rivera, “Implications of anthracycline-resistant and taxane-resistant metastatic breast cancer and new therapeutic options,” *Breast Journal*, vol. 16, no. 3, pp. 252–263, 2010, doi: 10.1111/j.1524-4741.2009.00896.x.
- [101] J. König *et al.*, “Expression and localization of human multidrug resistance protein (ABCC) family members in pancreatic carcinoma,” *Int. J. Cancer*, vol. 115, no. 3, pp. 359–367, 2005, doi: 10.1002/ijc.20831.

- [102] C. H. M. Versantvoort *et al.*, “Genistein modulates the decreased drug accumulation in non-p-glycoprotein mediated multidrug resistant tumour cells,” *Br. J. Cancer*, vol. 68, no. 5, pp. 939–946, 1993, doi: 10.1038/bjc.1993.458.
- [103] Y. Imai, S. Tsukahara, S. Asada, and Y. Sugimoto, “Phytoestrogens/flavonoids reverse breast cancer resistance protein/ABCG2-mediated multidrug resistance,” *Cancer Res.*, vol. 64, no. 12, pp. 4346–4352, 2004, doi: 10.1158/0008-5472.CAN-04-0078.
- [104] C. Lattrich, J. Lubig, A. Springwald, R. Goerse, O. Ortmann, and O. Treeck, “Additive effects of trastuzumab and genistein on human breast cancer cells,” *Anticancer. Drugs*, vol. 22, no. 3, pp. 253–261, 2011, doi: 10.1097/CAD.0b013e3283427bb5.
- [105] D. G. Pons, M. Nadal-Serrano, M. Torrens-Mas, J. Oliver, and P. Roca, “The Phytoestrogen Genistein Affects Breast Cancer Cells Treatment Depending on the ER α /ER β Ratio,” *J. Cell. Biochem.*, vol. 117, no. 1, pp. 218–229, 2016, doi: 10.1002/jcb.25268.
- [106] K. Sahin *et al.*, “Inhibitory effects of combination of lycopene and genistein on 7,12-Dimethyl benz(a)anthracene-induced breast cancer in rats,” *Nutr. Cancer*, vol. 63, no. 8, pp. 1279–1286, 2011, doi: 10.1080/01635581.2011.606955.
- [107] H. Marini *et al.*, “Effects of the phytoestrogen genistein on bone metabolism in osteopenic postmenopausal women: A randomized trial,” *Ann. Intern. Med.*, vol. 146, no. 12, 2007, doi: 10.7326/0003-4819-146-12-200706190-00005.
- [108] R. D’Anna *et al.*, “Effects of the phytoestrogen genistein on hot flushes, endometrium, and vaginal epithelium in postmenopausal women: A 1-year randomized, double-blind, placebo-controlled study,” *Menopause*, vol. 14, no. 4, 2007, doi: 10.1097/01.gme.0000248708.60698.98.
- [109] M. Razandi, A. Pedram, E. M. Rosen, and E. R. Levin, “BRCA1 Inhibits Membrane Estrogen and Growth Factor Receptor Signaling to Cell Proliferation in Breast Cancer,” *Mol. Cell. Biol.*, vol. 24, no. 13, 2004, doi: 10.1128/mcb.24.13.5900-5913.2004.
- [110] S. J. Boulton, “Cellular functions of the BRCA tumour-suppressor proteins,” *Biochemical*

- Society Transactions*, vol. 34, no. 5. 2006, doi: 10.1042/BST0340633.
- [111] G. Maskarinec, Y. Takata, A. A. Franke, A. E. Williams, and S. P. Murphy, “A 2-year soy intervention in premenopausal women not change mammographic densities,” *J. Nutr.*, vol. 134, no. 11, 2004, doi: 10.1093/jn/134.11.3089.
- [112] G. Maskarinec, A. E. Williams, and L. Carlin, “Mammographic densities in a one-year isoflavone intervention,” *Eur. J. Cancer Prev.*, vol. 12, no. 2, 2003, doi: 10.1097/00008469-200304000-00011.
- [113] G. Maskarinec, A. E. Williams, J. S. Inouye, F. Z. Stanczyk, and A. A. Franke, “A randomized isoflavone intervention among premenopausal women,” *Cancer Epidemiol. Biomarkers Prev.*, vol. 11, no. 2, 2002.
- [114] A. Daina, O. Michielin, and V. Zoete, “SwissADME: A free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules,” *Sci. Rep.*, vol. 7, 2017, doi: 10.1038/srep42717.
- [115] D. E. V. Pires, T. L. Blundell, and D. B. Ascher, “pkCSM: Predicting small-molecule pharmacokinetic and toxicity properties using graph-based signatures,” *J. Med. Chem.*, vol. 58, no. 9, pp. 4066–4072, 2015, doi: 10.1021/acs.jmedchem.5b00104.
- [116] F. Cheng *et al.*, “AdmetSAR: A comprehensive source and free tool for assessment of chemical ADMET properties,” *Journal of Chemical Information and Modeling*, vol. 52, no. 11. ACS Publications, pp. 3099–3105, 2012, doi: 10.1021/ci300367a.
- [117] N. Toomula, D. Sathish Kumar, A. Kumar, and M. Phaneendra, “Role of pharmacokinetic studies in drug discovery,” *Journal of Bioequivalence and Bioavailability*, vol. 3, no. 11. pp. 263–267, 2011, doi: 10.4172/jbb.1000097.