

Published in final edited form as:

*Molecules.* ; 18(4): 4357–4372. doi:10.3390/molecules18044357.

## Application Scope and Limitations of TADDOL-Derived Chiral Ammonium Salt Phase-Transfer Catalysts

Guddeangadi N. Gururaja, Richard Herchl, Antonia Pichler, Katharina Gratzner, and Mario Waser\*

Institute of Organic Chemistry, Johannes Kepler University Linz, Altenbergerstraße 69, 4040 Linz, Austria

### Abstract

We have recently introduced a new class of chiral ammonium salt catalysts derived from easily available TADDOLs. To get a full picture of the scope of application and limitations of our catalysts we tested them in a variety of different important transformations. We found that, although these compounds have recently shown their good potential in the asymmetric  $\alpha$ -alkylation of glycine Schiff bases, they clearly failed when we attempted to control more reactive nucleophiles like  $\beta$ -keto esters. On the other hand, when using them to catalyse the addition of glycine Schiff bases to different Michael acceptors it was found necessary to carefully optimize the reaction conditions for every single substrate class, as seemingly small structural changes sometimes required the use of totally different reaction conditions. Under carefully optimized conditions enantiomeric ratios up to 91:9 could be achieved in the addition of glycine Schiff bases to acrylates, whereas acrylamides and methyl vinyl ketone gave slightly lower selectivities (up to e.r. 77:23 in these cases). Thus, together with additional studies towards the syntheses of these catalysts we have now a very detailed understanding about the scope and limitations of the synthesis sequence to access our PTCs and about the application scope of these catalysts in asymmetric transformations.

### Keywords

asymmetric catalysis; tartaric acid;  $\alpha$ -alkylation; Michael addition

## 1. Introduction

Design, syntheses, and applications of chiral phase-transfer catalysts (PTCs) have attracted considerable interest over the last three decades [1–6]. The high potential of asymmetric phase-transfer catalysis can be attributed to several reasons (e.g., mild aqueous reaction conditions, operational simplicity, easily handled catalysts, scalability,...), making it a powerful and versatile methodology for a broad scope of different applications where other

© 2013 by the authors

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

\* Author to whom correspondence should be addressed; Mario.waser@jku.at; Tel.: +43-732-2468-8748; Fax: +43-732-2468-8747.

catalytic principles clearly fail. Among the different commonly employed catalytically active structural motives, chiral quaternary ammonium salts have found the most widespread applications so far [1–6]. Following the seminal reports of Wynberg [7] and a group of Merck scientists [8] employing cinchona alkaloid-derived quaternary ammonium salts for asymmetric epoxide formation [7] and methylation of a phenylindanone derivative [8], cinchona alkaloids remained the privileged source of chirality for syntheses and investigations concerning novel phase-transfer catalysts and applications thereof until the beginning of the 21st century. Pioneering work by the groups of O'Donnell [9,10], Lygo [11,12], and Corey [13,14] resulted in the development of several highly stereoselective applications using a variety of structurally carefully optimized cinchona alkaloid-based PTCs. Due to their high catalytic potential and broad application scope, catalysts based on this easily obtained naturally occurring chiral backbone still belong to the most commonly employed and most thoroughly investigated PTCs as shown in recent reports by the groups of Li Deng [15,16], Jørgensen [17,18], and others [19–29].

In 1999, Maruoka introduced a new designer catalyst system by using  $C_2$ -symmetric binaphthyl-based chiral spiro ammonium salts [30]. These Maruoka catalysts were found to be highly effective for a variety of asymmetric transformations (e.g., Michael additions,  $\alpha$ -amino acid syntheses, epoxidations, aldol-type reactions, isoxazoline syntheses,...), even using only minimum amounts of catalysts (<1 mol%) [4,5,30–35], thus belonging to the most powerful and versatile PTCs known to date. In addition, also Shibasaki's tartaric acid-derived bidentate PTCs [36–38] and Lygo's biphenyl-based spirocyclic catalysts [39,40] have proven their potential in different asymmetric applications.

However, despite more than three decades of active research in this field it is somewhat surprising that besides the already mentioned privileged catalyst structures (Figure 1) only a few other classes of chiral ammonium salt PTCs have been reported so far [41–45]. Despite sometimes very exhaustive and careful structure-activity based investigations and optimizations [44,45], none of these other classes has so far reached the catalytic potential and application scope of especially the cinchona-based catalysts and the Maruoka-type catalysts.

One of the main demands for novel catalysts is easy accessibility from readily available chiral starting materials. Among the easily available natural chiral sources, tartaric acid (**1**) has obtained a prominent position, especially due to the fact that both enantiomers are readily available in sufficient quantities. Although Shibasaki *et al.* have demonstrated the potential of tartaric acid-derived bidentate PTCs [36–38], others were less successful in their attempts to synthesize powerful tartaric acid-derived quaternary ammonium salt catalysts [41,42]. Based on the high potential of tartaric acid-derived easily obtainable tetraaryl-2,2-dimethyl-1,3-dioxolan-4,5-dimethanols (TADDOLs, **2**) as chiral ligands in (transition-) metal catalysis [46,47] we have recently carried out systematic investigations to use this unique structural motive for the syntheses of chiral *N*-spiroquaternary ammonium salt catalysts [48,49]. After a careful route development we were able to obtain more than 30 differently substituted  $C_1$ - or  $C_2$ -symmetric *N*-spiro catalysts **3**. The catalytic potential of these PTCs was initially tested for the benchmark  $\alpha$ -alkylation of glycine Schiff base **4** and the *p*-biphenyl containing acetonide-based catalyst **3a** turned out to be the most powerful

one therein, giving access to a variety of amino acid derivatives **5** in high yields and with satisfying enantioselectivities (Scheme 1). In contrast, testing this catalyst for the asymmetric epoxidation of chalcone **6** resulted in the formation of racemic **7** only [49].

To further elucidate the potential and the application scope of this novel and straightforwardly available class of catalysts we have now carried out a detailed screening of different other important test reactions. In addition further attempts to systematically modify the catalyst structures have been undertaken.

## 2. Results and Discussion

### 2.1. Late Stage Catalyst Modification

We have recently observed that the nature of the acetal protecting group of the catalyst has a strong influence on the catalyst performance in the benchmark  $\alpha$ -alkylation of **4** [49]. Unfortunately, our standard strategy to access these catalysts required introduction of the acetal group very early in the sequence already, making a rapid structural diversification tedious, especially as we found that, based on the nature of the acetal group, the subsequent steps sometimes proceeded significantly lower yielding (or were even not possible anymore) and purification of the final products also became difficult [49]. Thus, we targeted a late stage acetal-cleavage – acetal-formation sequence starting from readily available **3a** to access differently acetal-protected catalysts straightforwardly. Interestingly the dioxolane moiety was found to be exceptionally acid-stable and it required treatment with concentrated trifluoroacetic acid (TFA/H<sub>2</sub>O = 95:5) to obtain the diol **8** (Scheme 2). Initial attempts to test the free-OH containing ammonium salt **8** as a catalyst for the reaction of the glycine Schiff base *t*-butyl ester **4a** with benzylbromide (**9**) gave surprising results. First the enantioselectivity was rather low (e.r. 72:28) under the previously optimized conditions and, even more interesting, the product **5a** was only obtained in less than 20% yield. Furthermore the catalyst could not be recovered, but decomposed almost quantitatively under the basic reaction conditions. This pronounced base-sensitivity was also observed when we attempted an *O*-benzylation or *O*-methylation of **8** even using just bicarbonates as the bases, thus making syntheses of diether-derivatives of these catalysts impossible (these compounds were also not accessible using our standard procedure) [49]. In contrast, compound **8** was found to be rather acid-stable and dioxolane-formation with different ketones or aldehydes could be carried out in the presence of triflic acid. Noteworthy, these reactions only proceeded with an excess of this strong acid, whereas other strategies failed, thus giving the corresponding ammonium triflates **3** first. However, these catalysts absolutely failed in the test reaction as no turnover and only modest enantioselectivities were observed (an illustrative example using catalyst **3b** is given in Scheme 2 and similar results were obtained using other differently acetal-protected ammonium salts prepared by this strategy). As we recently observed a significant counter anion influence in this alkylation (e.g., changing Br<sup>-</sup> for other halides did not affect the activity, but using BF<sub>4</sub><sup>-</sup> or PF<sub>6</sub><sup>-</sup> reduced the catalytic potential dramatically [49]), we tried different counter anion exchange methods to obtain the ammonium bromides. However, only the use of HBr allowed us to replace the triflate anion to some extent, but always accompanied with significant decomposition, which made isolation and purification by standard methods very tedious and low yielding. Testing this

(not perfectly pure) catalyst **3c** an improved, but still not satisfying  $\alpha$ -alkylation result was obtained (see Scheme 2). Noteworthy, we have recently prepared catalyst **3c** via our conventional strategy (but were not able to obtain it in sufficient yield and purity either), which showed a better catalytic potential than material obtained by the new acetal transformation - counter anion exchange method [49]. Accordingly, although this late-stage modification strategy seemed promising at first, it did not allow us to obtain the targeted catalysts in sufficient quality and with strict control of the nature of the counter anion and thus could not be readily and reliably used for stereoselective applications.

## 2.2. Asymmetric $\alpha$ -Alkylation of $\beta$ -Keto Esters

As we have recently proven the high potential of our catalysts for asymmetric  $\alpha$ -alkylation reactions of glycine Schiff bases we also tested their applicability for the asymmetric  $\alpha$ -alkylation of  $\beta$ -keto esters. As a test reaction we choose the benzylation of esters **10** under a variety of different liquid/liquid or liquid/solid phase-transfer conditions (Scheme 3). Unfortunately, after an extensive screening of a variety of different conditions and also differently substituted esters **10** we were not able to obtain the products **11** with any reasonable enantiopurity. In contrast, the non-catalysed racemic background alkylation of this highly acidic starting material was found to be the dominating reaction therein. Thus, it seems reasonable that formation of the required chiral ion pair between the ammonium salt catalyst and the enolate of **10** is too slow compared to the non-catalysed racemic background reaction, thus explaining the low enantioselectivities observed in this specific test reaction.

## 2.3. Asymmetric Michael Addition Reactions of Glycine Schiff Bases

Besides asymmetric  $\alpha$ -alkylation reactions also the analogous Michael addition reactions have emerged as powerful applications of asymmetric PTCs in the past. To elucidate the potential of our catalysts for such transformations we investigated their use for the reaction of glycine Schiff bases **4** with a variety of different acrylates **12** next (Table 1 gives an overview of the most significant results obtained in a thorough screening of different reaction conditions, reagents, and catalysts). Initial experiments were carried out in analogy to our recent  $\alpha$ -alkylation protocol using the standard biphenyl catalyst **3a** (10 mol%) in toluene as the solvent and with aqueous KOH as the base at 0 °C (entries 1 and 2). Surprisingly, absolutely no enantioselectivity could be obtained in the addition of the *t*-butyl ester **4a** to methyl acrylate **12a**. Also the use of solid KOH (entry 3) or the use of weaker aqueous bases like K<sub>3</sub>PO<sub>4</sub> (entry 4) or different alkali carbonates (entries 5 and 6) gave racemic **13a** in low yields only. However, when we used an excess of solid Cs<sub>2</sub>CO<sub>3</sub> as the base, the product **13a** was obtained in modest enantioselectivity (e.r. 66:34) and with good yield (entry 7). Reducing the reaction temperature to -20 °C gave a slightly improved enantiomeric ratio of 71:29 (the e.r. could be increased to 75:25 upon using 20 mol% of catalyst). At this point we observed that the use of recovered catalyst (after extractive workup and column chromatography) resulted in a significantly reduced enantioselectivity compared to the use of freshly prepared catalyst (entry 9 vs. entry 8, this also explains why using 20 mol% of catalyst allowed us to obtain the product in higher yield and with better selectivity than using 10 mol%). As the only difference in these two cases seems to be the nature of the counter anion due to an exchange of the bromide to either carbonate or chloride

(due to brine extraction) we next tested the systematically modified catalysts **3d** (with  $\text{BF}_4^-$  as the counter anion) and **3e** ( $\text{PF}_6^-$ ) (entries 10 and 11). Unfortunately, in neither case an increased selectivity could be achieved [50,51]. Noteworthy, the use of those catalysts prepared via our acetal-deprotection – protection strategy having either a trifluoroacetate or a triflate counter anion (see Scheme 2) also did not allow us to obtain the Michael product in any reasonable quantity and enantiopurity. Accordingly, to obtain reproducible and comparable results for the rest of these studies we always used freshly prepared ammonium bromide catalysts (comparable results were obtained when recovered catalyst was refluxed in acetonitrile with an excess of KBr for 2 days, thus giving the corresponding ammonium bromide again). Next, a screening of different solvents revealed mesitylene to be the best-suited one (non-aromatic solvents were found to be not suitable). Interestingly, addition of different additives was found to have no beneficial effect. For example the use of molecular sieves significantly suppressed the yield and the enantioselectivity (entry 15) whereas on the other hand addition of a proton source (as described recently to be beneficial by Lygo *et al.* [52]) also did not allow us to achieve a higher selectivity (entry 16). Unfortunately also the addition of different inorganic salts (e.g., CsBr, KBr, CsF or others) did not have any beneficial effect.

Testing different esters **4** next, we observed a strong influence of the ester moiety (compare entries 14, 17, and 18). In contrast to our recent results obtained when we used these esters for asymmetric  $\alpha$ -alkylation reactions where we found the *t*-butyl ester Schiff base **4a** to give by far the best yields and highest selectivities [49], the opposite tendency was observed in the present Michael addition. Herein the methyl ester Schiff base **4c** was found to be the best suited one, giving **13c** with an e.r. of 78:22 at 0 °C (entry 18). Lowering the reaction temperature resulted in an improved selectivity, albeit with a significantly lower yield (entry 19), which could be overcome by using 20 mol% catalyst instead, giving **13c** with 90:10 e.r. and in 71% yield (entry 20). Under these optimized conditions we employed different acrylates **12** next (entries 20–23). Interestingly, whereas the methyl, *n*-butyl, and benzyl esters performed similarly well, no product was obtained when we used the *t*-butyl ester **12c**. Finally, to investigate the importance of the catalyst substituents we performed the reaction between methyl Schiff base **4c** and methyl acrylate **12a** in the presence of different  $C_1$  or  $C_2$ -symmetric PTCs **3** (The most illustrative results are summarized in entries 24–26, Table 1). In analogy to our recent alkylation results the phenyl-based catalyst **3f** performed less selective and lower yielding than catalyst **3a** (entry 24 vs. entry 20). Also the *n*-butyl containing  $C_2$ -symmetric catalyst **3g** was found to be less selective in both, alkylation [49] and Michael reaction (entry 25). Interestingly, the  $C_1$ -symmetric catalyst **3h** performed even slightly better in the Michael addition than **3a** (entry 26 vs. entry 20), which is in contrast to our recent alkylation results, where this catalyst was slightly less selective than **3a** [49]. Unfortunately, no further improvement could be achieved by using any other of our recently introduced catalysts anymore.

Having optimized the conditions for the asymmetric Michael addition of glycine Schiff bases **4** to acrylic acid esters **12** we then screened the use of other Michael acceptors like acrylamides **14** and methyl vinyl ketone (MVK, **15**) (Table 2 gives a comprehensive overview about the results obtained with catalyst **3a**). Use of acrylamides as acceptors in

PT-catalysed Michael reactions has only sparingly been reported in the past [40,55] and, interestingly enough, these seemingly subtle changes in the Michael acceptor resulted in a totally different behaviour in our test reaction with glycine Schiff bases **4**. Using the conditions that have been optimized for the addition to acrylates **12** first, we found that only small amounts of racemic product **16a** could be obtained when reacting Schiff base **4a** with the acrylamide **14a** (entries 1 and 2). Also the use of other weaker solid bases did not give any product (entries 3 and 4 give just two examples of the tested ones) whereas the use of solid KOH (entry 5) gave a significant conversion, but almost no enantioselectivity (the same was observed using other solid hydroxide bases). Interestingly, when we used aqueous KOH, the product was obtained in reasonable yield and with a low enantiomeric ratio of 65:35 (entry 6). Similar results were obtained using other aqueous alkali hydroxide bases (entries 7 and 8) with RbOH being the best-suited one (e.r. 69:31 at 0 °C). Noteworthy, the observed tendency that aqueous hydroxide bases perform better than solid bases is in sharp contrast to our observations with the Michael additions to acrylates (Table 1) where aqueous bases clearly failed whereas solid ones performed superior. Further testing of different solvents and different conditions showed that for this reaction toluene is the superior solvent. Finally running the reaction for 2 days the product **16a** was obtained in reasonable 65% yield and with a modest enantiomeric ratio of 75:25 (no further improvement was possible also due to the low reaction rate of this reaction at reduced temperature). Unfortunately, using the *N,N*-diphenyl acrylamide **14b** as an acceptor the enantioselectivity dropped significantly again (entries 11 and 12) and testing secondary amides (not in the table) no product was formed, thus illustrating that Michael addition to acrylamides under asymmetric phase-transfer conditions is a rather challenging transformation. Using MVK as the acceptor both liquid/liquid and liquid/solid conditions gave the product, but the milder liquid/liquid conditions were found to be slightly better suited to give the Michael product **17** with a modest enantiomeric ratio of 77:23 (entry 14).

### 3. Experimental

#### 3.1. General

<sup>1</sup>H- and <sup>13</sup>C-NMR spectra were recorded on a Bruker Avance III 300 MHz spectrometer. All NMR spectra were referenced on the solvent peak. High resolution mass spectra were obtained using an Agilent 6520 Q-TOF mass spectrometer with an ESI source and an Agilent G1607A coaxial sprayer. All analyses were made in the positive ionization mode. Purine (exact mass for [M+H]<sup>+</sup> = 121.050873) and 1,2,3,4,5,6-hexakis(2,2,3,3-tetrafluoropropoxy)-1,3,5,2,4,6-triazatriphosphinane (exact mass for [M+H]<sup>+</sup> = 922.009798) were used for internal mass calibration. IR spectra were recorded on a Shimadzu IR Affinity-1 Fourier Transform infrared spectrometer. Optical rotations were recorded on a Perkin Elmer Polarimeter Model 241 MC. HPLC was performed using a Dionex Summit HPLC system with a Chiralcel OD-H (250 × 4.6 mm, 5 μm) or a Chiralcel OD-R (250 × 4.6 mm, 10 μm) chiral stationary phase. All chemicals were purchased from commercial suppliers and used without further purification unless otherwise stated. All reactions were carried out under inert atmosphere (Ar). Catalysts **3** were prepared as described recently [48,49].

### 3.2. Conditions A: General Procedure for the Phase-Transfer Catalysed Michael-Reaction under Liquid/Solid Phase-Transfer Conditions

Reactions were carried out using 0.2 mmol of the Schiff base **4**. The catalyst **3** (10–20 mol %) and Schiff base **4** (1 eq.) were dissolved in degassed mesitylene (0.15 M) and Cs<sub>2</sub>CO<sub>3</sub> (20 eq.) was added. The vigorously stirred solution (>1200 rpm) was cooled to –20 °C and afterwards the corresponding electrophile (1.5 eq.) was added. After 22 h at –20 °C the reaction mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O, the combined organic phases were dried over Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness and purified by column chromatography (silica gel). The Michael-addition products were isolated using heptanes/EtOAc = 40:1 to 10:1 as the eluent.

### 3.3. Conditions B: General Procedure for the Phase-Transfer Catalysed Michael-Reaction under Liquid/Liquid Phase-Transfer Conditions

Reactions were carried out using 0.2 mmol of the Schiff base **4**. The catalyst **3** (20 mol%) and Schiff base **4** (1 eq.) were dissolved in degassed toluene (0.15 M) and aqueous RbOH (50%) (25 eq.) was added. The vigorously stirred solution (>1200 rpm) was cooled to –20 °C and afterwards the corresponding electrophile (2 eq.) was added. After 48 h at –20 °C the reaction mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O, the combined organic phases were dried over Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness and purified by column chromatography (silica gel). The Michael-addition products were isolated using heptanes/EtOAc = 10:1 as the eluent.

**(S)-(–)-13a**—Obtained as a colourless oil in 74% yield and with e.r. = 69:31 upon reacting Schiff base **4a** with acrylate **12a** in the presence of 10 mol% catalyst at 0 °C under conditions A. Analytical data are in full accordance with those reported in literature [52,53].  $[\alpha]_D^{20}$  (c = 0.35, CHCl<sub>3</sub>) = –32.8°; <sup>1</sup>H-NMR (δ, CDCl<sub>3</sub>, 298 K): 1.44 (s, 9H), 2.17–2.26 (m, 2H), 2.34–2.41 (m, 2H), 3.59 (s, 3H), 3.93–3.99 (m, 1H), 7.14–7.21 (m, 2H), 7.29–7.47 (m, 6H), 7.61–7.68 (m, 2H) ppm; <sup>13</sup>C-NMR (δ, CDCl<sub>3</sub>, 298 K): 28.0, 28.6, 30.5, 51.5, 64.8, 81.2, 127.8, 128.0, 128.4, 128.6, 128.8, 130.3, 136.5, 139.5, 170.8, 172.9, 173.6 ppm; IR (film):  $\nu = 2978, 2926, 1738, 1707, 1661, 1599, 1578, 1449, 1369, 1319, 1279, 1260, 1234, 1153, 943, 920, 849, 812$  cm<sup>–1</sup>; The enantioselectivity was determined by HPLC (Chiralcel OD-H, eluent: *n*-hexane/*i*-PrOH = 95:5, 0.5 mL/min, 10 °C, retention times: (+)-enantiomer 12.2 min, (–)-enantiomer 15.3 min); HRMS (ESI): *m/z* calcd for C<sub>23</sub>H<sub>27</sub>NO<sub>4</sub>: 382.2013 [M + H]<sup>+</sup>; found: 382.2013.

**(S)-(–)-13b**—Obtained as a colourless oil in 66% yield and with e.r. = 75:25 upon reacting Schiff base **4b** with acrylate **12a** in the presence of 10 mol% catalyst at 0 °C under conditions A. Analytical data are in full accordance with those reported in literature [54].  $[\alpha]_D^{20}$  (c = 0.22, CHCl<sub>3</sub>) = –34.4°; <sup>1</sup>H-NMR (δ, CDCl<sub>3</sub>, 298 K): 2.23–2.32 (m, 2H), 2.33–2.40 (m, 2H), 3.57 (s, 3H), 4.14 (t, *J* = 6.0 Hz, 1H), 5.16 (dd, *J* = 7.0 Hz, 12.5 Hz, 2H), 7.08–7.14 (m, 2H), 7.28–7.42 (m, 11H), 7.59–7.67 (m, 2H) ppm; <sup>13</sup>C-NMR (δ, CDCl<sub>3</sub>, 298 K): 28.5, 30.4, 51.5, 64.1, 66.6, 127.8, 128.0, 128.1, 128.2, 128.5, 128.7, 128.9, 130.5, 135.8, 136.1, 171.4 (2×), 173.4 ppm; IR (film):  $\nu = 3063, 2959, 2928, 2853, 1742, 1705, 1659, 1599, 1578, 1499, 1449, 1420, 1389, 1377, 1317, 1279, 1209, 1192, 1177, 1157, 922, 754, 706$  cm<sup>–1</sup>; The enantioselectivity was determined by HPLC (Chiralcel OD-H, eluent: *n*-hexane/*i*-PrOH = 99:1, 0.5 mL/min, 10 °C, retention times: (+)-enantiomer 68.4 min, (–)-

enantiomer 77.2 min); HRMS (ESI):  $m/z$  calcd for  $C_{26}H_{25}NO_4$ : 416.1856  $[M+H]^+$ ; found: 416.1858.

**(-)-13c**—Obtained as a colourless oil in 71% yield and with e.r. = 90:10 upon reacting Schiff base **4c** with acrylate **12a** using 20 mol% catalyst at  $-20\text{ }^\circ\text{C}$  under conditions A. Analytical data are in full accordance with those reported in literature [57].  $[\alpha]_D^{20}$  ( $c = 0.20$ ,  $CHCl_3$ ) =  $-53.0^\circ$ ;  $^1H$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 2.20–2.29 (m, 2H), 2.32–2.40 (m, 2H), 3.58 (s, 3H), 3.71 (s, 3H), 4.13 (t,  $J = 6.2$  Hz, 1H), 7.14–7.21 (m, 2H), 7.29–7.48 (m, 6H), 7.60–7.67 (m, 2H) ppm;  $^{13}C$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 28.6, 30.4, 51.6, 52.2, 64.1, 127.8, 128.1, 128.6, 128.8, 128.9, 130.6, 172.0, 172.2, 173.4 ppm; IR (film):  $\nu = 3057, 3051, 2992, 2955, 1736, 1624, 1576, 1445, 1437, 1316, 1265, 1204, 1172, 1074, 1028, 1001, 781, 731, 702$   $cm^{-1}$ ; The enantioselectivity was determined by HPLC (Chiralcel OD-H, eluent: *n*-hexane/*i*-PrOH = 99:1, 0.5 mL/min,  $10\text{ }^\circ\text{C}$ , retention times: (+)-enantiomer 48.5 min, (–)-enantiomer 62.4 min); HRMS (ESI):  $m/z$  calcd for  $C_{23}H_{21}NO_4$ : 340.1543  $[M+H]^+$ ; found: 340.1543.

**(-)-13d**—Obtained as a colourless oil in 68% yield and with e.r. = 89:11 upon reacting Schiff base **4c** with acrylate **12b** using 20 mol% catalyst at  $-20\text{ }^\circ\text{C}$  under conditions A.  $[\alpha]_D^{20}$  ( $c = 0.67$ ,  $CHCl_3$ ) =  $-50.8^\circ$ ;  $^1H$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 0.90 (t,  $J = 7.3$  Hz, 3H), 1.28–1.38 (m, 2H), 1.48–1.59 (m, 2H), 2.20–2.29 (m, 2H), 2.30–2.38 (m, 2H), 3.71 (s, 3H), 3.98 (t,  $J = 6.7$  Hz, 2H), 4.12 (t,  $J = 5.9$  Hz, 1H), 7.15–7.21 (m, 2H), 7.29–7.48 (m, 6H), 7.61–7.67 (m, 2H) ppm;  $^{13}C$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 13.7, 19.1, 28.6, 30.6, 30.6, 52.2, 64.2, 64.4, 127.8, 128.1, 128.6, 128.8, 128.9, 130.5, 136.1, 139.3, 171.2, 172.2, 173.0 ppm; IR (film):  $\nu = 3057, 2957, 2934, 2872, 1732, 1661, 1624, 1597, 1578, 1491, 1447, 1437, 1393, 1364, 1317, 1265, 1204, 1175, 1074, 1028, 1001, 941, 920, 781, 764, 737$   $cm^{-1}$ ; The enantioselectivity was determined by HPLC (Chiralcel OD-H, eluent: *n*-hexane/*i*-PrOH = 99:1, 0.5 mL/min,  $10\text{ }^\circ\text{C}$ , retention times: (+)-enantiomer 23.4 min, (–)-enantiomer 24.6 min); HRMS (ESI):  $m/z$  calcd for  $C_{23}H_{27}NO_4$ : 382.2013  $[M+H]^+$ ; found: 382.2015.

**(-)-13f**—Obtained as a colourless oil in 81% yield and with e.r. = 87:13 upon reacting Schiff base **4c** with acrylate **12d** using 20 mol% catalyst at  $-20\text{ }^\circ\text{C}$  under conditions A.  $[\alpha]_D^{20}$  ( $c = 0.63$ ,  $CHCl_3$ ) =  $-45.9^\circ$ ;  $^1H$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 2.23–2.32 (m, 2H), 2.38–2.45 (m, 2H), 3.71 (s, 3H), 4.14 (t,  $J = 6.0$  Hz, 1H), 5.02 (s, 2H), 7.13–7.19 (m, 2H), 7.28–7.46 (m, 11H), 7.61–7.67 (m, 2H) ppm;  $^{13}C$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 28.6, 30.6, 52.2, 64.1, 66.3, 127.8, 128.1, 128.2, 128.5, 128.6, 128.8, 128.9, 130.6, 135.9, 136.1, 172.1, 172.8 ppm; IR (film):  $\nu = 3059, 3036, 2951, 1732, 1659, 1622, 1597, 1578, 1491, 1447, 1420, 1385, 1316, 1265, 1206, 1159, 1074, 1028, 1001, 988, 974, 962, 943, 922, 912, 847, 735$   $cm^{-1}$ ; The enantioselectivity was determined by HPLC (Chiralcel OD-H, eluent: *n*-hexane/*i*-PrOH = 99:1, 0.5 mL/min,  $10\text{ }^\circ\text{C}$ , retention times: (+)-enantiomer 76.3 min, (–)-enantiomer 83.9 min); HRMS (ESI):  $m/z$  calcd for  $C_{26}H_{25}NO_4$ : 416.1856  $[M+H]^+$ ; found: 416.1861.

**(-)-16a**—Obtained as a colourless oil in 65% yield and with e.r. = 75:25 upon reacting Schiff base **4a** with acrylamide **14a** under conditions B.  $[\alpha]_D^{20}$  ( $c = 0.24$ ,  $CHCl_3$ ) =  $-15.8^\circ$ ;  $^1H$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 0.80 (s, 9H), 2.06–2.20 (m, 2H), 2.21–2.43 (m, 2H), 3.93 (dd,  $J = 5.6, 6.35$  Hz, 1H), 7.04–7.13 (m, 2H), 7.17–7.40 (m, 6H), 7.52–7.72 (m, 2H) ppm;  $^{13}C$ -NMR ( $\delta$ ,  $CDCl_3$ , 298 K): 28.1, 29.3, 29.6, 35.4, 37.3, 65.0, 81.1, 127.7, 128.0,

128.5, 128.6, 128.8, 130.3, 136.5, 139.6, 170.4, 171.2, 172.6 ppm; IR (film):  $\nu = 2960, 2880, 2560, 1720, 1640, 1520, 1440, 1400, 1360, 1320, 1280, 1160, 1080, 880 \text{ cm}^{-1}$ ; The enantioselectivity was determined by HPLC (Chiralcel OD-R, eluent: AcN/H<sub>2</sub>O = 55:45, 0.7 mL/min, 10 °C, retention times: (+)-enantiomer 11.6 min, (–)-enantiomer 13.1 min); HRMS (ESI):  $m/z$  calcd for C<sub>24</sub>H<sub>30</sub>N<sub>2</sub>O<sub>3</sub>: 395.2329 [M+H]<sup>+</sup>; found: 395.2325.

**S-(–)-17**—Obtained as a colourless oil in 97% yield and with e.r. = 77:23 upon reacting Schiff base **4c** with MVK (**15**) using 20 mol% catalyst at –20 °C under conditions A.  $[\alpha]_{\text{D}}^{20}$  ( $c = 0.46, \text{CHCl}_3$ ) = –34.1°; <sup>1</sup>H-NMR ( $\delta, \text{CDCl}_3, 298 \text{ K}$ ): 2.11 (s, 3H), 2.13–2.21 (m, 2H), 2.48–2.55 (m, 2H), 3.71 (s, 3H), 4.11 (t,  $J = 6.1 \text{ Hz}$ , 1H), 7.14–7.19 (m, 2H), 7.30–7.48 (m, 6H), 7.61–7.66 (m, 2H) ppm; <sup>13</sup>C-NMR ( $\delta, \text{CDCl}_3, 298 \text{ K}$ ): 27.6, 29.9, 39.7, 52.2, 64.0, 127.7, 128.1, 128.6, 128.8, 128.9, 130.5, 136.1, 139.3, 172.4 (2×), 208.0 ppm; IR (film):  $\nu = 3080, 3055, 2953, 2930, 2173, 1736, 1714, 1659, 1622, 1599, 1578, 1491, 1447, 1437, 1358, 1317, 1275, 1265, 1204, 1177, 1161, 1094, 1074, 1042, 1028, 1001, 941, 920, 810, 783, 766, 733, 698 \text{ cm}^{-1}$ ; The enantioselectivity was determined by HPLC (Chiralcel OD-H, eluent: *n*-hexane/*i*-PrOH = 99:1, 0.5 mL/min, 10 °C, retention times: (+)-enantiomer 69.0 min, (–)-enantiomer 84.7 min); HRMS (ESI):  $m/z$  calcd for C<sub>20</sub>H<sub>21</sub>NO<sub>3</sub>: 324.1594 [M+H]<sup>+</sup>; found: 324.1593.

## 4. Conclusions

Summarizing, the herein developed late stage acetal-transformation strategy did not allow us to obtain novel catalysts **3** in a reliable and straightforward fashion especially due to problems associated with the catalyst counter anion and the hereby formed hardly removable impurities. To get a detailed understanding of the application scope and limitations of our catalysts we tested them in a variety of different important transformations and found that, although these compounds have recently shown their good potential in the asymmetric  $\alpha$ -alkylation of glycine Schiff bases, they clearly failed when we attempted to control more reactive nucleophiles like  $\beta$ -keto esters. On the other hand using them to catalyse the Michael addition of glycine Schiff bases to different acceptors very interesting results have been obtained. It was found necessary to carefully optimize the reaction conditions for every single substrate class, as seemingly small structural changes required the use of totally different reaction conditions. Unfortunately, the strikingly different behaviour of different nucleophiles and different electrophiles and also the need for totally different reaction conditions compared to the standard alkylation reaction is not fully understood yet. This highlights again the necessity of carrying out careful screening studies and the problems of transferring knowledge gathered in one test system to another one, especially in complex heterogeneous reaction systems as usually employed in asymmetric phase-transfer catalysis. In addition, we observed again a very strong influence of the counter anions on the catalyst performance, thus making a strict control of the anion necessary. Under carefully optimized conditions enantiomeric ratios of up to 91:9 could be achieved in the addition of glycine Schiff bases to acrylates whereas acrylamides and methyl vinyl ketone were less well tolerated (up to e.r. 77:23 in these cases). Accordingly, we have now a rather detailed understanding about the scope and limitations of the synthesis sequence to access our PTCs and about the application scope of these catalysts in asymmetric transformations.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


This work was supported by the Austrian Science Funds (FWF): Project No. P22508-N17. Richard Herchl is recipient of a DOC-fellowship of the Austrian Academy of Sciences at the Institute of Organic Chemistry, JKU Linz. We are grateful to Christian Rückl for his support with the HPLC analysis. The used NMR spectrometers were acquired in collaboration with the University of South Bohemia (CZ) with financial support from the European Union through the EFRE INTERREG IV ETC-AT-CZ programme (project M00146, "RERI-uasb").

## References



1. O'Donnell MJ. The enantioselective synthesis of  $\alpha$ -amino acids by phase-transfer catalysis with achiral schiff base esters. *Acc. Chem. Res.* 2004; 37:506–517. [PubMed: 15311949]
2. Lygo B, Andrews BI. Asymmetric phase-transfer catalysis utilizing chiral quaternary ammonium salts: Asymmetric alkylation of glycine imines. *Acc. Chem. Res.* 2004; 37:518–525. [PubMed: 15311950]
3. Hashimoto T, Maruoka K. Recent development and application of chiral phase-transfer catalysis. *Chem. Rev.* 2007; 107:5656–5682. [PubMed: 18072806]
4. Ooi T, Maruoka K. Recent advances in asymmetric phase-transfer catalysis. *Angew. Chem. Int. Ed. Engl.* 2007; 46:4222–4266. [PubMed: 17525926]
5. Shirakawa S, Maruoka K. Recent developments in asymmetric phase-transfer reactions. *Angew. Chem. Int. Ed. Engl.* 2013 doi:10.1002/anie.201206835.
6. Novacek J, Waser M. Bifunctional chiral quaternary ammonium salt catalysts: A rapidly emerging class of powerful asymmetric catalysts. *Eur. J. Org. Chem.* 2013; 2013:637–648.
7. Helder R, Hummelen JC, Laane RWPM, Wiering JS, Wynberg H. Catalytic asymmetric induction in oxidation reactions. The synthesis of optically active epoxides. *Tetrahedron Lett.* 1976; 17:1831–1834.
8. Dolling UH, Davis P, Grabowski EJJ. Efficient catalytic asymmetric alkylations. 1. Enantioselective synthesis of (+)-indacrinone via chiral phase-transfer catalysis. *J. Am. Chem. Soc.* 1984; 106:446–447.
9. O'Donnell MJ, Bennett WD, Wu S. The stereoselective synthesis of  $\alpha$ -amino acids by phase-transfer catalysis. *J. Am. Chem. Soc.* 1989; 111:2353–2355.
10. O'Donnell MJ, Wu S, Huffman JC. A new active catalyst species for enantioselective alkylation by phase-transfer catalysis. *Tetrahedron.* 1994; 50:4507–4518.
11. Lygo B, Wainwright PG. A new class of asymmetric phase-transfer catalysts derived from *Cinchona* alkaloids—Application in the enantioselective synthesis of  $\alpha$ -amino acids. *Tetrahedron Lett.* 1997; 38:8595–8598.
12. Lygo B, Crosby J, Peterson JA. Enantioselective synthesis of bis- $\alpha$ -amino acid esters *via* asymmetric phase-transfer catalysis. *Tetrahedron Lett.* 1999; 40:1385–1388.
13. Corey EJ, Xu F, Noe MC. A rational approach to catalytic enantioselective enolate alkylation using a structurally rigidified and defined chiral quaternary ammonium salt under phase transfer conditions. *Am. Chem. Soc.* 1997; 119:12414–12415.
14. Corey EJ, Noe MC, Xu F. Highly enantioselective synthesis of cyclic and functionalized  $\alpha$ -amino acids by means of a chiral phase transfer catalyst. *Tetrahedron Lett.* 1998; 39:5347–5350.
15. Liu Y, Provencher BA, Bartelson KJ, Deng L. Highly enantioselective asymmetric Darzens reactions with a phase transfer catalyst. *Chem. Sci.* 2011; 2:1301–1304. [PubMed: 22174974]
16. Provencher BA, Bartelson KJ, Liu Y, Foxman BM, Deng L. Structural study-guided development of versatile phase-transfer catalysts for asymmetric conjugate additions of cyanide. *Angew. Chem. Int. Ed. Engl.* 2011; 50:10565–10569. [PubMed: 21919177]

17. Bella M, Kobbelgaard S, Jørgensen KA. Organocatalytic regio- and asymmetric C-selective  $S_NAr$  reactions—stereoselective synthesis of optically active spiro-pyrrolidone-3,3'-oxoindoles. *J. Am. Chem. Soc.* 2005; 127:3670–3671. [PubMed: 15771481]
18. Poulsen TB, Bernardi L, Aleman J, Overgaard J, Jørgensen KA. Organocatalytic asymmetric direct  $\alpha$ -alkynylation of cyclic  $\beta$ -ketoesters. *J. Am. Chem. Soc.* 2007; 129:441–449. [PubMed: 17212424]
19. Maciver EE, Knipe PC, Cridland AP, Thompson AL, Smith MD. Catalytic enantioselective electrocyclic cascades. *Chem. Sci.* 2012; 3:537–540.
20. Bernardi L, Indrigo E, Pollicino S, Ricci A. Organocatalytic trifluoromethylation of imines using phase-transfer catalysis with phenoxides. A general platform for catalytic additions of organosilanes to imines. *Chem. Commun.* 2012; 48:1428–1430.
21. Fiandra CD, Piras L, Fini F, Disetti P, Moccia M, Adamo MFA. Phase transfer catalyzed enantioselective cyclopropanation of 4-nitro-5-styrylisoxazoles. *Chem. Commun.* 2012; 48:3863–3865.
22. Johnson KM, Rattley MS, Sladojevich F, Barber DM, Nunez MG, Goldys AM, Dixon DJ. A new family of cinchona-derived bifunctional asymmetric phase-transfer catalysts: Application to the enantio- and diastereoselective nitro-Mannich reaction of amidosulfones. *Org. Lett.* 2012; 14:2492–2495. [PubMed: 22548631]
23. Bernal P, Fernández R, Lassaletta JM. Organocatalytic asymmetric cyanosilylation of nitroalkenes. *Chem. Eur. J.* 2010; 16:7714–7718. [PubMed: 20533468]
24. Haraguchi N, Ahamed P, Parvez M, Itsuno S. Synthesis of main-chain chiral quaternary ammonium polymers for asymmetric catalysis using quaternization polymerization. *Molecules.* 2012; 17:7569–7583. [PubMed: 22713348]
25. Hintermann L, Dittmer C. Asymmetric ion-pairing catalysis of the reversible cyclization of 2-hydroxychalcone to flavanone: Asymmetric catalysis of an equilibrating reaction. *Eur. J. Org. Chem.* 2012:5573–5584.
26. Herchl R, Waser M. Asymmetric cyclopropanation of chalcones using chiral phase-transfer catalysts. *Tetrahedron Lett.* 2013; 54:2472–2475. [PubMed: 24391288]
27. Kamlar M, Putaj P, Vesely J. Organocatalytic alkynylation of densely functionalized monofluorinated derivatives: C(sp<sup>3</sup>)–C(sp) coupling. *Tetrahedron Lett.* 2013; 54:2097–2100.
28. Denmark SE, Weintraub RC. Deconstructing quinine. Part 1. Toward an understanding of the remarkable performance of cinchona alkaloids in asymmetric phase-transfer catalysis. *Heterocycles.* 2011; 82:1527–1540.
29. Tanzer E-M, Schweizer WB, Ebert M-O, Gilmour R. Designing fluorinated cinchona alkaloids for enantioselective catalysis: Controlling internal rotation by a fluorine-ammonium ion gauche effect ( $\phi$ NCCF). *Chem. Eur. J.* 2012; 18:2006–2013. [PubMed: 22250061]
30. Ooi T, Kameda M, Maruoka K. Molecular design of a C<sub>2</sub>-symmetric chiral phase-transfer catalyst for practical asymmetric synthesis of  $\alpha$ -amino acids. *J. Am. Chem. Soc.* 1999; 121:6519–6520.
31. Shirakawa S, Liu K, Ito H, Maruoka K. Catalytic asymmetric synthesis of 1,1-disubstituted tetrahydro- $\beta$ -carboline by phase-transfer catalyzed alkylations. *Chem. Commun.* 2011; 47:1515–1517.
32. Kano T, Yamamoto A, Song S, Maruoka K. Catalytic asymmetric syntheses of isoxazoline-N-oxides under phase-transfer conditions. *Chem. Commun.* 2011; 47:4358–4360.
33. Hashimoto T, Sakata K, Maruoka K. Phase-transfer-catalyzed olefin isomerization/ $\alpha$ -alkylation of  $\alpha$ -alkynylcrotonates as a route for 1,4-enynes. *Adv. Synth. Catal.* 2010; 352:1653–1656.
34. Lan Q, Wang X, Shirakawa S, Maruoka K. Phase-transfer catalyzed asymmetric conjugate additions of  $\beta$ -ketoesters to acetylenic ketones. *Org. Process Res. Dev.* 2010; 14:684–686.
35. Shirakawa S, Liu K, Maruoka K. Catalytic asymmetric synthesis of axially chiral *o*-iodoanilides by phase-transfer catalyzed alkylations. *J. Am. Chem. Soc.* 2012; 134:916–919. [PubMed: 22208662]
36. Shibuguchi T, Fukuta Y, Akachi Y, Sekine A, Ohshima T, Shibasaki M. Development of new asymmetric two-center catalysts in phase-transfer reactions. *Tetrahedron Lett.* 2002; 43:9539–9543.
37. Okada A, Shibuguchi T, Ohshima T, Masu H, Yamaguchi K, Shibasaki M. Enantio- and diastereoselective catalytic Mannich-type reaction of a glycine schiff base using a chiral two-


- center phase-transfer catalyst. *Angew. Chem. Int. Ed. Engl.* 2005; 44:4564–4567. [PubMed: 15954147]
38. Shibuguchi T, Mihara H, Kuramochi A, Ohshima T, Shibasaki M. Catalytic asymmetric phase-transfer Michael reaction and Mannich-type reaction of glycine schiff bases with tartrate-derived diammonium Salts. *Chem. Asian. J.* 2007; 2:794–801. [PubMed: 17492797]
  39. Lygo B, Allbutta B, James SR. Identification of a highly effective asymmetric phase-transfer catalyst derived from  $\alpha$ -methylnaphthylamine. *Tetrahedron Lett.* 2003; 44:5629–5632.
  40. Lygo B, Allbutta B, Kirton EHM. Asymmetric Michael addition of glycine imines via quaternary ammonium ion catalysis. *Tetrahedron Lett.* 2005; 46:4461–4464.
  41. Arai S, Tsuji R, Nishida A. Phase-transfer-catalyzed asymmetric Michael reaction using newly-prepared chiral quaternary ammonium salts derived from L-tartrate. *Tetrahedron Lett.* 2002; 43:9535–9537.
  42. Kowtoniuk WE, MacFarland DK, Grover GN. Combining chiral elements: A novel approach to asymmetric phase-transfer catalyst design. *Tetrahedron Lett.* 2005; 46:5703–5705.
  43. Kumar S, Ramachandram U. The synthesis and applications of asymmetric phase-transfer catalysts derived from isomannide and isosorbide. *Tetrahedron.* 2005; 61:4141–4148.
  44. Denmark SE, Gould ND, Wolf LM. A systematic investigation of quaternary ammonium ions as asymmetric phase-transfer catalysts. Synthesis of catalyst libraries and evaluation of catalyst activity. *J. Org. Chem.* 2011; 76:4260–4336. [PubMed: 21446721]
  45. Denmark SE, Gould ND, Wolf LM. A systematic investigation of quaternary ammonium ions as asymmetric phase-transfer catalysts. Application of quantitative structure activity/selectivity relationships. *J. Org. Chem.* 2011; 76:4337–4357. [PubMed: 21446723]
  46. Seebach D, Keck AB, Heckel A. TADDOLs, their derivatives, and TADDOL analogues: Versatile chiral auxiliaries. *Angew. Chem. Int. Ed. Engl.* 2001; 40:92–138. [PubMed: 11169693]
  47. Pellissier H. Use of TADDOLs and their derivatives in asymmetric synthesis. *Tetrahedron.* 2008; 64:10279–10317.
  48. Waser M, Gratzner K, Herchl R, Müller N. Design, synthesis, and application of tartaric acid derived *N*-spiro quaternary ammonium salts as chiral phase-transfer catalysts. *Org. Biomol. Chem.* 2012; 10:251–254. [PubMed: 22038391]
  49. Gratzner K, Waser M. Investigations concerning the syntheses of TADDOL-derived secondary amines and their use to access novel chiral organocatalysts. *Synthesis.* 2012; 44:3661–3670.
  50. Chinchilla R, Mazon P, Najera C, Ortega FJ. The counterion effect in the phase-transfer catalyzed asymmetric synthesis of  $\alpha$ -amino acids promoted by anthryl-derived dimeric *Cinchona ammonium* salts. *Tetrahedron Asymmetry.* 2004; 15:2603–2607.
  51. Ohshima T, Gnanadesikan V, Shibuguchi T, Fukuta Y, Nemoto T, Shibasaki M. Enantioselective syntheses of Aeruginosin 298-A and its analogues using a catalytic asymmetric phase-transfer reaction and epoxidation. *J. Am. Chem. Soc.* 2003; 125:11206–11207. [PubMed: 16220936]
  52. Lygo B, Beynon C, Lumley C, McLeod MC, Wade CE. Co-catalyst enhancement of enantioselective PTC Michael additions involving glycine imines. *Tetrahedron Lett.* 2009; 50:3363–3365.
  53. Saito S, Tsubogo T, Kobayashi S. Chiral calcium complexes as Brønsted base catalysts for asymmetric addition of  $\alpha$ -amino acid derivatives to  $\alpha,\beta$ -unsaturated carbonyl compounds. *J. Am. Chem. Soc.* 2007; 129:5364–5365. [PubMed: 17425321]
  54. Shibuguchi T, Mihara H, Kuramochi A, Sakuraba S, Ohshima T, Shibasaki M. Short synthesis of (+)-Cylindricine C by using a catalytic asymmetric Michael reaction with a two-center organocatalyst. *Angew. Chem. Int. Ed. Engl.* 2006; 45:4635–4637. [PubMed: 16789053]
  55. Arai S, Tokumaru K, Aoyama T. Asymmetric Michael reaction promoted by new chiral phase-transfer catalysts. *Chem. Pharm. Bul.* 2004; 52:646–648.
  56. Akiyama T, Hara M, Fuchibe K, Sakamoto S, Yamaguchi K. Synthesis of a novel crown ether derived from *chiro*-inositol and its catalytic activity on the asymmetric Michael addition. *Chem. Commun.* 2003:1734–1735.
  57. Tsubogo T, Saito S, Seki K, Yamashita Y, Kobayashi S. Development of catalytic asymmetric 1,4-addition and [3 + 2] cycloaddition reactions using chiral calcium complexes. *J. Am. Chem. Soc.* 2008; 130:13321–13332. [PubMed: 18783222]



**Figure 1.**  
Privileged chiral ammonium salt PTCs.

**Scheme 1.**

Recently described synthesis of TADDOL-derived *N*-spiro ammonium salt catalysts **3** and their performance in initial test reactions [48,49].



**Representative screening results:**

catalyst **8** (X = CF<sub>3</sub>COO): 19%, *e.r.* 72:28


catalyst **3b** (R<sup>1</sup> = 2-Np, R<sup>2</sup> = Me, X = CF<sub>3</sub>SO<sub>3</sub>): 10%, *e.r.* 62:38

↓ HBr anion exchange

catalyst **3c** (R<sup>1</sup> = 2-Np, R<sup>2</sup> = Me, X = Br): 45%, *e.r.* 91:9

**Scheme 2.**



Late-stage acetal modification of **3a** to access **8** and differently substituted ammonium salts **3** and their catalytic potential.

**Scheme 3.**



Attempted **3a**-catalysed asymmetric  $\alpha$ -alkylation of  $\beta$ -keto esters **10**.

Table 1

Asymmetric Michael addition of glycine Schiff bases **4** to acrylates **12** catalysed by TADDOL-derived ammonium salts **3**.



Entry <sup>a</sup>	Cat. (mol%)	4	12	Solv.	Base (eq.)	T [°C]	13	Yield <sup>b</sup> [%]	e.r. <sup>c</sup> (conf.) <sup>d</sup>
1	3a (10%)	4a	12a	toluene	KOH (50%) (25×)	0	13a	94	51:49 ( <i>S</i> )
2	3a (10%)	4a	12a	toluene	KOH (50%) (1×)	0	13a	88	50:50
3	3a (10%)	4a	12a	toluene	KOH (s) (20×)	0	13a	76	50:50
4	3a (10%)	4a	12a	toluene	K <sub>3</sub> PO <sub>4</sub> (50%) (10×)	0	13a	34	50:50
5	3a (10%)	4a	12a	toluene	K <sub>2</sub> CO <sub>3</sub> (50%) (10×)	0	13a	18	52:48 ( <i>S</i> )
6	3a (10%)	4a	12a	toluene	Cs <sub>2</sub> CO <sub>3</sub> (70%) (10×)	0	13a	10	50:50
7	3a (10%)	4a	12a	toluene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13a	73	66:34 ( <i>S</i> )
8	3a (10%)	4a	12a	toluene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13a	56	71:29 ( <i>S</i> )
9 <sup>e</sup>	3a (10%) <sup>e</sup>	4a	12a	toluene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13a	62	61:39 ( <i>S</i> )
10	3d (10%)	4a	12a	toluene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13a	14	62:38 ( <i>S</i> )
11	3e (10%)	4a	12a	toluene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13a	82	64:36 ( <i>S</i> )
12	3a (10%)	4a	12a	benzene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13a	72	58:42 ( <i>S</i> )
13	3a (10%)	4a	12a	fluorobenzene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13a	89	54:46 ( <i>S</i> )
14	3a (10%)	4a	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13a	74	69:31 ( <i>S</i> )
15 <sup>f</sup>	3a (10%)	4a	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13a	33	57:43 ( <i>S</i> )
16 <sup>g</sup>	3a (10%)	4a	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13a	76	51:49 ( <i>S</i> )



Entry <sup>a</sup>	Cat. (mol%)	4	12	Solv.	Base (eq.)	T [°C]	13	Yield <sup>b</sup> [%]	e.r. <sup>c</sup> (conf.) <sup>d</sup>
17	3a (10%)	4b	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13b	66	75:25 (S)
18	3a (10%)	4c	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	0	13c	81	78:22
19	3a (10%)	4c	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13c	35	85:15
20	3a (20%)	4c	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13c	71	90:10
21	3a (20%)	4c	12b	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13d	68	89:11
22	3a (20%)	4c	12c	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13e	n.r.	n.d.
23	3a (20%)	4c	12d	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13f	81	87:13
24	3f (20%)	4c	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13c	51	80:20
25	3g (20%)	4c	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13c	56	86:14
26	3h (20%)	4c	12a	mesitylene	Cs <sub>2</sub> CO <sub>3</sub> (s) (20×)	-20	13c	68	91:9

<sup>a</sup>22 h reaction time under an Ar-atmosphere using 1.5 equiv. of the acrylate **12**;

<sup>b</sup>Isolated Yield;

<sup>c</sup>Determined by HPLC using a chiral stationary phase. In each case the (-)-enantiomer was the major one;

<sup>d</sup>Determined by comparison of the HPLC retention time and the optical rotation with values reported in literature (**13a** [53]; **13b** [54])

<sup>e</sup>Using recovered catalyst;

<sup>f</sup>Using 4Å molecular sieve as an additive

<sup>g</sup>Using mesitol as an additive.

