

No Association between 25 (OH) Vitamin D Level And Hypothyroidism among Females

Imad R. Musa¹, Gasim I. Gasim^{2*}, Sajjad Khan³, Ibrahim A. Ibrahim⁴, Hamdi Abo-alazm⁴, Ishag Adam⁵

¹Armed Forces Hospital at King Abdu Aziz Air Base, Dhahran, Saudi Arabia; ²Alneelain School of Medicine, Alneelain University, Khartoum, Sudan; ³Central Hospital, Arar, Saudi Arabia; ⁴Medical Laboratory Science, United Medical Laboratory, Saudi Arabia; ⁵Faculty of Medicine, Khartoum University, Khartoum, Sudan

Abstract

Citation: Musa IR, Gasim GI, Khan S, Ibrahim IA, Abo-alazm H, Adam I. No Association between 25 (OH) Vitamin D Level And Hypothyroidism among Females. Open Access Maced J Med Sci. 2017 Apr 15; 5(2):126-130. <https://doi.org/10.3889/oamjms.2017.029>

Keywords: hypothyroidism; vitamin D levels; females.

***Correspondence:** Gasim I. Gasim. Alneelain School of Medicine, Alneelain University, Khartoum, Sudan. E-mail: gasingsm@yahoo.com

Received: 26-Nov-2016; **Revised:** 16-Jan-2017; **Accepted:** 09-Feb-2017; **Online first:** 19-Mar-2017

Copyright: © 2017 Imad R. Musa, Gasim I. Gasim, Sajjad Khan, Ibrahim A. Ibrahim, Hamdi Abo-alazm, Ishag Adam. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Funding: This research did not receive any financial support.

Competing Interests: The authors have declared that no competing interests exist.

AIM: The aim was to investigate serum vitamin D (25-OH) level among females with hypothyroidism.

MATERIALS AND METHODS: A case-control study (58 in each arm) was conducted in Arar Central Hospital, Kingdom Saudi Arabia. The cases were females with hypothyroidism, and healthy females were controls. TSH, thyroid hormones: Free T3 (FT3) and Free T4 (FT4) and haemoglobin levels were measured in all participants. Serum vitamin D (25-OH) level was measured using the spectrophotometry.

RESULTS: While there was no significant difference in the age and haemoglobin level, body mass index (BMI) was significantly higher in the cases. Compared with the controls, cases had significantly higher TSH, had significantly lower T4, and there was no significant difference in FT3 and 25 (OH) vitamin D, [16.1 (8.8-26.7) vs. 14.0 (9.5-20.3 ng/ml; P = 0.577]. Linear regression showed no association between, age, BMI, haemoglobin, TSH, FT3, FT4 and the log of 25 (OH) vitamin D levels.

CONCLUSION: There was no significant difference in vitamin D level among females with hypothyroidism and healthy controls.

Introduction

Vitamin D deficiency is a global health problem where over a billion people worldwide have some vitamin D deficiency or insufficiency [1, 2]. The essential biological action of vitamin D is the regulation of calcium and phosphorus metabolism. Though the role of vitamin D in the regulation of immune system has recently been reported [3, 4]. Hypothyroidism is a common disease in the general population, most of which results as consequences of Hashimoto's thyroiditis with a tendency to increase with age and female are at 8 to 15 times at higher risk than men [5]. It has been observed that serum Vitamin D3 metabolites are elevated in hypothyroid subjects [6].

However, a reduced level has been reported in a patient with over-active thyroid gland [7]. Thyroid disorders may affect vitamin D status as subnormal

levels of vitamin D were observed in a patient with Graves' disease [8]. The role of vitamin D is explored as an anti-inflammatory, anti-proliferative factor and association with many clinical disorders [9, 10]. Vitamin D has a direct and indirect role in inhibiting the production of cytokines (IL-12 and IFN- γ), hence, enhances the immune and anti-inflammatory responses [11].

Vitamin D is circulating in the blood attached to a D-binding protein. Then hydroxylation takes place at the liver and the kidneys to produce 25(OH) D and the active metabolite, 1, 25 dihydroxy vitamin D (1, 25-(OH) 2 D) respectively [12]. The active form of vitamin D is called calcitriol or 1, 25-dihydroxyvitamin D3. Serum 25(OH) D is used to reflect vitamin D status because it is the major circulating precursor of active vitamin D and has a half-life, ranging from two to three weeks.

Interestingly, 1, 25-(OH) vitamin D has a short circulating half-life and is tightly under the influence of

parathyroid hormone, calcium and phosphate [13]. Moreover, low level of serum 1, 25-(OH) 2D may not observe until vitamin D deficiency is severe [14]. Hence 25(OH) level is used to assess vitamin D reserve. Vitamin D has a vital role in many physiological actions and an association with some clinical disorders.

Hence the current study was conducted to investigate serum vitamin D (25-OH) level among Saudi females with hypothyroidism.

Material and Methods

A case- control study was conducted at Arar Central Hospital, Kingdom of Saudi Arabia during the period 1st of November 2014 to 1st of May 2015. Cases were women with hypothyroidism. Moreover, those who had symptoms suggestive of hypothyroidism e.g. fatigue, cold intolerance, constipation, weight gain, dry skin and menstrual disturbance or those who had sounding features: puffy, hoarseness, depressed mood and bradycardia were screened and enrolled as new cases if proofed to have hypothyroidism. The cases were diagnosed by the high level of TSH (≥ 6.30 mIU/L) with lower levels of FT3 and FT4 than normal value. The controls were healthy females. Those with chronic medical diseases, on vitamin D supplementation, children, pregnant females and non-Saudi citizen, were excluded.

Ethical approval was obtained from medical education and research centre at Arar Central Hospital, KSA.

After signing an informed consent, height and weight were measured using the conventional method and were used to calculate the body mass index (BMI) by using the formula: weight in kilogrammes (kg) divided by height in square meters (m^2).

Blood (5 ml) was withdrawn, kept for 30 minutes at the room temperature to clot and centrifuged for five minutes. Then TSH, FT3 and FT4 were measured using commercially available kits (Roche Diagnostics, Mannheim, Germany). Reference values for normal range of TSH (0.48 – 6.30 mIU/l), FT3 (3.39-5.82, pmol/l) and FT4 (9.00-17.15, pmol/l) [15].

The 25-hydroxy vitamin D (25 (OH) vitamin D) was measured using spectrophotometry. The normal level of 25 (OH) D is 30 ng/ml or above [16]. Haemoglobin (Hb) level was checked too.

Fifty-eight participants in each arm of the study were calculated as to have the difference in the mean level of thyroid hormones and 25 (OH) vitamin D between the two groups and to have over 80%

power to detect a difference of 5% at $\alpha = 0.05$. We assumed that 10% of participants might not respond or have incomplete data.

Statistical analysis

Data was analysed using the Statistical Package for the Social Sciences (SPSS) version 22 (Chicago, IL, USA). Chi-square test was used to compare proportions between the two groups. Kolmogorov –Smirnov test was used for testing the normality of the data. Student's t-test and Mann-Whitney test were used to compare the continuous parametric and non-parametric (thyroid hormones and 25 (OH) vitamin D) data, respectively between the two groups. Simple linear regression models with the log of 25 (OH) vitamin D as a continuous dependent variable was used. Socio-demographic characteristics, BMI, and thyroid hormones levels were the independent predictors of interest. A p value of <0.05 was considered significant.

Results

While there was no significant difference in the mean (SD) of age [34.1 (13.2)] vs. [37.3 (10.2) years, $P = 0.140$] and haemoglobin [12.2 (1.4) vs. 12.6 (1.4) g/dl, $P = 0.192$] levels between the two groups, BMI was significantly higher in the cases compared with the controls, [33.2 (6.2) vs. 30.2 (6.1, kg/m^2 , $P = 0.01$], Table 1.

Table 1: Comparing the mean (SD) of the age, body mass index and haemoglobin level between the cases and the controls

Variables	Cases (n= 58)	Control (n= 58)	P
Age, years	34.1 (13.2)	37.3 (10.2)	0.140
Body mass index, kg/m^2	33.2 (6.2)	30.2 (6.1)	0.010
Haemoglobin, g/dl	12.2 (1.4)	12.6 (1.4)	0.192

Compared with the controls, cases had significantly higher TSH, had significantly lower FT4, and there was no significant difference in FT3 (4.7 (3.8 -5.4) vs. 4.5 (3.9 – 5.3) pmol/l, $P = 0.685$) and 25 (OH) vitamin D, 16.1(8.8– 26.7) vs. 14.0 (9.5– 20.3ng/ml; $P = 0.577$], Table 2, Figure 1.

Table 2: Median (interquartile) of thyroid hormones and 25 (OH) vitamin D level in the cases and controls

Variables	Cases (n= 58)	Control (n= 58)	P
TSH, mIU/l	4.9 (2.0 – 9.5)	2.2 (1.1– 3.4)	< 0.001
FT3, pmol/l	4.7 (3.8 – 5.4)	4.5 (3.9 – 5.3)	0.685
FT4, pmol/l	13.3 (11.1–15.0)	14.0 (12.0– 15.7)	0.028
25 (OH) vitamin D, ng/ml	16.1(8.8–26.7)	14.0 (9.5–20.3)	0.577

Linear regression showed no association between, age, BMI, haemoglobin, TSH, T3, T4 and the log of 25 (OH) vitamin D levels, Table3.




Figure 1: Median (interquartile) of 25 (OH) vitamin D level in the cases and the controls

Discussion

The main finding of the current study was a non-significant difference in the level of vitamin D among the women with hypothyroid compared to the control. This goes with Goswami et al., findings where they observed no significant difference between autoimmune thyroid disorders (AITD) and vitamin D levels [17]. Likewise, a non-significant association between vitamin D deficiency and autoimmune thyroid disorders (AITD) has been previously reported [18]. The aforementioned observation was further consolidated by a similar finding in hypothyroidism as the Hashimoto thyroiditis induced and Non-Hashimoto thyroiditis [18, 19].

Table 3: Factors affecting log of 25 (OH) vitamin D level using multiple linear regression analysis

Variables	Coefficient	Standard error	P
Age, years	0.003	0.003	0.189
Body mass index, kg.m ²	0.002	0.005	0.608
Haemoglobin, g/dl	0.013	0.021	0.525
TSH, mIU/l	-0.002	0.008	0.814
FT3, pmol/l	0.000	0.034	0.998
FT4, pmol/l	-0.001	0.011	0.913
Case (hypothyroidism) versus controls	0.023	0.068	0.741

The non-significant difference of low vitamin D level among both groups may be explained by the higher prevalence (61.5% to 83.6%) of vitamin D deficiency among Saudi population perhaps due and inadequate exposure to sensible sunlight or females are wearing black outer cloak which may reduce the benefit of sensible sunlight [20-22].

A significant correlation between vitamin D and hypothyroidism among female participants has recently been documented [23, 24]. Moreover, a

recent study showed, the serum 25-OH vitamin D levels were significantly lower in a group of hypothyroid cases compared to healthy controls [25]. Some different studies documented a significant negative association between vitamin D levels in patients with Hashimoto's thyroiditis compared to the control group [1, 6, 26-31]

Interestingly, significantly high levels of vitamin D were found in cases with subclinical hypothyroidism compared to a healthy control group in a similar study that was conducted in Saudi Arabia with the majority of cases being females: female (39) vs. males (3) [32].

The current study showed a significant difference in BMI between the cases and controls. This goes with one study conducted in the Kingdom of Saudi Arabia that showed similar findings [32]. In a similar context, another two studies proposed that vitamin D deficiency can predict BMI [33, 34].

In this study, the median (interquartile) of serum thyroid stimulating hormone (TSH) levels and FT4 levels showed a significant difference between the two groups. This goes with the previous finding in one study that was conducted on the same nationality of participants [35]. Serum 25(OH)D levels were found to be significantly negatively correlated with serum thyroid stimulating hormone (TSH) levels and positively with T4 in some recent clinical data [36-38].

The key role in the development of autoimmune process is genetically dependent on aberrant expression of HLA-DR and other antigens on the thyrocyte surface hence are become vulnerable to an autoimmune insult [39]. Likewise, a different gene in the Vitamin D receptor was found to be the culprit to autoimmune thyroid disease such as Graves' disease and Hashimoto's thyroiditis [26, 36]. An indirect effect can be due to a higher serum level of calcium, low level of circulating parathyroid hormone and high phosphorous levels which may lead to suppression of renal 25(OH) D1- α hydroxylase activity. Therefore 1, 25 (OH) 2D productions will be reduced and this will be accompanied with marked elevation in both serum 24, 25(OH) and serum thyroid hormone levels [6]. It has to be mentioned that the beneficial effects of vitamin D could be due to prevention/correction of the potential risk of hypothyroidism and an improvement in the thyroid functions [27, 40-42]. Also correction of vitamin D with anti-thyroid drugs or thyroid hormone replacement is associated with better treatment response, besides, suppressing the autoimmune reaction as reflected by reducing serum levels of thyroid autoantibodies [43]. Furthermore, vitamin D supplementation is associated with prevention and development of different autoimmune diseases in both humans and animal models [27]. Both vitamin D and thyroid hormone bind to analogous receptors called steroid hormone receptors. Tissues express vitamin D receptor (VDR) and 1-alpha-hydroxylase, have a paracrine and autocrine effects to allow upregulation

of calcitriol production, neither depends on parathormone nor renal influence. [6, 39] Mutation of genes encoding for tissue hydroxylases and VDR hire a risk for development of thyroid cancer. [44, 45] Low levels of vitamin D (1, 25-(OH) 2D3) were observed in different types of thyroid cancers and correlated with cancer staging (I-IV) [45]

On the other hand, low levels of vitamin D may be related to hypothyroidism disorder. We can justify the presence of low levels of vitamin D among subjects with hypothyroidism for poor intestinal absorption in some autoimmune disorders that are associated with hypothyroidism such as celiac disease and pernicious anaemia. Moreover, the seasonal variation may affect the prevalence as one report documented higher rates of low serum level of the vitamin during winter compared to spring [46]. The seasonal variation of vitamin D was not one of the objectives of the current study. On the contrary, a lower prevalence of vitamin D deficiency among males compared with females was previously observed, when gender issue was considered [22, 36].

One of the limitations of this study is; calcium, phosphorous, parathyroid hormones and thyroid antibodies were not investigated.

In conclusion, there was no significant difference in vitamin D level among females with hypothyroidism and healthy controls.

References

- Holick MF. Vitamin D Deficiency. *New England Journal of Medicine*. 2007; 357: 266–281. <https://doi.org/10.1056/NEJMra070553> PMID:17634462
- Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *The American journal of clinical nutrition*. 2008; 87:1080S–6S. PMID:18400738
- Lemire JM, Adams JS, Sakai R, Jordan SC. 1 alpha,25-dihydroxyvitamin D3 suppresses proliferation and immunoglobulin production by normal human peripheral blood mononuclear cells. *Journal of Clinical Investigation*. 1984;74(2):657-661. <https://doi.org/10.1172/JCI111465> PMID:6611355 PMID:PMC370520
- Rigby WF, Stacy T, Fanger MW. Inhibition of T lymphocyte mitogenesis by 1, 25-dihydroxyvitamin D3 (calcitriol). *The Journal of clinical investigation*. 1984; 74:1451–5. <https://doi.org/10.1172/JCI111557> PMID:6332829 PMID:PMC425314
- Kostoglou-Athanassiou I, Ntalles K. Hypothyroidism - new aspects of an old disease. *Hippokratia*. 2010;14(2):82-87. PMID:20596261 PMID:PMC2895281
- Mosekilde L, Lund B, Sørensen OH, Christensen MS, Melsen F. Serum-25 hydroxycholecalciferol in hyperthyroidism. *The Lancet*. 1977; 309:806–807. [https://doi.org/10.1016/S0140-6736\(77\)92995-6](https://doi.org/10.1016/S0140-6736(77)92995-6)
- Velentzas C, Oreopoulos D.G, From G, Porret B, Rapoport A. Vitamin-d levels in thyrotoxicosis. *The Lancet*. 1977; 309:370–371. [https://doi.org/10.1016/S0140-6736\(77\)91177-1](https://doi.org/10.1016/S0140-6736(77)91177-1)
- Alhuzaim ON, Aljohani N. Effect of Vitamin D3 on Untreated Graves' Disease with Vitamin D Deficiency. *Clinical Medicine Insights Case Reports*. 2014; 7:83-85. <https://doi.org/10.4137/CCRep.S13157> PMID:25187748 PMID:PMC4133032
- Lee JH, O'Keefe JH, Bell D, Hensrud DD, Holick MF. Vitamin D deficiency an important, common, and easily treatable cardiovascular risk factor? *Journal of the American College of Cardiology*. 2008;52:1949–56. <https://doi.org/10.1016/j.jacc.2008.08.050> PMID:19055985
- Melamed ML, Michos ED, Post W, Astor B. 25-hydroxyvitamin D levels and the risk of mortality in the general population. *Archives of internal medicine*. 2008; 168:1629–37. <https://doi.org/10.1001/archinte.168.15.1629> PMID:18695076 PMID:PMC2677029
- Almerighi C, Sinistro A, Cavazza A, Ciaprini C, Rocchi G, Bergamini A. 1Alpha,25-dihydroxyvitamin D3 inhibits CD40L-induced pro-inflammatory and immunomodulatory activity in human monocytes. *Cytokine*. 2009;45(3):190–7. <https://doi.org/10.1016/j.cyto.2008.12.009> PMID:19186073
- Lips P. Vitamin D physiology. *Progress in Biophysics and Molecular Biology*. 2006; 92:4–8. <https://doi.org/10.1016/j.pbiomolbio.2006.02.016> PMID:16563471
- Yetley EA, Brulé D, Cheney MC, Davis CD, Esslinger KA, Fischer PW et al. Dietary reference intakes for vitamin D: justification for a review of the 1997 values. *The American journal of clinical nutrition*. 2009; 89:719–27. <https://doi.org/10.3945/ajcn.2008.26903> PMID:19176741 PMID:PMC2667666
- Jones G. Pharmacokinetics of vitamin D toxicity. *The American journal of clinical nutrition*. 2008; 88:582S–586S. PMID:18689406
- Alqahatani M, Tamimi W, Aldaker M, Alenzi F, Tamim H, Alsadhan A. Young adult reference ranges for thyroid function tests on the Centaur immunoassay analyser. *British journal of biomedical science*. 2006;63:163–5. <https://doi.org/10.1080/09674845.2006.11732744> PMID:17201204
- Khan QJ, Fabian CJ. How I treat vitamin d deficiency. *Journal of oncology practice*. 2010; 6:97–101. <https://doi.org/10.1200/JOP.091087> PMID:20592785 PMID:PMC2835491
- Goswami R, Marwaha RK, Gupta N, Tandon N, Sreenivas V, Tomar N, et al. Prevalence of vitamin D deficiency and its relationship with thyroid autoimmunity in Asian Indians: a community-based survey. *British Journal of Nutrition*. 2009;102:382. <https://doi.org/10.1017/S0007114509220824> PMID:19203420
- Effraimidis G, Badenhop K, Tijssen JG, Wiersinga WM. Vitamin D deficiency is not associated with early stages of thyroid autoimmunity. *European journal of endocrinology / European Federation of Endocrine Societies*. 2012; 167:43–8. <https://doi.org/10.1530/EJE-12-0048> PMID:22518050
- Sezgin G, Ozer ME, Bayramicli OU, Ozel AM, Aksungar F, Nalbant S. Relationship of vitamin d deficiency and autoimmune thyroid diseases. *European Journal of Internal Medicine*. 2011; 22:S87. [https://doi.org/10.1016/S0953-6205\(11\)60355-5](https://doi.org/10.1016/S0953-6205(11)60355-5)
- Hussain AN, Alkhenizan AH, El Shaker M, Raef H, Gabr A. Increasing trends and significance of hypovitaminosis D: a population-based study in the Kingdom of Saudi Arabia. *Arch Osteoporos*. 2014;9(1):190. <https://doi.org/10.1007/s11657-014-0190-3> PMID:25213798
- Alkhenizan A, Mahmoud A, Hussain A, Gabr A, Alsoghayer S, Eldali A. The Relationship between 25 (OH) D Levels (Vitamin D) and Bone Mineral Density (BMD) in a Saudi Population in a Community-Based Setting. *Slominski AT, editor. PLoS One*. 2017;12(1):e0169122.
- Sadat-Ali M, AlElq A, Al-Turki H, Al-Mulhim F, Al-Ali A. Vitamin D levels in healthy men in eastern Saudi Arabia. *Annals of Saudi Medicine*. 2009;29:378-382. <https://doi.org/10.4103/0256-4947.55168> PMID:19700896 PMID:PMC3290044
- Fawzy E, Al-Sayed Mohamed S, El-Rabat AM. Hypovitaminosis D In Autoimmune Hypothyroidism. *Journal of*

American Science J Am Sci. 2013; 99:60–65.

24. Mackawy AMH, Al-ayed Bushra Mohammed, Al-rashidi Bashayer Mater. Vitamin D Deficiency and Its Association with Thyroid Disease. *International Journal of Health Sciences*. 2013;7(3):267-275. <https://doi.org/10.12816/0006054> PMID:24533019 PMCID:PMC3921055
25. Boyuk B, Altay M, Atalay H, Guzel S, Celebi A, Ekizoglu I. No relationship between 25 OH vitamin D and autoimmunity in women patients with hypothyroidism. *Acta Medica Mediterr*. 2016;32(3):779–84.
26. Kivity S, Agmon-Levin N, Zisapli M, Shapira Y, Szekanez Z. Thyroid disease and vitamin D deficiency. *Annals of the Rheumatic Diseases*. 2010;69:A62–A62. <https://doi.org/10.1136/ard.2010.129650q>
27. Baeke F, Takiishi T, Korf H, Gysemans C, Mathieu C. Vitamin D: modulator of the immune system. *Current opinion in pharmacology*. 2010;10:482–96. <https://doi.org/10.1016/j.coph.2010.04.001> PMID:20427238
28. Tamer G, Arik S, Tamer I, Coksert D. Relative vitamin D insufficiency in Hashimoto's thyroiditis. *Thyroid : official journal of the American Thyroid Association*. 2011; 21:891–6. <https://doi.org/10.1089/thy.2009.0200> PMID:21751884
29. Shin DY, Kim KJ, Kim D, Hwang S, Lee EJ. Low serum vitamin D is associated with anti-thyroid peroxidase antibody in autoimmune thyroiditis. *Yonsei medical journal*, 2014; 55:476–81. <https://doi.org/10.3349/yjm.2014.55.2.476> PMID:24532520 PMCID:PMC3936621
30. Choi YM, Kim WG, Kim TY, Bae SJ, Kim HK, Jang EK. et al. Low levels of serum vitamin D3 are associated with autoimmune thyroid disease in pre-menopausal women. *Thyroid : official journal of the American Thyroid Association*. 2014; 24:655–61. <https://doi.org/10.1089/thy.2013.0460> PMID:24320141 PMCID:PMC3993051
31. Wang J, Lv S, Chen G, Gao C, He J, Zhong H, Xu Y. et al. Meta-Analysis of the Association between Vitamin D and Autoimmune Thyroid Disease. *Nutrients*. 2015; 7:2485-2498. <https://doi.org/10.3390/nu7042485> PMID:25854833 PMCID:PMC4425156
32. Aljohani NJ, Al-Daghri NM, Al-Attas OS, Alokail MS, Alkharfy KM, Al-Othman A, Yakout S. et al. Differences and associations of metabolic and vitamin D status among patients with and without sub-clinical hypothyroid dysfunction. *BMC endocrine disorders*. 2013;13:31. <https://doi.org/10.1186/1472-6823-13-31> PMID:23962199 PMCID:PMC3751774
33. Al-Daghri NM, Al-Attas OS, Al-Okail MS, Alkharfy KM, Al-Yousef MA, Nadhrah HM. et al. Severe hypovitaminosis D is widespread and more common in non-diabetics than diabetics in Saudi adults. *Saudi medical journal*. 2010;31:775–80. PMID:20635011
34. Tamer G, Mesci B, Tamer I, Kilic D, Arik S. Is vitamin D deficiency an independent risk factor for obesity and abdominal obesity in women? *Endokrynologia Polska*. 2012;63:196–201. PMID:22744625
35. Saler T, Keşkek SO, Ahabab S, Cakir S, Ortoğlu G, Bankir M. Frequency of Hashimoto's thyroiditis in women with vitamin D deficiency: A cross sectional study. *American Journal of Internal Medicine*. 2014; 2:44–48. <https://doi.org/10.11648/j.ajim.20140203.12>
36. Zhang Q, Wang Z, Sun M, Mengdie Cao, Zhenxin Zhu, Qi Fu ,et al. Association of High Vitamin D Status with Low Circulating Thyroid-Stimulating Hormone Independent of Thyroid Hormone Levels in Middle-Aged and Elderly Males. *International Journal of Endocrinology*. 2014;2014:631819. <https://doi.org/10.1155/2014/631819> PMID:24693286 PMCID:PMC3947886
37. Kim, D. Low vitamin D status is associated with hypothyroid Hashimoto's thyroiditis. *Hormones (Athens, Greece)*. 2016;15(3):385-393. <https://doi.org/10.14310/horm.2002.1681>
38. Barchetta I, Baroni MG, Leonetti F, De Bernardinis M, Bertocchini L, Fontana M. et al. TSH levels are associated with vitamin D status and seasonality in an adult population of euthyroid adults. *Clinical and experimental medicine*. 2015;15:389–96. <https://doi.org/10.1007/s10238-014-0290-9> PMID:24925636
39. Vondra K, Stárka L, Hampl R. Vitamin D and Thyroid Diseases. *Physiol. Res*. 2015; 64:95–100.
40. Chaudhary S, Dutta D, Kumar M, Saha S, Mondal SA, Kumar A et al. Vitamin D supplementation reduces thyroid peroxidase antibody levels in patients with autoimmune thyroid disease: An open-labeled randomized controlled trial. *Indian journal of endocrinology and metabolism*. 2016; 20:391–8. <https://doi.org/10.4103/2230-8210.179997> PMID:27186560 PMCID:PMC4855971
41. Mansournia N, Mansournia MA, Saeedi S, Dehghan J. The association between serum 25OHD levels and hypothyroid Hashimoto's thyroiditis. *Journal of endocrinological investigation*. 2014; 37:473–6. <https://doi.org/10.1007/s40618-014-0064-y> PMID:24639121
42. Bársony J, Lakatos P, Földes J, Fehér T. Effect of vitamin D3 loading and thyroid hormone replacement therapy on the decreased serum 25-hydroxyvitamin D level in patients with hypothyroidism. *Acta endocrinologica*. 1986; 113:329–34. <https://doi.org/10.1530/acta.0.1130329>
43. Sworczak K, Wiśniewski P. The role of vitamins in the prevention and treatment of thyroid disorders. *Endokrynologia Polska*. 2011; 62:340–4. PMID:21879475
44. Clinckspoor I, Verlinden L, Mathieu C, Bouillon R, Verstuyf A, Decallonne B. Vitamin D in thyroid tumorigenesis and development. *Progress in Histochemistry and Cytochemistry*. 2013; 48:65–98. <https://doi.org/10.1016/j.proghi.2013.07.001> PMID:23890557
45. Penna-Martinez M, Ramos-Lopez E, Stern J, Kahles H, Hirsch N, Hansmann ML, Selinski I, et al. Impaired Vitamin D Activation and Association with CYP24A1 Haplotypes in Differentiated Thyroid Carcinoma. *Thyroid*. 2012; 22:709–716. <https://doi.org/10.1089/thy.2011.0330> PMID:22690899 PMCID:PMC3387756
46. Haney EM, Stadler D, Blizotes MM. Vitamin D insufficiency in internal medicine residents. *Calcified tissue international*. 2005;76: 11–6. <https://doi.org/10.1007/s00223-004-0025-0> PMID:15478001