

Breast Cancer Risk Among Male BRCA1 and BRCA2 Mutation Carriers

Yu Chuan Tai, Susan Domchek, Giovanni Parmigiani, Sining Chen

Men who carry germline mutations in the BRCA2 gene have a higher risk of developing breast carcinoma than men in the general population. Men who carry germline mutations in the BRCA1 gene may also be at a higher risk for breast carcinoma, but this association is not as well established. We evaluated the risks of developing breast carcinoma for male BRCA1 and BRCA2 mutation carriers in the US population based on data from 1939 families with 97 male subjects with breast carcinoma that were collected from eight centers across the National Cancer Institute's Cancer Genetics Network. At all ages, the cumulative risks of male breast cancer were higher in both BRCA1 and BRCA2 mutation carriers than in noncarriers. The relative risks of developing breast cancer were highest for men in their 30s and 40s and decreased with increasing age. Both the relative and cumulative risks were higher for BRCA2 mutation carriers than for BRCA1 mutation carriers. The estimated cumulative risk of breast carcinoma for male BRCA1 mutation carriers at age 70 years was 1.2% (95% confidence interval [CI] = 0.22% to 2.8%) and for BRCA2 mutation carriers, 6.8% (95% CI = 3.2% to 12%).

J Natl Cancer Inst 2007;99:1811-4

Male breast carcinoma accounts for less than 1% of all breast cancer cases in the United States (1,2). It is more common in men with a family history of breast cancer. Both genetic linkage and DNA sequencing data have demonstrated that male breast carcinoma is associated with deleterious germline mutations in BRCA2 (3,4). Several studies have examined the incidence of BRCA2 mutations in men with breast cancer (5-9). In one study that included 237 hereditary breast carcinoma families, 26 families had at least one male member who had been diagnosed with breast carcinoma and in 77%, the disease was linked to mutations in BRCA2 (5). Couch et al. (6) found that 14% of 50 male breast cancer patients who were not selected based on a family history of breast cancer had a deleterious BRCA2 mutation. Easton et al. (7) examined two large BRCA2-linked families that contained four male breast carcinoma cases. The estimated cumulative risk for male breast carcinoma by age 70 years was 6.3%. In a separate study of 164 families with BRCA2 mutations, the estimated cumulative risk of breast cancer for men who carried BRCA2 mutations was 2.8% by 70 years of age and 6.9% by 80 years of age (8). Recently, Risch et al. (9) estimated that

the relative risk of breast cancer for male BRCA2 carriers was 102 (95% confidence interval [CI] = 9.9 to 1050).

The association between male breast carcinoma and deleterious germline mutations in BRCA1 is less clear. Initial data suggested that inherited male breast cancer was not linked to germline mutations in BRCA1 (10) and BRCA1 mutations have been detected at very low frequency in several series of unselected male breast cancer patients (11-14). However, a more recent study (15) in a series of 10 000 individuals reported that 21 of 76 men with breast cancer had mutations in either BRCA1 or BRCA2; more than one-third of those mutations were in BRCA1. Brose et al. (16) estimated a lifetime risk of male breast cancer of 5.8% in BRCA1 mutation carriers, but this estimate was based on families who sought breast cancer risk counseling and was not adjusted for ascertainment. The purpose of our study was to evaluate the risk of breast carcinoma for male BRCA1 and BRCA2 mutation carriers by using a large sample of US family histories.

We used a database that contained detailed family history information for 1948 families with complete pedigrees that were collected from eight centers across the National Cancer

Institute's Cancer Genetics Network (CGN). The CGN has been described in detail (17). Nine families were excluded from the analysis because of invalid entries in the pedigree. Among the remaining 1939 families, 676 were of Ashkenazi Jewish ancestry. Seven of the eight centers were high-risk counseling clinics, and one was a population-based clinic. Therefore, most of the families were ascertained based on a family history of breast and/or ovarian cancer. The institutional review boards at each participating institution approved the study protocol.

Of the 1939 families included in this analysis, 87 families (4.5%) had one or more male members with breast carcinoma; the total number of male subjects with breast carcinoma was 97. Only one person in each family, i.e., the proband, was genotyped. In 45 families (52%), the male breast cancer patient was the proband. Of the 87 probands, 23 (26%) had detectable BRCA2 mutations and six (7%) had detectable BRCA1 mutations. The median age at diagnosis of male breast cancer was 64 years (range = 24-87 years). There were three male subjects with contralateral breast cancer. Table 1 presents information for the 97 male breast cancer subjects.

We estimated the risk of breast cancer for men who carried a BRCA1 or BRCA2 mutation by assuming that the age-conditional probability of developing cancer at age t was that of a noncarrier multiplied by the relative risk at that age $[RR(t)]$, which was defined as $\exp(\alpha + \beta \times t)$, where α quantifies the relative risk at age 0 and β quantifies the change in relative risk with respect to age. We assumed that the

Affiliations of authors: Institute for Human Genetics, University of California, San Francisco, CA (YCT); Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA (SD); Departments of Oncology (GP, SC) and Pathology (GP), Johns Hopkins School of Medicine, Baltimore, MD; Departments of Biostatistics (GP, SC) and Environmental Health Sciences (SC), Johns Hopkins Bloomberg School of Public Health, Baltimore, MD.

Correspondence to: Sining Chen, PhD, 615 North Wolfe Street, Rm W7041, Baltimore, MD 21205 (e-mail: sichen@jhsph.edu).

See "Funding" and "Notes" following "References."

DOI: 10.1093/jnci/djm203

© The Author 2007. Published by Oxford University Press. All rights reserved. For Permissions, please e-mail: journals.permissions@oxfordjournals.org.

CONTEXT AND CAVEATS

Prior knowledge

Men who carry germline mutations in the BRCA2 gene have a higher risk of developing breast carcinoma than men in the general population. However, the association between germline mutations in the BRCA1 gene and the risk of breast carcinoma in men is not as well established.

Study design

A retrospective study of the risk of developing breast carcinoma for male BRCA1 and BRCA2 mutation carriers in the United States based on data from 1939 families with 97 men with breast carcinoma that were collected from eight centers across the National Cancer Institute's Cancer Genetics Network.

Contribution

At all ages, the cumulative risks of male breast cancer were higher in both BRCA1 and BRCA2 mutation carriers than in non-carriers. The relative risks of developing breast cancer were highest for men in their 30s and 40s and decreased with increasing age. Both the relative and cumulative risks were higher for BRCA2 mutation carriers than for BRCA1 mutation carriers.

Implications

These risk estimates are important for determining appropriate risk management strategies for the male members of families with germline mutations in BRCA1 or BRCA2.

Limitations

No information about family members' ethnicity group other than their Ashkenazi Jewish ancestry was available, precluding estimates of the penetrance of mutations in BRCA1 and BRCA2 for African Americans.

age-conditional probabilities of developing cancer for noncarriers were the same as those for the US male population (18). We also assumed that the relative risks were the same for Ashkenazi Jewish and non-Ashkenazi Jewish individuals but that the relative risks differed between BRCA1 mutation carriers and BRCA2 mutation carriers.

Because our study included subjects who pursued mutation testing for BRCA1 and BRCA2, it included a greater representation of families with a history of breast and ovarian cancer than studies that have

Table 1. Male breast cancers, grouped by family

Mutated gene	Mutation	Ethnicity	Age at first diagnosis*, y	No. of female breast cancers in family†
BRCA1	633delC	Other	46‡	2
BRCA1	2931CC→G	Other	64‡	1
BRCA1	2985del5	Other	73	4
BRCA1	5149del5	Other	51	3
BRCA1	5296del4	Other	68	4
BRCA1	Exon 13 dup	Other	42	4
BRCA2	379delG	Other	79‡, 49	1
BRCA2	1128insG	Other	48‡	2
BRCA2	3034del4	Other	52‡, §	3
BRCA2	Y1894X	Other	64‡, 70	1
BRCA2	6174delT	Other	46‡	0
BRCA2	6174delT	Other	61‡	0
BRCA2	6174delT	Other	75‡	2
BRCA2	7989delC	Other	53‡, 70	5
BRCA2	1538del4	Other	66	6
BRCA2	1982delA	Ashkenazi Jewish	85, 71	4
BRCA2	2041insA	Other	51	4
BRCA2	3034del4	Other	47	4
BRCA2	3945delA	Other	66, 42	4
BRCA2	4075delGT	Ashkenazi Jewish	70	2
BRCA2	5482delC	Other	70	1
BRCA2	5849del4	Other	62, 65	7
BRCA2	5950delCT	Other	65	1
BRCA2	6051delA	Other	58	2
BRCA2	6174delT	Other	49	3
BRCA2	6174delT	Ashkenazi Jewish	61	1
BRCA2	6659delA	Other	49	4
BRCA2	9538delAA	Other	50	5
BRCA2	Q3066X	Other	26	5
None found	–	18 Ashkenazi Jewish	60 (13.7)	1.56 (1.29)
None found	–	44 Other	64 (12.2)	1.45 (1.38)

* When more than one breast cancer was diagnosed in a single man, the age at each diagnosis is presented for BRCA1 and BRCA2 mutation carriers, and the mean age at diagnosis (standard deviation) is presented for noncarriers.

† The mean number of female breast cancers (standard deviation) is presented for noncarriers.

‡ Carriage of a mutation was confirmed by genotype analysis.

§ This man had contralateral breast cancer. Cancer in the second breast was diagnosed at age 55 years.

|| This group includes two men with contralateral male breast cancer: one was diagnosed at ages 58 and 59 years, and the other was diagnosed at age 45 years for both breasts. Both patients were genotyped, but no mutation was found.

examined unselected cases of male breast cancer. To correct for possible selection bias, we used a retrospective likelihood approach (19), in which each family's contribution to the likelihood of penetrance parameters is the probability of the genetic test results conditioning on the observed family history. This approach produces unbiased estimates of risk provided that the probability of including a family in the study depends only on the family history (19) and not on other information relevant to the presence of a genetic mutation (e.g., linkage score). Because all subjects included in this study were the first person in their family to be tested for a BRCA1 or

BRCA2 mutation, this approach gives unbiased estimates in our study, even though the selection of subjects into the study involves, to some extent, the subjects' own preferences.

Table 2 presents the estimated age-specific cumulative risks and relative risks of developing breast cancer for male BRCA1 and BRCA2 mutation carriers. Prospective cancer risks are provided in the BRCAPRO program (20,21). At all ages, the cumulative risks of male breast cancer were higher in both BRCA1 and BRCA2 mutation carriers than in noncarriers. The estimated cumulative risk of breast carcinoma for male BRCA1 mutation carriers at age 70 years

Table 2. Age-specific cumulative risks and relative risks of developing breast cancer for male BRCA1 and BRCA2 mutation carriers and noncarriers*

Age, y	Cumulative breast cancer risk, % (95% CI)			RR (95% CI)	
	BRCA1 mutation carrier	BRCA2 mutation carrier	No BRCA1 or BRCA2 mutation†	BRCA1 mutation carrier	BRCA2 mutation carrier
30	1.7×10^{-2} (6.8×10^{-4} to 0.12)	0.18 (0.027 to 0.85)	1.2×10^{-4}	89 (6 to 505)	936 (201 to 3277)
40	0.12 (0.012 to 0.58)	1.2 (0.3 to 3.6)	1.9×10^{-3}	41 (6 to 170)	396 (129 to 986)
50	0.3 (0.052 to 1.2)	2.7 (0.94 to 6.6)	8.5×10^{-3}	23 (5 to 61)	178 (76 to 344)
60	0.62 (0.13 to 1.7)	4.7 (1.9 to 9.5)	2.7×10^{-2}	16 (3 to 40)	84 (38 to 158)
70	1.2 (0.22 to 2.8)	6.8 (3.2 to 12)	6.7×10^{-2}	12 (1 to 34)	42 (15 to 91)
80	1.8 (0.3 to 4.5)	8.3 (3.9 to 15)	0.12	11 (0.3 to 34)	22 (5 to 57)

* CI = confidence interval; RR = relative risk.

† The cumulative risk estimates for noncarriers were obtained from the Surveillance, Epidemiology, and End Results database, which does not provide 95% CIs.

was 1.2% (95% CI = 0.22% to 2.8%), and for BRCA2 mutation carriers, 6.8% (95% CI = 3.2% to 12%). The relative risks of developing male breast cancer were highest for men in their 30s and 40s and decreased with increasing age. This trend was particularly pronounced in BRCA2 mutation carriers: the relative risk at age 30 years was 22.3 times that at age 70 years.

Our study has several strengths. First, the US cohort that we used to examine male breast cancer risk is the largest to date, contains families of Ashkenazi Jewish and other ancestries, and provided risk estimates for BRCA1 and BRCA2 mutation carriers in the US population. Such risk estimates are important for determining appropriate risk management strategies for the male members of families with germline mutations in BRCA1 or BRCA2. Our cumulative risk estimates for BRCA2 mutation carriers are similar to those reported by Easton et al. (7) and Risch et al. (9). Second, to our knowledge, this study is the first to characterize the risks for male BRCA1 mutation carriers in an unbiased fashion with a large cohort. Our demonstration that BRCA1 mutations are associated with an elevated risk of male breast cancer, but that those risks are still substantially lower than those in BRCA2 mutations carriers, confirms previous suggestions (2,10) that the link between BRCA1 mutations and male breast carcinoma is weaker than that between BRCA2 mutations and male breast cancer.

A potential limitation of our approach is that the values we used for the BRCA1 and BRCA2 mutated allele frequencies were fixed at values derived from the literature instead of estimated from the data.

For individuals of Ashkenazi Jewish ancestry, we used mutated allele frequencies of 0.0060 and 0.0067 for BRCA1 and BRCA2, respectively, according to a meta-analysis of published estimates (22). For individuals who were not of Ashkenazi Jewish ancestry, we used mutated allele frequencies of 0.00058 and 0.00067 for BRCA1 and BRCA2, respectively (23,24). Risch et al. (9) recently published mutated allele frequencies of 0.0016 and 0.0035 for BRCA1 and BRCA2, respectively, for individuals not of Ashkenazi Jewish ancestry living in Ontario, Canada. This BRCA2 mutated allele frequency is fivefold larger than the frequency we used. To test the robustness of our risk estimates with respect to allele frequency, we repeated the analysis with the higher allele frequencies. As a result, the cumulative risks by age 80 years decreased from 1.8% (95% CI = 0.3% to 4.5%) to 1.5% (95% CI = 0.3% to 3.9%) for BRCA1 and from 8.3% (95% CI = 3.9% to 15%) to 5.1% (95% CI = 2.0% to 11.0%) for BRCA2. The relative risks at younger ages were also smaller than the ones we estimated in the original analysis (e.g., the relative risks for BRCA1 and BRCA2 mutation carriers at age 30 years were 24 [95% CI = 2.4 to 102] and 430 [95% CI = 68 to 1759], respectively). The relative risks gradually decreased with increasing age, and the relative risks at age 80 years (e.g., 11 [95% CI = 1.7 to 34] for BRCA1 mutation carriers and 16 [95% CI = 5.1 to 42] for BRCA2 mutation carriers) were similar to the ones estimated in our original analysis (Table 2). These similar risk estimates suggest that our approach was robust across a range of allele frequencies.

Another limitation of our study is that we had no information about family

members' ethnicity group other than their Ashkenazi Jewish ancestry. Therefore, we were not able to estimate the penetrance of mutations in BRCA1 and BRCA2 for African Americans, for whom the male-to-female ratio for breast carcinoma is higher than it is for white populations (2,25).

The male penetrance estimates reported here have been incorporated into the current version of the BayesMendel genetic counseling software and which is available to genetic counselors through the CaGene software package (26).

References

- Landis SH, Murray T, Bolden S, Wingo PA. Cancer statistics, 1999. *CA Cancer J Clin* 1999; 49:8-31.
- Lynch HT, Watson P, Narod SA. The genetic epidemiology of male breast carcinoma. *Cancer* 1999;86:744-6.
- Thorlacius S, Tryggvadottir L, Olafsdottir GH, Jonasson JG, Ogmundsdottir HM, Tulinius H, et al. Linkage to BRCA2 region in hereditary male breast cancer. *Lancet* 1995; 346:544-5.
- Wooster R, Bignell G, Lancaster J, Swift S, Seal S, Mangion J, et al. Identification of the breast cancer susceptibility gene BRCA2. *Nature* 1995;378:789-92.
- Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, et al. Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. *Am J Hum Genet* 1998;62:676-89.
- Couch FJ, Farid LM, DeShano ML, Tavtigian SV, Calzone K, Campeau L, et al. BRCA2 germline mutations in male breast cancer cases and breast cancer families. *Nat Genet* 1996;13:123-5.
- Easton DF, Steele L, Fields P, Ormiston W, Averill D, Daly PA, et al. Cancer risks in two large breast cancer families linked to BRCA2 on chromosome 13q12-13. *Am J Hum Genet* 1997;61:120-8.

- (8) Thompson D, Easton D. Variation in cancer risks, by mutation position, in BRCA2 mutation carriers. *Am J Hum Genet* 2001;68:410–19.
- (9) Risch HA, McLaughlin JR, Cole DE, Rosen B, Bradley L, Fan I, et al. Population BRCA1 and BRCA2 mutation frequencies and cancer penetrances: a kin-cohort study in Ontario, Canada. *J Natl Cancer Inst* 2006;98:1694–706.
- (10) Stratton MR, Ford D, Neuhasen S, Seal S, Wooster R, Friedman LS, et al. Familial male breast cancer is not linked to the BRCA1 locus on chromosome 17q. *Nat Genet* 1994;7:103–7.
- (11) Struewing JP, Coriaty ZM, Ron E, Livoff A, Konichezky M, Cohen P, et al. BRCA1/2 mutations among male patients with breast cancer in Israel. *Am J Hum Genet* 1999;65:1800–2.
- (12) Sverdlov RS, Barshack I, Bar Sade RB, Baruch RG, Hirsh-Yehezkel G, Dagan E, et al. Genetic analyses of male breast cancer in Israel. *Genet Test* 2000;4:313–7.
- (13) Friedman LS, Gayther SA, Kurosaki T, Gordon D, Noble B, Casey G. Mutation analysis of BRCA1 and BRCA2 in a male breast cancer population. *Am J Hum Genet* 1997;60:313–9.
- (14) Csokay B, Udvarhelyi N, Sulyok Z, Besznyak I, Ramus S, Ponder B, et al. High frequency of germline BRCA2 mutations among Hungarian male breast cancer patients without family history. *Cancer Res* 1999;59:995–8.
- (15) Frank TS, Deffenbaugh AM, Reid JE, Hulick M, Ward BE, Lingenfelter B, et al. Clinical characteristics of individuals with germline mutations in BRCA1 and BRCA2: analysis of 10,000 individuals. *J Clin Oncol* 2002;20:1480–90.
- (16) Brose MS, Rebbeck TR, Calzone KA, Stopfer JE, Nathanson KL, Weber BL. Cancer risk estimates for BRCA1 mutation carriers identified in a risk evaluation program. *J Natl Cancer Inst* 2002;94:1365–72.
- (17) Chen S, Iversen ES, Friebel T, Finkelstein D, Weber BL, Eisen A, et al. Characterization of BRCA1 and BRCA2 mutations in a large United States sample. *J Clin Oncol* 2006;24:863–71.
- (18) Fay MP, Pfeiffer R, Cronin KA, Le C, Feuer EJ. Age-conditional probabilities of developing cancer. *Stat Med* 2003;22:1837–48.
- (19) Kraft P, Thomas DC. Bias and efficiency in family-based gene-characterization studies: conditional, prospective, retrospective, and joint likelihoods. *Am J Hum Genet* 2000;66:1119–31.
- (20) Parmigiani G, Berry DA, Aguilar O. Determining carrier probabilities for breast cancer susceptibility genes BRCA1 and BRCA2. *Am J Hum Genet* 1998;62:145–58.
- (21) Chen S, Wang W, Broman K, Katki HA, Parmigiani G. BayesMendel: an R environment for Mendelian risk prediction. *Stat Appl Genet Mol Biol* 2004;3(1):Article 21.
- (22) Chen S, Parmigiani G. Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol* 2007;25:1329–33.
- (23) Andersen TI. Genetic heterogeneity in breast cancer susceptibility. *Acta Oncol* 1996;35:407–10.
- (24) Antoniou AC, Pharoah PDP, McMullan G, Day NE, Stratton MR, Peto J, et al. A comprehensive model for familial breast cancer incorporating BRCA1, BRCA2 and other genes. *Br J Cancer* 2002;86:76–83.
- (25) Sasco AJ, Lowenfels AB, Pasker-De Jong P. Review article: epidemiology of male breast cancer. A meta-analysis of published case-control studies and discussion of selected aetiological factors. *Int J Cancer* 1993;53:538–49.
- (26) Euhus DM, Smith KC, Robinson L, Stucky A, Olopade OI, Cummings S, et al. Pretest prediction of BRCA1 or BRCA2 mutation by risk counselors and the computer model BRCAPRO. *J Natl Cancer Inst* 2002;94:844–51.

Funding

Cancer Genetics Network at Duke University (U24 CA78157) and at The Johns Hopkins University (U24 CA78148); The Johns Hopkins Specialized Projects of Oncology Research Excellence in Breast Cancer (P50CA88843 to G. P.); National Cancer Institute (R01CA105090-01A1 to S. C., G. P., and Y. C. T. and P50CA62924-05 to G. P.).

Notes

The authors wish to thank the following members of the CGN BRCA1/2 Models Validation Study, who contributed to the original collection of the cohort used: Edwin S. Iversen; Tara M. Friebel; Dianne M. Finkelstein; Barbara L. Weber; Andrea Eisen; Leif E. Peterson; Joellen M. Schildkraut; Claudine Isaacs; Camille Corio; Leoni Leondaridis; Gail Tomlinson; Christopher I. Amos; Louise C. Strong; Donald A. Berry; Jeffrey N. Weitzel; Sharon Sand; Debra Dutton; Rich Kerber; Beth N. Peshkin; and David M. Euhus. The CGN steering committee reviewed the design of the study. The CGN funded part of the study and is responsible for good research practices and for data storage. The CGN and the other funding sources, listed in the Funding section, had no other role in the design, conduct, or reporting of the study.

Manuscript received February 21, 2007; revised September 24, 2007; accepted September 25, 2007.