

Development and implementation of multilocus sequence typing to study the diversity of the yeast *Kluyveromyces marxianus* in Italian cheeses

Fabrizia Tittarelli,^{1,2} Javier A. Varela,² Loughlin Gethins,² Catherine Stanton,^{3,4} R.P. Ross,⁴ Giovanna Suzzi,¹ Luigi Grazia,⁵ Rosanna Tofalo¹ and John P. Morrissey^{2,4,*}

Abstract

The yeast *Kluyveromyces marxianus* possesses advantageous traits like rapid growth, GRAS (generally regarded as safe) status and thermotolerance that make it very suitable for diverse biotechnological applications. Although physiological studies demonstrate wide phenotypic variation within the species, there is only limited information available on the genetic diversity of *K. marxianus*. The aim of this work was to develop a multilocus sequence typing (MLST) method for *K. marxianus* to improve strain classification and selection. Analysis of housekeeping genes in a number of sequenced strains led to the selection of five genes, *IPP1*, *TFC1*, *GPH1*, *GSY2* and *SGA1*, with sufficient polymorphic sites to allow MLST analysis. These loci were sequenced in an additional 76 strains and used to develop the MLST. This revealed wide diversity in the species and separation of the culture collection and wild strains into multiple distinct clades. Two subsets of strains that shared sources of origin were subjected to MLST and split decomposition analysis. The latter revealed evidence of recombination, indicating that this yeast undergoes mating in the wild. A public access web-based portal was established to allow expansion of the database and application of MLST to additional *K. marxianus* strains. This will aid understanding of the genetic diversity of the yeast and facilitate biotechnological exploitation.

DATA SUMMARY

Accession numbers are provided in Table 1. The software used was as follows:

Oligo Analyzer (<https://eu.idtdna.com/calc/analyzer>); UGENE (<http://ugene.net/>); Mega 7 (<http://megasoftware.net/>); SnapGene (<https://snapgene.com/>); SplitsTree (<http://splitstree.org/>).

INTRODUCTION

Kluyveromyces marxianus is a homothallic yeast that, along with its sister species *Kluyveromyces lactis*, belongs to the subphylum Saccharomycotina. Some strains are able to utilize lactose as a carbon source but, unlike *K. lactis*, it can also secrete enzymes that degrade inulin, a fructan found in plants such as Jerusalem artichokes and agave [1, 2]. The capacity to assimilate lactose means that *K. marxianus* is

frequently isolated from dairy products, such as fermented milk, yoghurt, kefir, koumiss and cheese, although it can also be isolated from fermenting fruit and plant material [3–7]. The broad metabolic capacity combined with a rapid growth rate, thermotolerance (up to 52 °C) and a high secretory capacity (e.g. secretion of lytic enzymes) have led to multiple applications in biotechnology; for example, production of bioingredients and aroma compounds, production of ethanol by fermentation, removal of lactose from food, and bioremediation [8, 9]. Because of its long and safe association with foods, *K. marxianus* is a GRAS (generally regarded as safe) organism in the USA and is on the European Food Safety Authority QPS (qualified presumption of safety) list.

There is not a unique *K. marxianus* strain that has been extensively characterized as a ‘model’ and those strains that

Received 20 October 2017; Accepted 2 January 2018

Author affiliations: ¹Faculty of BioScience and Technology for Food, Agriculture and Environment, University of Teramo, Teramo, Italy; ²School of Microbiology, Environmental Research Institute, Centre for Synthetic Biology and Biotechnology, University College Cork, Cork T12YN60, Ireland; ³Teagasc Research Centre, Moorepark, Ireland; ⁴School of Microbiology, APC Microbiome Institute, University College Cork, Cork, Ireland; ⁵Department of Science and Technology for Food and Agriculture (DISTAL), Alma Mater Studiorum, University of Bologna, Bologna, Italy.

*Correspondence: John P. Morrissey, j.morrissey@ucc.ie

Keywords: MLST; dairy products; *Kluyveromyces marxianus*; population.

Abbreviations: DST, diploid sequence type; IUPAC, International Union of Pure and Applied Chemistry; MLST, multilocus sequence typing; SNP, single nucleotide polymorphism; UPGMA, unweighted pair-group method with average.

Data statement: All supporting data, code and protocols have been provided within the article or through supplementary data files. Two supplementary tables and four supplementary figures are available with the online version of this article.

have been studied display substantial physiological diversity [2, 10]. Although a challenge for detailed molecular and metabolic studies, this diversity is likely to be a very important enabler of a wide spectrum of biotechnological applications and, thus, can be considered positively. Nonetheless, full exploitation of this potential requires an improved understanding of the genetic diversity, and a means to identify and categorize strains. Sequence variation in the D1/D2 region of the 26S rRNA gene is universally accepted as the best way to distinguish yeast species from each other [11], but since this sequence is invariant within a species, this is not useful for intraspecific studies. Several researchers in recent years have applied molecular approaches to address this question. Differences at the intraspecific level in *K. marxianus* have been studied by restriction analysis of the non-transcribed spacer (NTS) region of the rRNA gene cluster, RAPD (random amplified polymorphic DNA) PCR fingerprinting and variability of the insertion of the long terminal repeat (LTR) retrotransposon Tkm1 [12–16]. Karyotyping of *K. marxianus* strains from different isolation sources and geographical origins suggest diverse chromosome numbers, although analysis of genomes of sequenced strains is consistent with a standard complement of eight chromosomes [17, 18]. This fits the emerging view that stable haploid and diploid *K. marxianus* strains occur in natural environments [19]. It is notable, however, that a wide haplotype diversity was observed through RFLPs of mitochondrial DNA [15, 20]. A rapid, easy to use, scalable method in yeast and fungi is microsatellite analysis. Microsatellites are short tandem DNA repeats that can be found in all genomes and are used as molecular markers able to

IMPACT STATEMENT

Kluyveromyces marxianus is emerging as one of the most important non-*Saccharomyces* yeasts under development for biotechnological applications. It has traits such as rapid growth, high yield and thermotolerance that make it relevant for exploitation. In addition, genome engineering tools are now available to allow construction of new strains. There is, however, little information on the genetic diversity of the species, or on the relationship between strains. This is resolved in this study through the development of an multilocus sequence typing system. To facilitate use by the research community, a web-based database and tool was implemented to allow any researchers to easily type and classify more *K. marxianus* strains in the future.

distinguish strains. They have been applied for typing a variety of yeasts, such as *Saccharomyces uvarum* and *Saccharomyces cerevisiae*. These studies showed that the patterns are stable and that the method is discriminatory; thus, it can be used for population structure studies [21, 22].

Comparison of the relationships among *K. marxianus* strains is difficult and does not yield consistent patterns [23]. Furthermore, none of those methods is based on phylogeny, and so including new isolates and strains is problematic. These drawbacks can be overcome with the use of methods that are based on genomic sequence variation,

Table 1. *K. marxianus* strains used in this study and their origin

Strain	Isolation source	Culture collection/accession no.
LM3, LM6, LM9, LM14, LM15, LM17, LM20, LM28, LM30, LM32, LM36, LM38, LM42, LM44, LM47, LM48, LM50, LM54, LM72, LM92, LM96, LM114, LM116, LM127, LM133, LM136, LM139, LM141, LM142, LM148, LM153, LM154, LM161, LM167, LM169, LM174	Parmigiano Reggiano natural whey starter culture	Department of Agricultural and Food Sciences, University of Bologna, Italy [33]
6M2, 1SC4, K326	Parmigiano Reggiano cheese	
M12, M14, M38, M41, M48, M68, M81, M83, M123, M131, M135, M166, M169	Pecorino di Farindola cheese	Faculty of BioScience and Technology for Food, Agriculture, and Environment, University of Teramo, Italy [13]
VG1, VG4, VG6	Cow's milk whey	
FM09	Fermented milk	
CBS 4857, CBS 1555, CBS 5795, CBS 397, CBS 2762, CBS 6432, CBS 6556, CBS 712, CBS 7894, CBS 7858, CBS 1596, CBS 2233, CBS 4354, CBS 608, CBS 5668, CBS 745, CBS 834 ^T		CBS-KNAW, Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands
NCYC 179, NCYC 1424		National Collection of Yeast Cultures, UK
UCKM1	Unpasteurized cow's milk	School of Microbiology, University College Cork, Ireland [6]
DMKU 3-1042	Genome sequence	AP012220.1
NBRC 1777	Genome sequence	AP014606.1
CCT 7735	Genome sequence	CP009310.1
DMB1	Genome sequence	BBIL00000000.1
KCTC 17555	Genome sequence	AKFM00000000.2
IPE453	Genome sequence	LDJA00000000.1
BO339	Genome sequence	LXZY00000000.1

especially multilocus sequence typing (MLST), which is based on analysing polymorphic sites in a number of conserved housekeeping genes within a species, permits easy comparison and sharing of the results among laboratories, and minimizes the possibility of errors resulting from local experimental conditions [24]. MLST has frequently been used to evaluate the genetic structure of different bacteria and yeasts [25, 26]. For example, this technique has been applied to study population structure and evolution of *S. cerevisiae* [27–29] and of the pathogenic yeast *Candida albicans* [30–32]. The aim of this work was to develop MLST for *K. marxianus* to enable strain identification and classification. As proof of principle, this method was applied to study the diversity of a collection of *K. marxianus* strains that had been isolated from two Italian cheese fermentations.

METHODS

Origin of yeast strains

The 83 *K. marxianus* strains included in this study are listed in Table 1, which also lists the strains for which genome sequences were available for this study. Most of the isolates were from dairy products from different geographical areas in Italy, and were obtained from previous studies [13, 33]. In addition, 19 *K. marxianus* strains were from the Westerdijk Fungal Biodiversity Institute (Utrecht, The Netherlands) and from the National Collection of Yeast Cultures (Norwich, UK), and 1 was from the University College Cork collection. Yeasts were maintained as culture stocks in YPD (1%, w/v, yeast extract, 2%, w/v, peptone, 2%, w/v, dextrose) medium containing 20% (v/v) glycerol at -80°C . All the strains were routinely grown in YPD medium overnight at 28°C in shake flasks.

Choice of loci and design of primers

Five housekeeping genes were retrieved from the *K. marxianus* genomes of seven sequenced strains (CCT 7735, DMB1, DMKU 3-1042, NBRC 1777, KCTC 17555, IPE453, BO339), and a multiple sequence alignment was carried out to identify single nucleotide polymorphisms (SNPs). These genes, *IPP1*, *TFCl*, *GPH1*, *GSY2* and *SGA1*, showed good SNP distribution and were used in this study. Primers were designed with SnapGene software to hybridize to non-variable regions of each gene, thereby allowing amplification of the polymorphic region in any *K. marxianus* strain. The primer sequences and amplicon sizes are shown in Table 2. A BLAST (basic alignment search tools) search was used to check the specificity for *K. marxianus* of each primer set. Furthermore, the properties of each primer were verified by Oligo Analyzer 3.1. *K. lactis* CBS 683^T was used as a negative control to evaluate primer specificity by PCR.

DNA extraction, amplification and nucleotide sequencing

DNA was extracted with the QIAprep Spin miniprep kit, following the manufacturer's guidelines (Qiagen), modified as follows: 5 ml overnight culture was resuspended in 250 μl

resuspension buffer [50 mM Tris-HCl (pH 8.0), 10 mM EDTA and 10 μg RNase A ml^{-1}] and vortexed for 5 min with glass beads (0.15–0.6 mm in diameter; Sigma). DNA quantification was carried out with a NanoDrop ND-1000 spectrophotometer. PCR analysis was performed with a ProFlex PCR system thermal cycler (Applied Biosystems). The reaction mixture (25 μl) contained 100 ng genomic DNA, 13 μl GoTaq Green master mix (Promega) and 10 μM each primer. The PCR conditions were: initial denaturation at 95°C for 5 min; followed by 30 cycles of denaturation at 95°C for 30 s, annealing at 56°C for 15 s and elongation at 72°C for 1 min; followed by a final extension step at 72°C for 5 min. PCR products were separated in 1% (w/v) agarose gel in $1\times$ TAE buffer [40 mM Tris (pH 7.6), 20 mM acetic acid, 1 mM EDTA (pH 8.0)] at 80 V for 1 h, stained with SafeView (NBS Biologicals) and visualized under UV light. DNA fragment sizes were determined using a 1 kb DNA HyperLadder (BioLine). The amplified products were purified with a commercial PCR purification system (QIAquick PCR purification kit; Qiagen), and amplicons were sequenced in each direction by GATC (Constance, Germany).

Data analysis

The sequence analysis for each gene fragment was performed with Mega 7 [34] and Unipro UGENE programs [35]. For each fragment, the sequences obtained from the 83 strains were manually edited and compared with those from the 7 *K. marxianus* reference sequences. Defined regions in all sequences obtained from a single locus were aligned using Muscle [36]. All the polymorphic sites were visually inspected, and the polymorphic nucleotide sites within these sequences were identified. In some cases, there were two coincident, equivalently sized peaks in the forward and reverse sequences, indicating heterozygosity at that position. To avoid erroneous interpretation of double peaks, we followed previous suggestions, and routinely took account of noise in neighbouring positions and compared the chromatograms for both sequenced strands [37]. The one-letter code for nucleotides from the International Union of Pure and Applied Chemistry (IUPAC) nomenclature was used to designate homozygous and heterozygous polymorphic sites. Genotype numbers were assigned to each unique sequence. Each strain was defined by the combination of numbers corresponding to the genotype at the loci analysed, which is a genotype profile or diploid sequence type (DST). Sequences different even at a single nucleotide site were considered distinct genotypes.

Similarities among allelic profiles were displayed with a dendrogram generated by using the unweighted pair-group method with average (UPGMA) algorithm. This data analysis was performed by using START software [38]. For the computation of the phylogenetic tree, based on the concatenated sequence of the five loci, a distance matrix was created and a phylogenetic tree was reconstructed with the program Unipro UGENE by using the maximum-likelihood

Table 2. Characteristics of the five housekeeping genes and primer sequences

Gene	Molecular function	Gene length (bp)	Primer (5'→3')	Amplicon size (bp)	5' (3') position start	Size of sequenced fragment (bp)	No. of genotypes identified	No. of variable sites		
								Polymorphic nucleotide site	Non-synonymous polymorphic amino acid site	Synonymous polymorphic amino acid site
<i>IPPI1</i>	Inorganic diphosphatase activity	864	Fwd: ATCGGTGCCAAGAA-CACCTT Rev: ITGTCGATTGGCTCG	803	44 (807)	763	28	25	4	21
<i>TFC1</i>	RNA Pol III transcription factor activity	1857	Fwd: AAGCCGATTGGG TCTGG TCAAAAC Rev: TCTGGGACTCAGAG TTATGC	832	992 (1783)	791	37	44	19	25
<i>GPH1</i>	Glycogen phosphorylase activity	2712	Fwd: TGAACACTGTGAAG-CAGCA Rev: ITTCGTCAGCGTAC TCCTGG	902	1797 (2659)	862	30	34	8	26
<i>GSY2</i>	Glycogen synthase activity	2106	Fwd: CACCCATGAGA TCCCTCA Rev: CGTCCTCTCGTCCG TCATCC	891	1197 (2068)	871	34	49	19	30
<i>SGA1</i>	Glucan 1,4- α -glucosidase activity	1788	Fwd: CTCCGATGGTTCCGGG TCAAT Rev: TCATGGGTCAAGGTAC TGGC	926	179 (1085)	906	26	44	13	31

Table 3. Genotypes of the 83 strains of *K. marxianus* analysed by MLST

Strain name	Genotype					DST
	<i>IPP1</i>	<i>TFC1</i>	<i>GPH1</i>	<i>GSY2</i>	<i>SGA1</i>	
LM127*, LM30*, LM38*	1	1	1	1	1	1
LM3*, LM50*, LM96*	1	1	1	1	14	2
LM47*	1	1	1	1	18	3
LM32*	1	1	1	2	1	4
LM6*	1	1	1	4	2	5
LM42*	1	1	1	17	16	6
LM36*	1	1	6	1	1	7
CBS 1555, CBS 712, CBS 7858, CBS 608	1	1	8	1	1	8
6M2*	1	1	10	3	4	9
CBS 834 ^T	1	1	11	1	4	10
VG6†	1	1	16	17	2	11
K326*	1	1	26	16	4	12
M169‡, M38‡	1	1	26	17	4	13
LM139*	1	4	1	1	1	14
LM133*	1	4	1	1	15	15
CBS 7894	1	3	9	22	1	16
1SC4*	1	6	13	12	15	17
LM116*	1	7	1	7	15	18
LM72*	1	12	4	1	1	19
LM28*	1	19	5	28	1	20
LM169*	1	20	27	13	24	21
LM142*	1	21	7	1	25	22
CBS 2762	1	25	28	8	14	23
CBS 1596	1	32	21	1	1	24
LM141*	1	33	1	1	1	25
LM92*	1	33	8	20	1	26
LM174*	1	34	2	19	22	27
LM9*	1	35	1	1	14	28
LM14*	2	1	1	1	3	29
LM48*	3	1	1	5	1	30
LM136*	4	9	4	6	15	31
LM17*	5	15	1	10	23	32
LM114*	6	23	1	18	1	33
LM153*	7	17	3	11	21	34
VG1†, VG4†	8	11	22	22	11	35
M41‡	8	13	22	17	4	36
M83‡	8	14	22	17	4	37
M12‡, M123‡, M131‡, M135‡, M166‡, M48‡, M68‡	9	1	26	17	4	38
M14‡	9	1	26	17	5	39
M81‡	10	13	22	17	4	40
LM44*	11	1	1	1	15	41
CBS 397	12	1	8	1	1	42
LM161*	13	26	14	14	20	43
LM15*	13	27	12	15	17	44
LM154*	13	37	16	32	20	45
FM09§	14	31	29	29	15	46
NCYC 179	15	1	19	1	7	47
CBS 6432	15	1	24	1	7	48
UCKM1	15	36	17	33	6	49
CCT 7735	16	22	26	30	15	50
NCYC 1424	16	28	14	9	14	51

Table 3. cont.

Strain name	Genotype					DST
	<i>IPPI</i>	<i>TFC1</i>	<i>GPH1</i>	<i>GSY2</i>	<i>SGA1</i>	
CBS 5795	16	29	14	9	14	52
LM54*	17	12	4	1	1	53
CBS 5668	19	24	15	25	14	54
BO339	18	3	30	34	8	55
KCTC 17555 [†] , CBS 6556	20	5	25	21	13	56
NBRC 1777	21	1	22	22	11	57
CBS 4857	21	10	18	23	9	58
DMKU 3-1042	22	11	22	22	10	59
LM148*	23	11	18	31	26	60
IPE453	22	11	22	22	11	61
CBS 745	24	30	20	26	11	62
DMB1	25	11	18	22	12	63
LM20*	26	18	13	24	19	64
LM167*	27	16	18	27	11	65
CBS 2233, CBS 4354	28	8	23	23	11	66

*From Parmigiano Reggiano, Italy.

†From cow's milk whey, Italy.

‡From Pecorino di Farindola, Italy.

§From fermented milk, Italy.

||From unpasteurized milk, Ireland.

method. The bootstrap analysis was conducted on 1000 replicates.

The MLST data were used to predict genetic exchanges with the program SplitsTree [39]. The split decomposition method was used to assess the degree of tree-like structure present in the alleles found in all the five loci for the strains isolated from the Italian cheeses. Three types of statistical analysis were applied to the data: (1) the *p*-distance to check mean identity to estimate reliability of the alignment; (2) the standardized index of association (I^S_A) to estimate multilocus linkage disequilibrium [40, 41]; (3) the *phi*-test for detecting the presence of recombination. The tests were carried out with Mega 7, START and SplitsTree programs, respectively.

Website development

A website was created to allow users to compare a strain of interest against the 83 *K. marxianus* strains shown in this study, through a maximum-likelihood phylogenetic tree. The website takes a multi FASTA file containing the *IPPI*, *TFC1*, *GPH1*, *GSY2* and *SGA1* sequences as input. Then, it runs a custom Biopython script that computes the phylogenetic tree. The program checks for sequence format, runs a multiple alignment against a prebuilt alignment profile using MUSCLE v 3.8 and reconstructs the phylogenetic tree using PhyML v 3.1. The website was constructed using the HTML v 5.0 mark-up language and CSS v 3. Complementary scripts were written in Biopython v 1.69 and implemented into the website through a CGI Python script.

RESULTS AND DISCUSSION

The availability of an accurate and reproducible method to classify strains within a species is necessary to establish phylogenetic relationships and study biogeography, population structure and evolution. There are a wide variety of methods that have been developed and applied for this purpose but, depending on the method, there can be problems with technical difficulty, poor reproducibility between groups and limited scalability [42, 43]. The most powerful methods are based on DNA sequence variation; thus, comparison of whole genome sequences can be considered the gold standard. However, despite the reduction in sequence costs, this is not always feasible when large numbers of strains are involved and other methods that are rapid, easy to use and scalable, while still accurate, have been applied over the years to study diversity [16–18, 20].

Choice of loci and sequence polymorphisms

The focus of this study was to develop an MLST analysis method for *K. marxianus* that could be used to distinguish strains. The utility of MLST relies on the selectively neutral variability of the housekeeping genes selected [24]; therefore, the choice of genes is critical.

To identify loci that would be suitable for MLST, five genes that had been previously described as housekeeping genes and used as controls in molecular studies in *S. cerevisiae* were considered [44]. The sequences of these genes were retrieved from the seven *K. marxianus* genome sequences available at that time. The dN/dS (non-synonymous to

genotype numbers, it was possible to describe the DSTs as the unique combination of the genotypes of the five loci studied. From the 83 strains, 66 DSTs were defined, whereby 58 strains were the sole representative of their DST, while the other 8 DSTs contained 2–7 representatives. This high level of differentiation confirmed the discriminatory power of MLST as a strain-typing procedure (Table 3).

To investigate the relationship between the 83 strains, the 5 loci of each strain were concatenated and aligned using Muscle. Allelic profiles evaluated by the UPGMA algorithm are shown in Fig. S1. A phylogenetic tree was then reconstructed by the maximum-likelihood method using Unipro UGENE (Fig. 1). This illustrated the diverse nature of the *K. marxianus* species, and showed that the culture collections and sequenced strains are interspersed in different clades. Split decomposition analysis was applied to further analyse the population structure of the 52 *K. marxianus* strains that were isolated from two different Italian cheeses, Parmigiano Reggiano and Pecorino di Farindola. These strains were part of sets previously analysed by different typing methods that did not yield consistency in classification [13, 18, 33]. The concatenated *IPPI1*, *TFC1*, *GPH1*, *GSY2* and *SGA1* gene sequence was analysed and graphically displayed with SplitsTree, which calculates a phylogenetic network that permits the visualization of genetic exchanges in addition to mutations (Fig. 2). The SplitsTree graphs for both the Parmigiano Reggiano strains (Fig. 2a) and the Pecorino di Farindola strains (Fig. 2b) presented network-like structures, separating the strains into groups.

The same groups are found in the maximum-likelihood tree shown in Fig. 1. Concerning the origin of the strains, all *K. marxianus* isolated from Pecorino di Farindola clustered together, although strains M41, M81 and M83 appeared in a separated branch in a similar way as in Figs 1 and 2(b). However, Parmigiano Reggiano *K. marxianus* strains were found to be more different, showing the presence of different groups in Fig. 1. In Fig. 2(a) the majority of the strains clustered together and just seven strains (LM154, LM169, LM161, LM15, LM20, LM167 and LM148) were differentiated. Two different analyses were employed to check for recombination in the two sets of strains. First, the *phi*-test gave *p*-values of 1.337^{-5} with $K=2$ (Parmigiano Reggiano strains) and 2.158^{-5} with $K=1$ (Pecorino di Farindola strains), which provides statistical support for recombination in both groups of strains. Second, the level of linkage disequilibrium between alleles was assessed using the I_A value [45], which is calculated as the ratio of the observed variance with the expected variance [40]. To avoid dependence on the number of loci, the I_A^S , which is expected to be zero when alleles are in linkage equilibrium (free recombination), was used [41]. The I_A^S value was 0.993 for the entire population, indicating the presence of groups of strains corresponding to distinct populations. The I_A^S was then calculated for the two sub-populations separately (Parmigiano Reggiano and Pecorino di Farindola strains). The values were 0.208 and 0.269 for Parmigiano Reggiano and Pecorino di Farindola strains, respectively, indicating recombination within the subpopulations. Both the *phi*-test




Fig. 2. Split decomposition analysis based on the allelic profiles of the 39 strains isolated from Parmigiano Reggiano (a), and the 13 strains isolated from Pecorino di Farindola (b).

and the linkage disequilibrium measure I^S_A are consistent with the idea that there is recombination within these populations of strains. This should not be unexpected since most strains were diploids and *K. marxianus* possesses an intact mating system and can readily be crossed in laboratory studies [19, 46].

To allow the MLST developed here to be applied to more strains, a straightforward website was set up where custom Biopython scripts carry out the typing and generate a phylogenetic tree as well as a PhyML text output file in Newick format. That output file can be imported into other tree visualization software such as MEGA or TreeMe. This website is hosted on a University College Cork server at (kmarxianusMLST.ucc.ie). To type any additional *K. marxianus* strains, the *IPPI*, *TFC1*, *GPH1*, *GSY2* and *SGA1* genes need to be sequenced using the primers specified in Table 2. The sequences are then uploaded in a multi FASTA file and the analysis is automatically performed. Illustrations of the outputs are available as supplementary data (Figs S2, S3 and S4).

In conclusion, in addition to shedding light on the population structure of the *K. marxianus* species, this study provides a framework useful to understanding genetic divergence. Our results showed that MLST of *K. marxianus* is highly reproducible and discriminatory, and superior to other methods used to date to classify strains. To enhance the applicability of the method, a website was constructed to allow typing and comparison of additional *K. marxianus* strains in the future.

Funding information

L.G. was supported by Teagasc Walsh fellowship grant number 2013046. The research leading to these results has received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013, under REA grant agreement no. 606795.

Acknowledgements

We thank Paddy Mulligan for assistance with installing and hosting the MLST server.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Data bibliography

1. The accession numbers of the genome sequences are listed in Table 1.

References

1. Arrizon J, Morel S, Gschaedler A, Monsan P. Fructanase and fructosyltransferase activity of non-*Saccharomyces* yeasts isolated from fermenting musts of Mezcal. *Bioresour Technol* 2012;110:560–565.
2. Lane MM, Morrissey JP. *Kluyveromyces marxianus*: a yeast emerging from its sister's shadow. *Fungal Biol Rev* 2010;24:17–26.
3. Lachance MA. *Kluyveromyces* van der Walt (1971). In: *The Yeasts: A Taxonomic Study*, 5th ed. Amsterdam: NHPC; 2011. pp. 471–481.
4. Mayoral MB, Martín R, Sanz A, Hernández PE, González I et al. Detection of *Kluyveromyces marxianus* and other spoilage yeasts in yoghurt using a PCR-culture technique. *Int J Food Microbiol* 2005;105:27–34.
5. Amaya-Delgado L, Herrera-López EJ, Arrizon J, Arellano-Plaza M, Gschaedler A. Performance evaluation of *Pichia kluyveri*, *Kluyveromyces marxianus* and *Saccharomyces cerevisiae* in industrial tequila fermentation. *World J Microbiol Biotechnol* 2013;29:875–881.
6. Gethins L, Rea MC, Stanton C, Ross RP, Kilcawley K et al. Acquisition of the yeast *Kluyveromyces marxianus* from unpasteurised milk by a kefir grain enhances kefir quality. *FEMS Microbiol Lett* 2016;363:fnw165.
7. Coloretti F, Chiavari C, Luise D, Tofalo R, Fasoli G et al. Detection and identification of yeasts in natural whey starter for Parmigiano Reggiano cheese-making. *Int Dairy J* 2017;66:13–17.
8. Fonseca GG, Heinzle E, Wittmann C, Gombert AK. The yeast *Kluyveromyces marxianus* and its biotechnological potential. *Appl Microbiol Biotechnol* 2008;79:339–354.
9. Morrissey JP, Etschmann MM, Schrader J, de Billerbeck GM. Cell factory applications of the yeast *Kluyveromyces marxianus* for the biotechnological production of natural flavour and fragrance molecules. *Yeast* 2015;32:3–16.
10. Rocha SN, Abrahão-Neto J, Gombert AK. Physiological diversity within the *Kluyveromyces marxianus* species. *Antonie van Leeuwenhoek* 2011;100:619–630.
11. Kurtzman CP, Robnett CJ. Phylogenetic relationships among yeasts of the '*Saccharomyces* complex' determined from multi-gene sequence analyses. *FEMS Yeast Res* 2003;3:417–432.
12. Pérez-Brito D, Tapia-Tussell R, Quijano-Ramayo A, Larqué-Saavedra A, Lappe P. Molecular characterization of *Kluyveromyces marxianus* strains isolated from *Agave fourcroydes* (Lem.) in Yucatan, Mexico. *Mol Biotechnol* 2007;37:181–186.
13. Tofalo R, Fasoli G, Schirone M, Perpetuini G, Pepe A et al. The predominance, biodiversity and biotechnological properties of *Kluyveromyces marxianus* in the production of Pecorino di Farindola cheese. *Int J Food Microbiol* 2014;187:41–49.
14. Lopandic K, Zelger S, Bánszky LK, Eliskases-Lechner F, Prillinger H. Identification of yeasts associated with milk products using traditional and molecular techniques. *Food Microbiol* 2006; 23:341–350.
15. Suzzi G, Lombardi A, Lanorte MT, Caruso M, Andrighetto C et al. Phenotypic and genotypic diversity of yeasts isolated from water-buffalo mozzarella cheese. *J Appl Microbiol* 2000;88:117–123.
16. Sohier D, Le Dizes A-S, Thuault D, Neuveglise C, Coton E et al. Important genetic diversity revealed by inter-LTR PCR fingerprinting of *Kluyveromyces marxianus* and *Debaryomyces hansenii* strains from French traditional cheeses. *Dairy Sci Technol* 2009; 89:569–581.
17. Belloch C, Barrio E, García MD, Querol A. Inter- and intraspecific chromosome pattern variation in the yeast genus *Kluyveromyces*. *Yeast* 1998;14:1341–1354.
18. Fasoli G, Tofalo R, Lanciotti R, Schirone M, Patrignani F et al. Chromosome arrangement, differentiation of growth kinetics and volatile molecule profiles in *Kluyveromyces marxianus* strains from Italian cheeses. *Int J Food Microbiol* 2015;214:151–158.
19. Lane MM, Burke N, Karreman R, Wolfe KH, O'Byrne CP et al. Physiological and metabolic diversity in the yeast *Kluyveromyces marxianus*. *Antonie van Leeuwenhoek* 2011;100:507–519.
20. Belloch C, Barrio E, Uruburu F, Garcia MD, Querol A. Characterisation of four species of the genus *Kluyveromyces* by mitochondrial DNA restriction analysis. *Syst Appl Microbiol* 1997;20: 397–408.
21. Masneuf-Pomarede I, Salin F, Börlin M, Coton E, Coton M et al. Microsatellite analysis of *Saccharomyces uvarum* diversity. *FEMS Yeast Res* 2016;16:fow002.
22. Hennequin C, Thierry A, Richard GF, Lecointre G, Nguyen HV et al. Microsatellite typing as a new tool for identification of *Saccharomyces cerevisiae* strains. *J Clin Microbiol* 2001;39:551–559.
23. Pérez-Brito D, Tapia-Tussell R, Quijano-Ramayo A, Larqué-Saavedra A, Lappe P. Molecular characterization of

- Kluyveromyces marxianus* strains isolated from *Agave fourcroydes* (Lem.) in Yucatan, Mexico. *Mol Biotechnol* 2007;37:181–186.
24. Cooper JE, Feil EJ. Multilocus sequence typing – what is resolved? *Trends Microbiol* 2004;12:373–377.
 25. Taylor JW, Fisher MC. Fungal multilocus sequence typing – it's not just for bacteria. *Curr Opin Microbiol* 2003;6:351–356.
 26. Maiden MC. Multilocus sequence typing of bacteria. *Annu Rev Microbiol* 2006;60:561–588.
 27. Fay JC, Benavides JA. Evidence for domesticated and wild populations of *Saccharomyces cerevisiae*. *PLoS Genet* 2005;1:e5–71.
 28. Fay JC, Benavides JA. Hypervariable noncoding sequences in *Saccharomyces cerevisiae*. *Genetics* 2005;170:1575–1587.
 29. Aa E, Townsend JP, Adams RI, Nielsen KM, Taylor JW et al. Population structure and gene evolution in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 2006;6:702–715.
 30. Bounoux ME, Morand S, D'Enfert C. Usefulness of multilocus sequence typing for characterization of clinical isolates of *Candida albicans*. *J Clin Microbiol* 2002;40:1290–1297.
 31. Bounoux ME, Tavanti A, Bouchier C, Gow NA, Magnier A et al. Collaborative consensus for optimized multilocus sequence typing of *Candida albicans*. *J Clin Microbiol* 2003;41:5265–5266.
 32. Tavanti A, Gow NA, Senesi S, Maiden MC, Odds FC. Optimization and validation of multilocus sequence typing for *Candida albicans*. *J Clin Microbiol* 2003;41:3765–3776.
 33. Fasoli G, Barrio E, Tofalo R, Suzzi G, Belloch C. Multilocus analysis reveals large genetic diversity in *Kluyveromyces marxianus* strains isolated from Parmigiano Reggiano and Pecorino di Farindola cheeses. *Int J Food Microbiol* 2016;233:1–10.
 34. Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 2016;33:1870–1874.
 35. Okonechnikov K, Golosova O, Fursov M, UGENE team. Unipro UGENE: a unified bioinformatics toolkit. *Bioinformatics* 2012;28:1166–1167.
 36. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 2004;32:1792–1797.
 37. Odds FC, Jacobsen MD. Multilocus sequence typing of pathogenic *Candida* species. *Eukaryot Cell* 2008;7:1075–1084.
 38. Jolley KA, Feil EJ, Chan MS, Maiden MC. Sequence type analysis and recombinational tests (START). *Bioinformatics* 2001;17:1230–1231.
 39. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol* 2006;23:254–267.
 40. Smith JM, Smith NH, O'Rourke M, Spratt BG. How clonal are bacteria? *Proc Natl Acad Sci USA* 1993;90:4384–4388.
 41. Haubold B, Hudson RR. LIAN 3.0: detecting linkage disequilibrium in multilocus data. Linkage analysis. *Bioinformatics* 2000;16:847–849.
 42. Saghrouni F, Ben Abdeljelil J, Boukadida J, Ben Said M. Molecular methods for strain typing of *Candida albicans*: a review. *J Appl Microbiol* 2013;114:1559–1574.
 43. Pérez MA, Gallego FJ, Hidalgo P. Evaluation of molecular techniques for the genetic characterization of *Saccharomyces cerevisiae* strains. *FEMS Microbiol Lett* 2001;205:375–378.
 44. Teste MA, Duquenne M, François JM, Parrou JL. Validation of reference genes for quantitative expression analysis by real-time RT-PCR in *Saccharomyces cerevisiae*. *BMC Mol Biol* 2009;10:99.
 45. Maiden MC, Bygraves JA, Feil E, Morelli G, Russell JE et al. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci USA* 1998;95:3140–3145.
 46. Yarimizu T, Nonklang S, Nakamura J, Tokuda S, Nakagawa T et al. Identification of auxotrophic mutants of the yeast *Kluyveromyces marxianus* by non-homologous end joining-mediated integrative transformation with genes from *Saccharomyces cerevisiae*. *Yeast* 2013;30:485–500.

Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at microbiologyresearch.org.