

Effect of *Foeniculum vulgare* Mill. (fennel) on menopausal symptoms in postmenopausal women: a randomized, triple-blind, placebo-controlled trial

Fatemeh Rahimikian, MSc,^{1,2} Roja Rahimi, PhD,^{3,4} Parvin Golzareh, MSc,⁵ Reza Bekhradi, MD,⁶ and Abbas Mehran, MSc⁷

Abstract

Objective: Preliminary data suggest that *Foeniculum vulgare* (fennel) can be an effective treatment for menopausal symptoms. This trial was designed to assess the efficacy of fennel in the management of menopausal symptoms in postmenopausal women.

Methods: In this triple-blind, placebo-controlled trial, 90 postmenopausal women aged 45 to 60 years in Tehran were randomly assigned to treatment (n = 45) or placebo (n = 45) groups. The participants received 8 weeks of treatment with soft capsules containing 100 mg fennel or a placebo (2 per day for each group). The participants were followed for 2 weeks postintervention to assess the continuance of the effect of intervention. The Menopause Rating Scale (MRS) questionnaire was used to assess changes in menopausal symptoms at baseline and at 4, 8, and 10 weeks after onset of intervention.

Results: The groups recorded similar mean scores on the MRS questionnaire before intervention. After intervention, the treatment group showed a significant decrease in the mean MRS score. The results of the Friedman test showed significant differences between the mean score at baseline and those at 4, 8, and 10 weeks after onset of intervention in the treatment group ($P < 0.001$), whereas there were no significant differences in the placebo group. When the fennel and the placebo groups were compared, the independent t test showed significant differences in mean scores between groups at 4, 8, and 10 weeks (2 weeks postintervention; $P < 0.001$).

Conclusions: Fennel is an effective and safe treatment to reduce menopausal symptoms in postmenopausal women without serious side effects. More clinical trials with larger populations are required to confirm this result.

Key Words: Fennel – *Foeniculum vulgare* – Menopause Rating Scale – Menopausal symptoms.

Menopause is the period in the life of an adult woman in which the ovaries gradually become inactive and menstruation ceases.¹ The average age of US women at menopause is 51 years²; however, in

some countries, the age of menopause is lower. For example, the average age of menopause for Indian women is 46.2 years³ and for Iranian women is 48.2 years.^{4,5} Most women will spend more than one-third of their lives in a postmenopausal period.²⁻⁵ The aging of the global population means that millions of women worldwide enter menopause every year. Although some women have few problems, three out of four will experience symptoms resulting from changes in estrogen and progesterone levels during this time. One in four will experience major symptoms that can affect quality of life.^{6,7}

The postmenopausal symptoms can be vasomotor, urogenital, and psychological.^{1,2,7} Research has shown that the most prevalent complaints reported by postmenopausal women are joint and muscular discomfort, physical and mental exhaustion, sleeping problems, hot flashes, sweating, irritability, dryness of the vagina, anxiety, and depression.⁸⁻¹⁰ Hormone therapy (HT) with estrogen and progesterone can improve these symptoms, but can lead to serious side effects such as breast and endometrial cancer and cardiovascular disease.^{2,11,12} Globally, women have turned to complementary and alternative medicine (CAM) for relief of their menopausal symptoms. The most popular CAM modality is herbal medicine (especially phytoestrogens and evening

Received November 22, 2016; revised and accepted January 30, 2017.

From the ¹Nursing and Midwifery Care Research Center, ²Department of Midwifery, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran; ³Department of Traditional Pharmacy, School of Traditional Medicine, Tehran University of Medical Sciences, Tehran, Iran; ⁴Phyto Pharmacology Interest Group (PPIG), Universal Scientific Education and Research Network (USERN), Tehran, Iran; ⁵School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran; ⁶Barij Medicinal Herbs Research Center, Kashan, Iran; and ⁷Statistical Department, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran.

Author contributions: F.R. had designed and managed the study. P.G. conducted the study and wrote the paper. R.R. had a role in traditional and herbal medicine counseling, handled pharmaceutical part of project and edited the manuscript. R.B. handled pharmaceutical part of project. A.M. analyzed data.

Funding/support: This study has been partially supported by the Nursing and Midwifery Care Research Center of Tehran University of Medical Sciences (TUMS); Grant No 31096-99-04-94.

Financial disclosure/conflicts of interest: None reported.

Address correspondence to: Parvin Golzareh, MSc, Nursing and Midwifery Faculty, Tehran University of Medical Sciences (TUMS), Tehran 13185-1678, Iran. E-mail: golzareh23@yahoo.com

primrose oil).¹³⁻¹⁵ Phytoestrogens and a phytoestrogenic diet can help prevent and treat menopausal symptoms such as skin aging, osteoporosis, cancer, cardiovascular disease, vaginal atrophy, sleep disturbances, and cognitive disorders.¹⁶⁻¹⁸

Foeniculum vulgare Mill., commonly known as fennel, is a medicinal plant from the family Umbelliferae that is used all over the world. Fennel fruit contains essential oils that are mainly composed of trans-anethole, fenchone, and estragole. The fruit also contains phenolic compounds including flavonoids (flavonoid glycoside and flavonoid aglycon), phenolic acid, hydroxycinnamic acid, coumarin, and tannin.¹⁹ Fennel fruit has phytoestrogenic properties and has been effective for increasing breast milk,²⁰ improving premenstrual symptoms,²¹ decreasing climacteric symptoms in men, increasing libido,²² and improving menopausal symptoms and sexual activity in postmenopausal women.^{23,24} Few clinical studies have examined the effect of fennel on menopausal symptoms. This study was designed to evaluate the effects of an herbal preparation made from fennel on the menopausal symptoms of postmenopausal women.

METHODS

Study design

This study was a randomized, triple-blind, placebo-controlled clinical trial that was approved by the Ethics Committee of Tehran University of Medical Sciences and has been registered in the Iranian Registry of Clinical Trials (code: IRCT2016050827788N1). The aim of this study was to determine the effects of fennel on menopausal symptoms in postmenopausal women.

Participants

This study follows similar previous studies on the effect of phytoestrogens on menopausal symptoms which hypothesized $P_0 = 50\%$ (ratio of severe menopausal symptoms) and projected a decrease in symptoms of 20% ($P_1 = 20\%$) after intervention. Assuming $\alpha = 0.05$ (confidence interval [CI] = 95%) and $\beta = 0.2$ (statistical power = 80%), with the use of the following theorem, a total of 40 participants was calculated for each group. To accommodate possible withdrawal of participants during the study, 45 participants were enrolled in each group.

$$n = \frac{2 \times (Z_{1-\alpha/2} + Z_{1-\beta})^2 \times pq}{(P_0 - P_1)^2}$$

$$= \frac{2 \times (1.96 + 0.842)^2 \times 0.35 \times 0.65}{(0.5 - 0.2)^2} = 40 \quad (1)$$

$$(P = \frac{P_0 + P_1}{2} = 0.35 \rightarrow q = 0.65)$$

The participants were selected from women referred to the Farmanfarmayian Public Health Center affiliated with Tehran University of Medical Sciences in Tehran, Iran, in 2016. This center was selected because it had the highest number of postmenopausal women referred for testing of blood sugar,

lipids, and blood pressure. The inclusion criteria were women aged 45 to 60 years who were married and in the first 1 to 5 years of the postmenopausal period (a woman was defined as postmenopausal beginning 1 year after her last menstrual period).²⁵ They had to achieve a minimum score of 9 on the Menopause Rating Scale (MRS) questionnaire (medium severity of menopausal symptoms) and also had to have a negative history of physical or psychological disease, HT, CAM for menopausal symptoms, allergies to herbal medicine, sedative or antidepressant drug use, addiction, and smoking. The participants were required to be able to read and write. Exclusion criteria were allergies to fennel/placebo during the intervention, worsening of symptoms during the intervention, poor cooperation, failure to use fennel/placebo for a total of 6 days,²⁶ and use of other remedies for menopausal symptoms during the study.

Randomization

After approval and registration of the proposed study plan by the Ethics Committee of Tehran University of Medical Sciences, the study sample of 90 women was selected. The randomization method was the use of random numbers as implemented on the Randomizer website (www.randomizer.org). A set of 45 random numbers between 1 and 90 was generated by the site and assigned to group A. The remaining numbers between 1 and 90 were assigned to group B. Using the numbers allocated to each group, the participants were divided into the A and B groups (each with 45 participants). This randomly generated two groups of 45 women.

Intervention

The soft fennel and placebo capsules were produced by Barij Essence Pharmaceutical Company (Iran). The essential oil was derived from fennel fruits using steam distillation, and the fennel essential oil (FEO) was formulated into soft capsules. Each 100 mg soft capsule contained 30% FEO (71-90 mg anethole) and 0.02% butylated hydroxytoluene, combined with qs to 100% wt/wt with sunflower oil. There was no difference between the fennel and placebo capsules in terms of color, odor, shape, and size, but the placebo capsules were fully filled with sunflower oil. The study was triple-blind such that the participants, investigators, and the person analyzing the data were blind. Women in the intervention group received two soft capsules containing 100 mg of fennel daily for 8 weeks, whereas those in the placebo group received two soft capsules containing 100 mg of sunflower oil daily for 8 weeks. None of the participants in either group engaged in other interventions during the study.

Outcome measures

Sociodemographic data of the women participating in the study were collected by questionnaire. The menopausal symptoms were assessed as total symptoms, somatovegetative symptoms (hot flashes, heart discomfort, sleep problems, and joint or muscular discomfort), psychological symptoms (depression, irritability, anxiety, and physical or mental exhaustion), and urogenital symptoms (sexual dysfunction,




FIG. 1. Consolidated standards of reporting trials diagram reflecting flow of participants through the study.

bladder problems, and vaginal dryness) using the MRS questionnaire at baseline and at weeks 4, 8, and 10 of the study. The MRS is a standardized and formally validated scale that is used worldwide to measure the severity of 11 symptoms associated with menopause and to assess health-related quality of life of menopausal women.^{27,28}

At the onset of the study and after completing the demographic and MRS questionnaires, the soft capsules for the first 4 weeks of intervention and a checklist to record daily capsule consumption were given to the participants. At the end of the first 4 weeks, a second meeting was set in which the participants handed in the first checklist, completed the second MRS questionnaire, and received the capsules and checklist for the second 4 weeks. At the end of the second phase, women participated in the third meeting, again completing the MRS questionnaire and the second checklist was handed in. At completion of 8 weeks of intervention, no further capsules or checklists were provided, but participants again completed the MRS questionnaire at 10 weeks (10 weeks after the onset of intervention).

One researcher called participants weekly to ask about capsule consumption and possible side effects. Participants were provided with the researcher’s phone number should any questions or problems arise.

Statistical analysis

The data were analyzed using SPSS software (version 22). Participants who did not complete the trial were not included in the statistical analysis. The independent *t* test was used to compare the means between groups, and the Friedman test was used to compare the means of total scores before and after treatment. Demographic and clinical characteristics were analyzed using Fisher’s exact test, and an α value of 0.05 was considered statistically significant.

RESULTS

In all, 40 women in the intervention and 39 women in the placebo group completed the study (Fig. 1). No serious adverse events were recorded during or after the intervention.

The minor side effects were one case of allergic reaction and one case of feeling severe heat in the intervention group and one case of digestive problems in the control group. The epidemiological and clinical characteristics of age, menopausal age, number of pregnancies, number of children, body mass index (BMI), education, occupation, and economic status were similar in both groups and there were no significant differences between groups (Table 1).

In the intervention group, the mean MRS score was 20.02 ± 6.18 at baseline, which decreased to 11.20 ± 4.92 at week 4, 9.35 ± 4.54 at week 8, and 13.05 ± 4.94 at week 10. There was a significant statistical difference between the scores at baseline and weeks 4, 8, and 10 ($P < 0.001$; Table 2). In the control group, the mean MRS score was 20.37 ± 5.51 at baseline, which decreased to 19.23 ± 5.42 at week 4, 18.58 ± 6.11 at week 8, and 19.20 ± 5.97 at week 10. There were no significant differences between scores at baseline and at weeks 4, 8, and 10 ($P = 0.402$; Table 2) in the control group. When the fennel and the placebo groups were compared, there was no significant difference in mean MRS score at baseline

TABLE 1. Demographic and clinical characteristics of menopausal women and their comparisons in two studied groups

Group characteristic	<i>Foeniculum vulgare</i> (n = 40)	Placebo (n = 39)	P
Age (mean \pm SD)	52.33 \pm 3.12	51.91 \pm 3.72	0.563
Menopausal age (mean \pm SD)	49.20 \pm 2.69	48.22 \pm 3.22	0.122
Number of pregnancies (mean \pm SD)	3.18 \pm 1.41	3.11 \pm 1.265	0.815
Number of children (mean \pm SD)	3.07 \pm 1.38	2.84 \pm 1.04	0.393
BMI (mean \pm SD)	28.33 \pm 3.47	27.14 \pm 4.87	0.073
Education status			
Primary or guidance school (%)	57.8	55.5	
High school or diploma (%)	37.8	35.6	0.797
Academic (%)	4.4	8.9	
Job status			
Homemaker (%)	97.8	95.6	1
Employed (%)	2.2	4.4	
Economic status			
Low (%)	11.1	11.1	
Middle (%)	75.6	84.5	0.376
High (%)	13.3	4.4	

BMI, body mass index; SD, standard deviation.

TABLE 2. Comparison of total score of Menopause Rating Scale (MRS) questionnaire between the two groups in four different periods of treatment

Group	Before intervention		4 weeks after intervention		8 weeks after intervention		10 weeks after intervention (2 weeks after cutting the intervention)		P
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
<i>Foeniculum vulgare</i> (n = 40)	20.02	6.18	11.20	4.92	9.35	4.54	13.05	4.94	<0.001
Placebo (n = 39)	20.37	5.51	19.23	5.42	18.58	6.11	19.20	5.97	0.402
P	0.77		<0.001		<0.001		<0.001		

SD, standard deviation.

between the two groups ($P > 0.05$; Table 2), but there were significant differences in the scores between groups at weeks 4, 8, and 10 ($P \leq 0.001$; Table 2), and scores in the fennel group were lower than the placebo group.

DISCUSSION

Phytoestrogens are estrogen-like chemicals in plants¹ that have been employed as a treatment for menopausal symptoms. Studies have demonstrated the estrogenic properties of fennel. It has been shown that fennel can increase the weight of the mammary glands, oviducts, cervix, and vagina in rats.²⁹ Human studies have shown that fennel can increase breast milk,²⁰ decrease hirsutism,³⁰ improve dysmenorea,³¹ and treat amenorea³² in women. Fennel has also been shown to improve menopausal symptoms and sexual activity in postmenopausal women.²³

In this randomized, placebo-controlled clinical trial, the effects of fennel fruit as a medicinal herb with phytoestrogenic properties was investigated in postmenopausal women suffering from climacteric complaints. A significant improvement was observed in menopausal symptoms during the 8 weeks of intervention and 2 weeks of follow-up in fennel group. At week 10 (2 weeks postintervention), the total score had increased slightly in the fennel group, but was still markedly lower when compared with baseline.

Studies have reported that phytoestrogens can reduce the mean MRS score. The results of one 12-week study indicated that soy-derived isoflavone treatment improved general menopausal symptoms and decreased MRS scores when compared with baseline.³³ Another 12-week study that examined the effect of red clover on menopausal symptoms showed that the total MRS score decreased significantly in the intervention group.³⁴ Rostock et al³⁵ found a reduction in total MRS score after treatment with black cohosh.

An 8-week placebo-controlled study investigated the effect of fennel and St John's wort (*Hypericum perforatum*) on climacteric symptoms. Both fennel and St John's wort were found to significantly reduce menopausal symptoms when compared with placebo.²³ Another 8-week study indicated that fennel vaginal cream was effective for vaginal atrophy in postmenopausal women.²⁴

The results of the present study indicate that fennel significantly reduced the total MRS score of the intervention group when compared with the placebo group, which is

similar to the findings of the foregoing studies. Although many studies have confirmed the positive effects of phytoestrogens on postmenopausal symptoms,^{36,37} some have shown no difference between the phytoestrogens and the placebo. One study found that pomegranate seed oil did not significantly reduce hot flashes over a 12-week observation period.³⁸ Amato et al³⁹ reported that soy isoflavone supplementation offered no benefit to quality of life in postmenopausal women. del Giorno et al⁴⁰ found that 12 months of treatment with *Trifolium pratense* had no significant effect on menopausal symptoms and sexual satisfaction. Liu et al⁴¹ reported that whole soy and purified daidzein had no significant effect on alleviation of menopausal symptoms.

The limitations of this study were its short duration and small sample size. Studies with longer durations and larger populations are required to confirm the results of this study.

CONCLUSIONS

The results of this study show that fennel as a phytoestrogen is effective in reducing menopausal symptoms with no serious side effects. It can be considered for decreasing the menopausal symptoms in women with low estrogen levels, and those who have experienced early menopause or have had a hysterectomy or oophorectomy. Further clinical trials with a larger population and of longer duration are recommended to obtain more conclusive results.

Acknowledgments: We would like to thank Tehran University of Medical Sciences for the research grant, the helpful personnel of the Farmanfarmayian Public Health Center, and the patients for all their cooperation and support.

REFERENCES

1. Fritz MA, Speroff L. *Clinical Gynecology Endocrinology and Infertility*. 8th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2011:673-857.
2. Berek JS. *Berek and Novak's Gynecology*. 15th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2012:1320-1340.
3. Ahuja M. Age of menopause and determinants of menopause age: a PAN India survey by IMS. *J Midlife Health* 2016;7:126-131.
4. Rajaeefard A, Mohammad-Beigi A, Mohammad-Salehi N. Estimation of natural age of menopause in Iranian women: a meta-analysis study. *Koomesh* 2011;13:1-7.
5. Nahidi F, Karman N, Vallaei N, Fazli Z. Studying incidence of menopause and its effective factors in Tehran. *Res Med* 2010;33:258-265.
6. Bayer HealthCare Pharmaceuticals. Menopausal complaints. Available at: <http://pharma.bayer.com/en/treatment-care/womens-health/menopausal-complaints/>. Accessed November 17, 2016.

7. Manson JE, Bassuk SS. *Hot Flashes, Hormones, and Your Health (Harvard Medical School Guides). Hard Cover Edition*. New York: McGraw-Hill; 2007;1-17.
8. Rahman SA, Zainudin SR, Mun VL. Assessment of menopausal symptoms modified Menopause Rating Scale (MRS) among middle age women in Kuching, Sarawak, Malaysia. *Asia Pac Fam Med* 2010;9:5.
9. Ashrafi M, Kazemi Ashtiani S, Malekzadeh F, et al. Symptoms of natural menopause among Iranian women living in Tehran, Iran. *Iran J Reproduct Med* 2010;8:29-32.
10. Jin F, Tao M, Teng Y, Shao H, Li C, Mills E. Knowledge and attitude towards menopause and hormone replacement therapy in Chinese woman. *Gynecol Obstet Invent* 2015;79:5-40.
11. Edelman J. *Menopause Matters*. United States of America; 2010;41-53.
12. Sjögren LL, Mørch LS, Løkkegaard E. Hormone replacement therapy and the risk of endometrial cancer: a systematic review. *Maturitas* 2016; 91:25-35.
13. Buhling KJ, Daniels BV, Studnitz FS, Eulenbug C, Mueck AO. The use of complementary and alternative medicine by women transitioning through menopause in Germany: results of a survey of women aged 45-60 years. *Complement Ther Med* 2014;22:94-98.
14. Cardini F, Lesi G, Lombardo F, et al. The use of complementary and alternative medicine by women experiencing menopausal symptoms in Bologna. *BMC Womens Health* 2010;10:7.
15. Posadzki P, Lee MS, Moon TW, Choi TY, Park TY, Ernst E. Prevalence of complementary and alternative medicine (CAM) use by menopausal women: a systematic review of surveys. *Maturitas* 2013;75:34-43.
16. Sirotkin AV, Harrath AH. Phytoestrogens and their effects. *Eur J Pharmacol* 2014;741:230-236.
17. Bedell S, Nachtigall M, Naftolin F. The pros and cons of plant estrogens for menopause. *J Steroid Biochem Mol Bio* 2014;139:225-236.
18. Franco OH, Chowdhury R, Troup J, et al. Use of plant-based therapies and menopausal symptoms: a systematic review and meta-analysis. *JAMA* 2016;315:2554-2563.
19. Rahimi R, Ardekani MR. Medicinal properties of *Foeniculum vulgare* Mill. In traditional Iranian medicine and modern phytotherapy. *Chin J Integr Med* 2013;19:73-79.
20. Honarvar F, Tadayon M, Afshari P, Namjooyan F, Haghghi M. The effect of *Foeniculum vulgare* on serum prolactin level in lactating women. *Iran J Obstet Gynecol Infertil* 2013;16:18-24.
21. Delaram M, Heydarnejad S. Herbal remedy for premenstrual syndrome with fennel (*Foeniculum vulgare*): randomized, placebo-controlled study. *Adv Clin Exp Med* 2011;20:509-512.
22. Puleo M. Fennel and anise as estrogenic agents. *J Ethnopharmacol* 1980;2:337-344.
23. Abdali K, Dowran P, Emamghoreishi M, Kasraian M, Tabatabaei H. Comparison of the effect of *Foeniculum vulgare* and St John's wort (*Hypericum perforatum*) on the climacteric symptoms and sexual activity in menopausal woman. *IJBR* 2016;7:148-154.
24. Yaralizadeh M, Abedi P, Najari S, Namjooyan F, Saki A. Effect of *Foeniculum vulgare* (fennel) vaginal cream on vaginal atrophy in postmenopausal women: a double-blind randomized placebo-controlled trial. *Maturitas* 2016;84:75-80.
25. Diagnosis of perimenopause and menopause. In: Menopause: Full Guideline. National Collaborating Centre for Women's and Children's Health (UK). London: National Institute for Health and Care Excellence (UK); NICE Guideline, 2015;23:49.
26. Tice JA, Ettinger B, Ensrud K, Wallace R, Blackwell T, Cummings SR. Phytoestrogen supplements for the treatment of hot flashes: the Isoflavone Clover Extract (ICE) study: a randomized controlled trial. *JAMA* 2003; 290:207-213.
27. Heinemann Lothar AJ, Potthoff P, Schneider Hermann PG. International versions of the Menopause Rating Scale (MRS). *Health Qual Life Outcomes* 2003;1:1-4.
28. Tao M, Shao H, Li C, Teng Y. Correlation between the modified Kupperman Index and the Menopause Rating Scale in Chinese women. *Patient Prefer Adherence* 2013;7:223-229.
29. Malini T, Vanithakumari G, Megala N, Anusya S, Devi K, Elango V. Effect of *Foeniculum vulgare* Mill. seed extract on the genital organs of male and female rats. *Indian J Physiol Pharmacol* 1985;29: 21-26.
30. Javidnia K, Dastgheib L, Mohammadi Samani S, Nasiri A. Antihirsutism activity of fennel (fruits of *Foeniculum vulgare*) extract: a double-blind placebo controlled study. *Phytomedicine* 2003;10:455-458.
31. Ghodsi Z, Asltooghi M. The effect of fennel on pain quality, symptoms and menstrual duration in primary dysmenorrhea. *J Pediatr Adolesc Gynecol* 2014;27:283-286.
32. Mohebbi-Kian E, Mohammad-Alizadeh-Charandabi S, Bekhradi R. Efficacy of fennel and combined oral contraceptive on depot medroxyprogesterone acetate-induced amenorrhea: a randomized placebo-controlled trial. *Contraception* 2014;90:440-446.
33. Chedraui P, San Miguel G, Schwager G. The effect of soy-derived isoflavones over hot flushes, menopausal symptoms and mood in climacteric women with increased body mass index. *Gynecol Endocrinol* 2011; 27:307-313.
34. Shakeri F, Taavoni S, Goushegir A, Haghani H. Effectiveness of red clover in alleviating menopausal symptoms: a 12-week randomized, controlled trial. *Climacteric* 2015;18:568-573.
35. Rostock M, Fischer J, Mumm A, Stammwitz U, Saller R, Bartsch HH. Black cohosh (*Cimicifuga racemosa*) in tamoxifen-treated breast cancer patients with climacteric complaints: a prospective observational study. *Gynecol Endocrinol* 2011;27:844-848.
36. Thomas AJ, Ismail R, Taylor-Swanson L, et al. Effects of isoflavones and amino acid therapies for hot flashes and co-occurring symptoms during the menopausal transition and early postmenopause: a systematic review. *Maturitas* 2014;78:263-276.
37. Chen M-N, Lin C-c, Liu C-f. Efficacy of phytoestrogens for menopausal symptoms: a meta-analysis and systematic review. *Climacteric* 2015; 18:260-269.
38. Auerbach L, Rakus J, Bauer C, et al. Pomegranate seed oil in women with menopausal symptoms: a prospective randomized, placebo-controlled, double-blinded trial. *Menopause* 2012;19:426-432.
39. Amato P, Young RL, Steinberg FM, et al. Effect of soy isoflavone supplementation on menopausal quality of life. *Menopause* 2013;20: 443-447.
40. del Giorno C, Fonseca AM, Bagnoli VR, Assis JS, Soares JM Jr, Baracat EC. Effects of *Trifolium pratense* on the climacteric and sexual symptoms in postmenopause women. *Rev Assoc Med Bras (1992)* 2010;56: 558-562.
41. Liu ZM, Ho SC, Woo J, Chen YM, Wong C. Randomized controlled trial of whole soy and isoflavone daidzein on menopausal symptoms in equal-producing Chinese postmenopausal women. *Menopause* 2014;21: 653-660.