

Regulation of Adipocyte 11 β -Hydroxysteroid Dehydrogenase Type 1 (11 β -HSD1) by CCAAT/Enhancer-Binding Protein (C/EBP) β Isoforms, LIP and LAP

Cristina L. Esteves¹, Val Kelly¹, Valérie Bégay³, Tak Y. Man¹, Nicholas M. Morton², Achim Leutz³, Jonathan R. Seckl¹, Karen E. Chapman^{1*}

1 Endocrinology Unit, University/BHF Centre for Cardiovascular Science, Queen's Medical Research Institute, The University of Edinburgh, Edinburgh, United Kingdom, **2** Molecular Metabolism Group, University/BHF Centre for Cardiovascular Science, Queen's Medical Research Institute, The University of Edinburgh, Edinburgh, United Kingdom, **3** Max Delbrück Center for Molecular Medicine, Berlin, Germany

Abstract

11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1) catalyses intracellular regeneration of active glucocorticoids, notably in liver and adipose tissue. 11 β -HSD1 is increased selectively in adipose tissue in human obesity, a change implicated in the pathogenesis of metabolic syndrome. With high fat (HF)-feeding, adipose tissue 11 β -HSD1 is down-regulated in mice, plausibly to counteract metabolic disease. Transcription of 11 β -HSD1 is directly regulated by members of the CCAAT/enhancer binding protein (C/EBP) family. Here we show that while total C/EBP β in adipose tissue is unaltered by HF diet, the ratio of the C/EBP β isoforms liver-enriched inhibitor protein (LIP) and liver-enriched activator protein (LAP) (C/EBP β -LIP:LAP) is increased in subcutaneous adipose. This may cause changes in 11 β -HSD1 expression since genetically modified C/EBP $\beta^{(+/-)}$ mice, with increased C/EBP β -LIP:LAP ratio, have decreased subcutaneous adipose 11 β -HSD1 mRNA levels, whereas C/EBP $\beta^{\Delta\text{ORF}}$ mice, with decreased C/EBP β -LIP:LAP ratio, show increased subcutaneous adipose 11 β -HSD1. C/EBP β -LIP:LAP ratio is regulated by endoplasmic reticulum (ER) stress and mTOR signalling, both of which are altered in obesity. In 3T3-L1 adipocytes, 11 β -HSD1 mRNA levels were down-regulated following induction of ER stress by tunicamycin but were up-regulated following inhibition of mTOR by rapamycin. These data point to a central role for C/EBP β and its processing to LIP and LAP in transcriptional regulation of 11 β -HSD1 in adipose tissue. Down-regulation of 11 β -HSD1 by increased C/EBP β -LIP:LAP in adipocytes may be part of a nutrient-sensing mechanism counteracting nutritional stress generated by HF diet.

Citation: Esteves CL, Kelly V, Bégay V, Man TY, Morton NM, et al. (2012) Regulation of Adipocyte 11 β -Hydroxysteroid Dehydrogenase Type 1 (11 β -HSD1) by CCAAT/Enhancer-Binding Protein (C/EBP) β Isoforms, LIP and LAP. PLoS ONE 7(5): e37953. doi:10.1371/journal.pone.0037953

Editor: Kang Sun, Fudan University, China

Received: February 28, 2012; **Accepted:** April 30, 2012; **Published:** May 25, 2012

Copyright: © 2012 Esteves et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a Wellcome Trust Programme Grant (grant number WT083184 to JRS and KEC). NMM was funded by a Wellcome Trust Career Development Fellowship (grant 079660/z/06/z). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Karen.Chapman@ed.ac.uk

Introduction

11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1) is highly expressed in liver and adipose tissue where it catalyses the regeneration of active glucocorticoids (corticosterone, cortisol) from inert 11keto- forms (11-dehydrocorticosterone, cortisone) thus increasing intracellular glucocorticoid action [1]. 11 β -HSD1 expression is elevated selectively in adipose tissue of obese humans and in monogenic rodent genetic obesity, whereas levels in liver are unaffected or even decreased [2,3,4]. Transgenic over-expression of 11 β -HSD1 in adipose tissue recapitulates the metabolic syndrome in mice, with visceral obesity, dyslipidemia, insulin resistance/diabetes and hypertension [2,5]. In contrast, 11 β -HSD1-deficiency or inhibition causes insulin-sensitization (including in humans), lowers fasting plasma glucose and lipid levels, reduces visceral adipose tissue mass and attenuates atherosclerosis [6,7,8]. Unexpectedly, high fat (HF) diet down-regulated 11 β -HSD1 selectively in adipose tissue in mice and rats [9,10,11]. This down-regulation is greatest in obesity-resistant strains [9] suggesting it may be a mechanism to minimise metabolic disease with adiposity. Understanding the mechanisms

of adipose-specific control of 11 β -HSD1 is crucial to dissecting the pathogenesis of sensitivity/resistance to obesity.

Transcription of 11 β -HSD1 is directly regulated by members of the CCAAT/enhancer binding protein (C/EBP) family of transcription factors in all tissues and cells studied [12,13,14,15,16]. The family comprises 6 members; C/EBP α , β , δ , γ , ϵ and ζ (or CHOP) [17]. C/EBP α , β , δ and CHOP are essential for adipocyte differentiation and function *in vitro* and *in vivo* [18,19,20,21]. C/EBP α and β each occur as distinct isoforms arising from differential translation initiation or proteolysis [22,23,24]. C/EBP α produces 42 kDa (p42) and 30 kDa (p30) isoforms [22,24,25]. C/EBP β exists as three major isoforms; the 38 kDa and 35 kDa liver-enriched activator protein isoforms (LAP* and LAP, respectively), which stimulate transcription, and the 20 kDa liver-enriched inhibitor protein (LIP) [24]. LIP lacks the transcription activation domain of C/EBP β and is typically a dominant-negative regulator of C/EBP function [24]. The LIP:LAP ratio is thus an important determinant of C/EBP β action [26,27,28].

In vivo, C/EBP α is the major known inducer of 11 β -HSD1 transcription in liver, where C/EBP β acts as a relative repressor

[12]. In contrast, in adipose tissue, C/EBP β is an activator [29]. Moreover, C/EBP β is required for glucocorticoid-induction of 11 β -HSD1 in A549 cells [16] as well as regulation by IL-1 β , cAMP and ceramide in fibroblasts and 3T3-L1 preadipocytes [13,14,30]. However, the crucial transcriptional regulation of 11 β -HSD1 in adipose tissue and mature adipocytes remains unexplored. Here, we have tested the hypothesis that C/EBPs are modulated by diet in mice and mediate the regulation of 11 β -HSD1 in adipose.

Materials and Methods

Animals

C57BL/6J mice (Harlan UK Ltd., Oxon, UK) were housed in standard conditions on a 12 h light, 12 h dark cycle (lights on at 0700 h) at $21 \pm 1^\circ\text{C}$. Adult male mice ($n = 16/\text{group}$) were fed control diet (11% calories as fat; diet D12328, Research Diets, Inc., New Brunswick, NJ) or HF diet (58% calories as fat; diet D123331, Research Diets) for 6 weeks. Adult male C/EBP β mutant mice, heterozygous C/EBP $\beta^{+/L}$ and homozygous C/EBP $\beta^{\Delta\text{uORF}}$ and respective wild-type (WT) littermate control mice ($n = 6-7/\text{group}$) were generated as previously described [31,32] and fed standard chow diet. All animal experiments were conducted in strict accord with accepted standards of humane animal care under the auspices of the Animal (Scientific Procedures) Act UK 1986 and following prior approval by the Home Office in UK or following prior approval by the Institutional Animal Care and Use Committee in Berlin, Germany.

3T3-L1 Adipocyte Differentiation and Transfection with siRNA

3T3-L1 murine preadipocytes (ATCC, American Type Culture Collection) were maintained and differentiated into mature adipocytes as previously described [33]. Briefly, 2d after reaching confluence, 3T3-L1 cells were induced to differentiate by the addition of 0.5 μM dexamethasone, 500 μM 3-isobutyl-1-methylxanthine and 5 $\mu\text{g}/\text{ml}$ insulin for 2d. Thereafter, 3T3-L1 cell differentiation continued in medium supplemented with 5 $\mu\text{g}/\text{ml}$ insulin alone. Experiments were performed 8–12d following induction of differentiation, unless otherwise stated. 3T3-L1 adipocytes were transfected with siRNA using DeliverX Plus (Panomics Inc., Fremont, CA) according to the manufacturer's protocol. Fully differentiated adipocytes were gently detached by a brief wash with trypsin followed by collagenase type I (0.5 mg/ml; Invitrogen, Paisley, UK), and reseeded at 1.5×10^5 cells per well in a 12-well plate then transfected the next day with siRNA (32 pmol) in serum-free medium for 3 h followed by 21 h incubation in growth medium. All siRNAs were purchased from Applied Biosystems (Warrington, UK) and were scrambled (AM4611, negative control), C/EBP α (ID 101889), C/EBP β (ID 288793) and CHOP (ID 288792). In the experiments designed to induce ER stress, 3T3-L1 adipocytes were treated with tunicamycin (4 $\mu\text{g}/\text{ml}$). Rapamycin (100 nM and 500 nM, 24 h) was used to inhibit the mammalian target of rapamycin (mTOR) in 3T3-L1 adipocytes.

RNA Extraction and Analysis

Following dissection, subcutaneous adipose tissue (inguinal depot) was frozen on dry ice and stored at -70°C . Tissues were weighed on a balance model AA-160 (Denver Instrument Company, NY). RNA was extracted using Qiagen RNeasy Lipid Tissue Mini kit (Qiagen, West Sussex, UK). 3T3-L1 cells were harvested in Trizol. RNA (2 μg) was reverse transcribed using SuperScript III (Invitrogen) and quantified by real-time PCR on a LightCycler 480 (Roche) as described [16]. Primer-probe sets were purchased from Applied Biosystems: 11 β -HSD1 (Mm00476182_m1), C/EBP α

(Mm00514283_s1), C/EBP β (Mm00843434_s1), C/EBP δ (Mm00786711_s1) and CHOP (Mm00492097_m1). Routinely, at least two internal standards were used from among 18S (Hs99999901_s1), β -actin (Mm00607939_s1) and TATA binding protein (TBP; Mm00446973_m1).

Western Blot Analysis

Subcutaneous adipose tissue (inguinal depot) was homogenized in lysis buffer (0.125M Tris-HCL, pH 6.8, 2% sodium dodecyl sulphate, and 10% glycerol) in the presence of a protease inhibitor cocktail (P2714; Sigma-Aldrich, Dorset, UK) and heated at 100°C . 3T3-L1 adipocytes were harvested directly into lysis buffer and processed similarly. Electrophoresis was carried out on 4–12% NuPage Bis-Tris gels (Invitrogen). After transfer, blots were probed with antibodies to C/EBP α , β , δ (1:500 dilution of stock 200 $\mu\text{g}/\text{ml}$; Santa Cruz Biotechnology Inc., Santa Cruz, CA), CHOP (1:1000 dilution; Abcam, Cambridge, UK) and β -tubulin (1:10000 dilution; Chemicon/Millipore, Watford, UK) and secondary antibodies anti-rabbit IgG-HRP (1:2000 dilution from stock 400 $\mu\text{g}/\text{ml}$ Santa Cruz Biotechnology Inc.), anti-mouse IgG-HRP (1:4000 dilution; Cell signaling, Danvers, MA), anti-rabbit IgG Alexa Fluor 700 (Invitrogen) and anti-mouse IgG (Rockland Immunochemicals Inc. Gilbertsville, PA). The resulting bands were analyzed by ImageJ (NIH, USA) or the Odyssey Infrared Imaging System (LI-COR Biosciences Ltd, Cambridge, UK), as appropriate.

Chromatin Immunoprecipitation (ChIP) Assays

ChIP assays were carried out using an Upstate EZ ChIP kit (Millipore, Billerica, MA) according to the manufacturer's protocol. Briefly, differentiating 3T3-L1 cells (8 h after induction of differentiation) or fully mature adipocytes (differentiated for 9d) in 15 cm tissue culture dishes were incubated with dimethyl adipimidate.2 HCL (10 mM in PBS; Thermo Fisher Scientific Cramlington, UK) for 30 min in the dark followed by 1% (v/v) formaldehyde. Cells were lysed in sodium dodecyl sulphate buffer (provided in the kit). Chromatin was sheared by sonication using a Soniprep 150 (MSE; Beckenham, Kent, UK) with eight 10 s pulses, at maximum amplitude, keeping cells on ice between pulses. Immunoprecipitations were carried out with 5 μg C/EBP or control rabbit IgG antibody (Santa Cruz). After reversal of cross-links, purified DNA was amplified by standard PCR (35 cycles of 94°C 30 s, 56°C 1 min, 72°C 1 min) or real-time PCR (95°C 5 s, 65°C 1 min) as above, using primers spanning the following C/EBP-binding sites in the 11 β -HSD1 promoter [12]: footprint FP1 and FP2 (5'-CTAGTGCTGCCTGAGACTACTCC-3' and 5'-TTTTTCATAACTGCCATCAAACAG-3'), FP3 and FP4 (5'-CAGTAGGAGATGCTCAGGAACC-3' and 5'-AGAGCAACGATTGGCTTTTGG-3'), FP9 (5'-ACTTGCTGAGAGTCTTGAAGT-3' and 5'-AGCAAAA-TAAGCTCAAGGGAAAA-3') and a C/EBP β binding site in the C/EBP α promoter (5'-GGTGACTTAGAGGCTTAAAGGAG-3' and 5'-GTGACTTTCCAAGCGGTGAGT-3'). Primers spanning a 11 β -HSD1 intronic sequence not predicted to bind C/EBP were used as a control (5'-TCAAACATC-CAGGTTCTTTTCATT-3' and 5'-CATACCACATAAACCA-CACATGG-3'). Data are expressed relative to levels of input DNA in immunoprecipitations.

Statistical Analysis

All data were analyzed by Student's *t* test or ANOVA followed by *post hoc* Tukey, Fisher LSD or Dunnett tests using SigmaStat 2.03 statistical software. Significance was set at $p \leq 0.05$.




Figure 1. Expression of 11 β -HSD1, C/EBP α , β , δ and CHOP in adipose tissue of HF diet-fed mice compared to control diet-fed mice. (A) Real-time PCR measurements of levels of mRNA encoding 11 β -HSD1, C/EBP α , β , δ , and CHOP in adipose tissue of HF-fed (black bars) and control diet-fed (white bars) mice. Levels of mRNA, normalized to TBP, are expressed relative to levels in control diet-fed mice, arbitrarily set to 100%. (B-E) Representative western blots (50 μ g protein/lane) and quantification of blots probed with; (B) C/EBP α antibody showing levels of the 42 kDa (p42) and 30 kDa (p30) isoforms of C/EBP α as well as total C/EBP α (p42+p30) levels (the intermediate immunoreactive protein indicated by* is likely to be a proteolytic product of p42); (C) C/EBP β antibody showing levels of LAP (38 kDa LAP* +35 kDa LAP) and the 20 kDa LIP isoforms of C/EBP β as well as total C/EBP β (LAP+LIP) levels (inset shows LIP:LAP ratio); (D) C/EBP δ antibody and (E) CHOP antibody showing levels of respective proteins in adipose tissue of mice fed HF diet (HF, black bars) or control diet (C, white bars). Blots were stripped and reprobed with β -tubulin antibody, as loading control. In C/EBP β and CHOP westerns, all samples were analysed in the same gel but not all in adjacent lanes. C/EBP levels were quantified relative to β -tubulin levels using ImageJ analysis and are expressed in arbitrary units (AU). All values are mean \pm SEM; n = 16/group (A) or 4/group (B-E). *, p \leq 0.05; **, p $<$ 0.001. doi:10.1371/journal.pone.0037953.g001

Results

Effect of HF Diet on 11 β -HSD1 and C/EBP Expression in Mouse Adipose Tissue

Mice fed HF diet for 6 weeks were heavier (HF, 30.8 \pm 0.52 *vs* control diet, 26.3 \pm 0.34 g; p $<$ 0.01) with increased subcutaneous

adipose tissue weight (HF, 0.022 \pm 0.001 *vs* control, 0.012 \pm 0.001 (w/w) corrected for body weight; p $<$ 0.01), while liver weight was unchanged (HF, 0.049 \pm 0.001 *vs* control, 0.046 \pm 0.003 (w/w) corrected for body weight). 11 β -HSD1 mRNA was down-regulated in adipose tissue by HF diet both in subcutaneous (Fig. 1A) and visceral (mesenteric) depots (data not shown), but

Fig. 2




Figure 2. Transfection of fully differentiated 3T3-L1 adipocytes with C/EBP α or C/EBP β siRNA decreases 11 β -HSD1 mRNA levels, whilst CHOP siRNA has no effect. (A) Real-time PCR measurement of levels of mRNA encoding C/EBP α (left panel), C/EBP β (centre panel) and CHOP (right panel), 24 h after transfection of fully differentiated 3T3-L1 adipocytes with scrambled RNA (Scr; as control) or siRNAs (32 pmol) targeting C/EBP α , β or CHOP. Data are normalized to TBP and expressed relative to levels in cells transfected with scrambled RNA (arbitrarily set to 100%). Values are mean \pm SEM of 3 different experiments (independent adipocyte differentiations) with each siRNA treatment tested in triplicate. *, Significantly different from scrambled RNA, $p \leq 0.05$. (B) Representative western blots showing levels of the 42 kDa (p42) and 30 kDa (p30) isoforms of C/EBP α (left panel; 20 μ g protein/lane), 38 kDa LAP*, 35 kDa LAP and 20 kDa LIP isoforms of C/EBP β (centre panel; 40 μ g protein/lane) and CHOP (right panel; 20 μ g protein/lane), 24 h after transfection of 3T3-L1 adipocytes with scrambled RNA or respective siRNAs targeting C/EBP α , β or CHOP. Blots were stripped and reprobed with β -tubulin antibody, as loading control. In C/EBP β and CHOP westerns all samples were analysed in the same gel but not all in adjacent lanes. (C) Real-time PCR measurement of mRNA encoding 11 β -HSD1, 24 h after transfection of 3T3-L1 mature adipocytes with scrambled RNA or siRNA (32 pmol) targeting C/EBP α , β or CHOP. Data are normalized to TBP and expressed relative to levels in cells transfected with scrambled RNA (arbitrarily set to 100%). Values are mean \pm SEM of 3 to 4 different experiments (independent adipocyte differentiations) with each siRNA treatment tested in triplicate. *, Significantly different from scrambled RNA, $p \leq 0.05$. doi:10.1371/journal.pone.0037953.g002

was unchanged in liver (data not shown), consistent with previous data showing down-regulation of 11 β -HSD1 mRNA and enzyme activity in adipose tissue of mice fed HF diet [9,11]. To test whether altered C/EBP expression may underlie the dietary regulation of 11 β -HSD1, we examined C/EBP α , β , δ and CHOP expression in adipose tissue. HF diet did not change C/EBP α , β or δ mRNA levels, while CHOP mRNA

levels were increased in subcutaneous adipose tissue (Fig. 1A) and mesenteric [11] adipose tissue (data not shown). Because 11 β -HSD1 and C/EBP mRNA levels showed the same pattern of changes in both adipose depots of HF-fed (vs control) animals, subcutaneous adipose, which is more abundant, was used for subsequent analyses. Consistent with mRNA levels, western blot analysis showed an increase in CHOP protein




Figure 4. Altered C/EBP β -LIP:LAP ratio *in vivo* affects 11 β -HSD1 expression in adipose tissue. (A) Representative western blot (40 μ g protein/lane) showing levels of C/EBP β -LAP (38 kDa LAP* +35 kDa LAP) and -LIP (20 kDa) isoforms in the adipose tissue of C/EBP β mutants C/EBP $\beta^{+/L}$ (+/L), C/EBP $\beta^{\Delta uORF}$ (Δ uORF) and control (Con) mice. Blots were stripped and reprobed with β -tubulin antibody, as loading control. All samples were analysed in the same gel but not all in adjacent lanes. (B) Real-time PCR measurement levels of mRNA encoding 11 β -HSD1 in adipose tissue of wild type control mice (Con, white bar), C/EBP $\beta^{+/L}$ (+/L, black bar) and C/EBP $\beta^{\Delta uORF}$ (Δ uORF, grey bar). C/EBP $\beta^{+/L}$ mice are heterozygous for an allele of C/EBP β in which the normal gene has been replaced by C/EBP β -LIP (a “knock-in”) [31] and C/EBP $\beta^{\Delta uORF}$ is homozygous for the deletion of upstream ORF codon [32]. Adipose 11 β -HSD1 mRNA levels, normalized to TBP, are expressed relative to levels in control mice (arbitrarily set to 100%) and are mean \pm SEM; n=6–9/group. *, p \leq 0.05; **, p<0.001. doi:10.1371/journal.pone.0037953.g004

LIP:LAP ratio (scrambled RNA, 0.5 ± 0.03 vs C/EBP β siRNA 0.7 ± 0.02). The results also showed that reduced expression of one of the C/EBPs in mature 3T3-L1 adipocytes did not affect mRNA levels of the other C/EBPs tested (Fig. 2A). Both C/EBP α and C/EBP β siRNA reduced 11 β -HSD1 mRNA levels in 3T3-L1 adipocytes (Fig. 2C), demonstrating that both these members of the C/EBP family play a key role in maintaining adipocyte expression of 11 β -HSD1. In contrast, although levels of CHOP were reduced in 3T3-L1 adipocytes by siRNA (Fig. 2A, B), this did not alter 11 β -HSD1 mRNA levels (Fig. 2C).

Fig. 4

C/EBP β Locates to the Promoter of 11 β -HSD1 throughout Adipocyte Differentiation Whilst C/EBP α Binds Only in Differentiated Adipocytes

Previous reports showed C/EBP β binding to the 11 β -HSD1 promoter in undifferentiated 3T3-L1 preadipocytes [13,14], but this has not been described in adipocytes. To determine if C/EBPs interact with the 11 β -HSD1 promoter in immature and fully mature adipocytes, binding of C/EBP α and β was assayed by chromatin immunoprecipitation (ChIP). The proximal promoter of the rat 11 β -HSD1 gene contains 4 C/EBP binding sites between -196 and +44; footprints (FP) 1 to 4, which are conserved in the mouse promoter [12,13] (Fig. 3A). C/EBP β was bound to the proximal 11 β -HSD1 promoter early during adipocyte differentiation (when 11 β -HSD1 expression is low) (Fig. 3B) as well as in fully differentiated 3T3-L1 adipocytes (Fig. 3C). In contrast, C/EBP α was only bound to the 11 β -HSD1 proximal promoter in fully differentiated 3T3-L1 adipocytes, which highly express 11 β -HSD1 [33]. Neither C/EBP α nor C/EBP β were detected at FP9 in the promoter of 11 β -HSD1 (also conserved between rat and mouse) (Fig. 3B, C). Thus C/EBP β interacts directly with the 11 β -HSD1 promoter in mature adipocytes.

Increased C/EBP β -LIP:LAP Ratio *in vivo* Decreases 11 β -HSD1 mRNA Levels in Adipose Tissue, an Effect Reversed in Mice with Decreased C/EBP β -LIP:LAP Ratio

To test whether altered C/EBP β -LIP:LAP ratio can similarly alter adipose 11 β -HSD1 expression *in vivo*, we measured 11 β -HSD1 mRNA levels in subcutaneous adipose tissue of C/EBP $\beta^{+/L}$ and C/EBP $\beta^{\Delta uORF}$ mice which have increased and decreased C/EBP β -LIP:LAP ratio, respectively. In C/EBP $\beta^{+/L}$ mice, a “knock-in” of C/EBP β -LIP (L allele) replaces the normal C/EBP β gene [31], increasing C/EBP β -LIP:LAP ratio (Fig. 4A). In C/EBP $\beta^{\Delta uORF}$ mice, deletion of the upstream open reading frame (Δ uORF allele) prevents translation of C/EBP β -LIP [32], thus decreasing C/EBP β -LIP:LAP ratio (Fig. 4A). 11 β -HSD1 mRNA levels were significantly lower in adipose tissue of C/EBP $\beta^{+/L}$ mice (increased C/EBP β -LIP:LAP ratio) than in wild-type controls (Fig. 4B). The reverse was observed in C/EBP $\beta^{\Delta uORF}$ mice (Fig. 4B), demonstrating that, *in vivo*, the C/EBP β -LIP:LAP ratio controls the transcription of 11 β -HSD1.

11 β -HSD1 mRNA Levels are Down-regulated by ER Stress in 3T3-L1 Adipocytes

Elevated C/EBP β -LIP:LAP ratio and CHOP levels, observed in adipose tissue of HF-fed mice, may be a result of ER stress [34,35]. Treatment of differentiated 3T3-L1 adipocytes with tunicamycin, an inducer of ER stress [35], significantly reduced 11 β -HSD1 mRNA levels at 6 h, with a more pronounced effect at 16 h (Fig. 5A). Similar results were obtained with an alternative ER stress-inducer, thapsigargin (data not shown). Down-regulation of 11 β -HSD1 mRNA was accompanied by up-regulation of genes increased by ER stress; CHOP and the 78 kDa glucose-regulated protein (GRP78) (Fig. 5C, E), though calreticulin mRNA levels were unaffected at this stage (Fig. 5D). C/EBP β mRNA levels were not altered (Fig. 5B) whilst, as expected, C/EBP β -LIP:LAP ratio was increased, and both C/EBP β -LAP and -LIP were up-regulated (Fig. 5F). These results show that induction of ER stress, which increases the C/EBP β -LIP:LAP ratio, also down-regulates 11 β -HSD1 in 3T3-L1 adipocytes.

mTOR Regulates the Expression of 11 β -HSD1 in 3T3-L1 Adipocytes

The mTOR pathway, which is responsive to nutrient availability and growth factors, controls many aspects of cellular




Figure 5. Tunicamycin-induced ER stress reduces 11 β -HSD1 expression in 3T3-L1 adipocytes. (A–E) Real-time PCR measurement of levels of mRNA encoding (A) 11 β -HSD1, (B) C/EBP β , (C) CHOP, (D) calreticulin and (E) GRP78 in untreated differentiated 3T3-L1 adipocytes (white bars) or treated with tunicamycin (4 μ g/ml, black bars) for 6 or 16 h. Data are normalized to TBP and expressed relative to levels of control cells (arbitrarily set to 100%). (F) Representative western blot (40 μ g protein/lane) and quantification showing levels of C/EBP β -LAP (38 kDa LAP* +35 kDa LAP) and -LIP (20 kDa) isoforms. Inset shows C/EBP β -LIP:LAP ratio in untreated 3T3-L1 adipocytes (C, white bar) or treated with tunicamycin for 0.5 h (0.5 h, grey bar) or 2 h (2 h, black bar). Blots were stripped and reprobed with β -tubulin antibody, as loading control. All samples were analysed in the same gel but not in adjacent lanes. C/EBP β levels were quantified relative to β -tubulin and are expressed in arbitrary units (AU). Values are mean \pm SEM of 2–3 independent adipocyte differentiations with each treatment tested in duplicate or triplicate. *, Significantly different from control. *, $p \leq 0.05$; **, $p < 0.001$; ***, $p < 0.0001$.

doi:10.1371/journal.pone.0037953.g005




Figure 6. Inhibition of mTOR in 3T3-L1 adipocytes reduces C/EBP β -LIP:LAP ratio and increases 11 β -HSD1 expression. Real-time PCR measurement of levels of mRNA encoding (A) C/EBP β and (B) 11 β -HSD1 in 3T3-L1 adipocytes (white bars) or following treatment with 100 nM (grey bars) or 500 nM (black bars) rapamycin for 24 h. Data are normalized to TBP and expressed relative to levels of control cells (arbitrarily set to 100%). (C) Representative western blot (40 μ g protein/lane) and quantification showing levels of C/EBP β -LAP (38 kDa LAP* +35 kDa LAP) and -LIP (20 kDa) isoforms and inset showing C/EBP β -LIP:LAP ratio in untreated 3T3-L1 adipocytes (C, white bars) or following treatment with 100 nM (100, grey bars) or 500 nM (500, black bars) rapamycin for 24 h. Blots were stripped and reprobed with β -tubulin antibody, as loading control. All samples were analysed in the same gel but not in adjacent lanes. C/EBP β levels were quantified relative to β -tubulin and are expressed in arbitrary units (AU). Values are mean \pm SEM of up to 3 independent adipocyte differentiations, each treatment tested in triplicate. *, Significantly different from control. *, $p \leq 0.05$; **, $p < 0.001$.

doi:10.1371/journal.pone.0037953.g006

metabolism and regulates the production of C/EBP β -LIP and -LAP isoforms [24]. To test the importance of this pathway for the transcriptional regulation of 11 β -HSD1, rapamycin was used to inhibit mTOR in differentiated 3T3-L1 adipocytes. Rapamycin did not alter C/EBP β mRNA levels (Fig. 6A) or C/EBP β -LAP, but decreased C/EBP β -LIP, resulting in a lower C/EBP β -LIP:LAP ratio (Fig. 6C). This was associated with increased 11 β -HSD1 mRNA levels (Fig. 6B). These results show that the mTOR signalling pathway regulates 11 β -HSD1 transcription in adipocytes, plausibly via modulation of the C/EBP β -LIP:LAP ratio.

Discussion

This study emphasises the key role played by members of the C/EBP family in the regulation of 11 β -HSD1 transcription in adipocytes. However, the main finding here is that the striking down-regulation of 11 β -HSD1 in adipose tissue of mice in response to HF feeding is likely due to changes in post-transcriptional processing of C/EBP β generating an increased

C/EBP β -LIP:LAP ratio. Indeed, knock-down of C/EBP β in fully differentiated mature 3T3-L1 adipocytes showed that C/EBP β is required for normal 11 β -HSD1 expression in differentiated adipocytes, consistent with a previous report of reduced 11 β -HSD1 expression in adipose tissue of C/EBP β -deficient mice [29]. In differentiating 3T3-L1 adipocytes, C/EBP β binds to the C/EBP α promoter and induces its expression [20]. Importantly, in contrast, knock-down of C/EBP β in fully differentiated adipocytes did not affect levels of C/EBP α , suggesting a direct role for C/EBP β in adipocyte 11 β -HSD1 expression, independent of its effect on C/EBP α expression. Indeed, C/EBP α mRNA and protein levels were unchanged in adipose tissue of HF diet-fed mice. However, it cannot be excluded that C/EBP β may also exert its effect by regulating C/EBP α transactivation potential. Nevertheless, supporting a direct role, C/EBP β was bound to the 11 β -HSD1 promoter in mature adipocytes as well as in 3T3-L1 preadipocytes during the differentiation process. C/EBP α , in contrast, was only bound to the 11 β -HSD1 promoter in

differentiated adipocytes, as expected from its later appearance during differentiation [20]. Thus, both C/EBP α and C/EBP β play a direct and positive role in regulation of 11 β -HSD1 transcription in differentiated adipocytes, but C/EBP β alone is crucial in differentiating cells. This is likely to impact on the 11 β -HSD1 mRNA levels in these cells, where C/EBP α is absent or in low levels. In adipose tissue, C/EBP β and CHOP were the only C/EBPs changed by HF diet. Although CHOP generally represses transcription [36], it can also be a co-activator [37]. However, siRNA-mediated knock-down of CHOP showed that it plays no role in 11 β -HSD1 expression in fully differentiated 3T3-L1 adipocytes, although we cannot rule out that increased CHOP levels may contribute to the down-regulation of 11 β -HSD1 in adipose tissue with HF diet.

An altered C/EBP β -LIP:LAP ratio, like the HF-induced increase in CHOP, might be a response to ER stress as a consequence of a high demand for ER activity with nutrient excess [34]. The C/EBP β -LIP:LAP ratio is controlled by the double stranded RNA-dependent protein kinase (PKR, a target of ER stress [38]) and mTOR signalling pathways through the eukaryotic translation factors eIF-2 α and eIF-4E, respectively [24]. Indeed, diet-induced obesity in mice results in the activation of ER stress pathways in metabolically relevant tissues, such as adipose and liver, with increased phosphorylation of the PKR-like kinase and its substrate eIF-2 α , which are key indicators of ER stress [34]. mTOR may contribute to the ER stress response as well [39]. Our results show that both induction of ER stress and inhibition of mTOR signalling alter C/EBP β -LIP:LAP ratio and modulate 11 β -HSD1 mRNA levels in 3T3-L1 adipocytes. ER stress and mTOR signalling pathways are sensitive to cellular nutrients and energy homeostasis, suggesting that the increased C/EBP β -LIP:LAP ratio in obese adipose tissue may result from activation of these nutrient/stress sensing mechanisms leading to the down-regulation of 11 β -HSD1.

References

- Seckl JR, Morton NM, Chapman KE, Walker BR (2004) Glucocorticoids and 11 β -hydroxysteroid dehydrogenase in adipose tissue. *Recent Prog Horm Res* 59: 359–393.
- Masuzaki H, Paterson J, Shinyama H, Morton NM, Mullins JJ, et al. (2001) A transgenic model of visceral obesity and the metabolic syndrome. *Science* 294: 2166–2170.
- Lindsay RS, Wake DJ, Nair S, Bunt J, Livingstone DE, et al. (2003) Subcutaneous adipose 11 β -hydroxysteroid dehydrogenase type 1 activity and messenger ribonucleic acid levels are associated with adiposity and insulinemia in Pima Indians and Caucasians. *J Clin Endocrinol Metab* 88: 2738–2744.
- Wake DJ, Rask E, Livingstone DE, Soderberg S, Olsson T, et al. (2003) Local and systemic impact of transcriptional up-regulation of 11 β -hydroxysteroid dehydrogenase type 1 in adipose tissue in human obesity. *J Clin Endocrinol Metab* 88: 3983–3988.
- Masuzaki H, Yamamoto H, Kenyon CJ, Elmquist JK, Morton NM, et al. (2003) Transgenic amplification of glucocorticoid action in adipose tissue causes high blood pressure in mice. *J Clin Invest* 112: 83–90.
- Morton NM, Paterson JM, Masuzaki H, Holmes MC, Staels B, et al. (2004) Novel adipose tissue-mediated resistance to diet-induced visceral obesity in 11 β -hydroxysteroid dehydrogenase type 1-deficient mice. *Diabetes* 53: 931–938.
- Hermanowski-Vosatka A, Balkovec JM, Cheng K, Chen HY, Hernandez M, et al. (2005) 11 β -HSD1 inhibition ameliorates metabolic syndrome and prevents progression of atherosclerosis in mice. *J Exp Med* 202: 517–527.
- Rosenstock J, Banarer S, Fonseca V, Inzucchi S, Hollis GF, et al. (2010) The 11 β -hydroxysteroid dehydrogenase type 1 inhibitor INCB13739 improves hyperglycemia in patients with type 2 diabetes inadequately controlled by metformin monotherapy. *Diabetes Care* 33: 1516–1522.
- Morton NM, Ramage L, Seckl JR (2004) Down-regulation of adipose 11 β -hydroxysteroid dehydrogenase type 1 by high-fat feeding in mice: a potential adaptive mechanism counteracting metabolic disease. *Endocrinology* 145: 2707–2712.
- Drake AJ, Livingstone DE, Andrew R, Seckl JR, Morton NM, et al. (2005) Reduced adipose glucocorticoid reactivation and increased hepatic glucocorticoid clearance as an early adaptation to high-fat feeding in Wistar rats. *Endocrinology* 146: 913–919.
- Man TY, Michailidou Z, Gokcel A, Ramage L, Chapman KE, et al. (2011) Dietary manipulation reveals an unexpected inverse relationship between fat mass and adipose 11 β -hydroxysteroid dehydrogenase type 1. *Am J Physiol Endocrinol Metab* 300: E1076–1084.
- Williams IJ, Lyons V, MacLeod I, Rajan V, Darlington GJ, et al. (2000) C/EBP regulates hepatic transcription of 11 β -hydroxysteroid dehydrogenase type 1. A novel mechanism for cross-talk between the C/EBP and glucocorticoid signaling pathways. *J Biol Chem* 275: 30232–30239.
- Gout J, Tirard J, Thevenon C, Riou JP, Begout M, et al. (2006) CCAAT/enhancer-binding proteins (C/EBPs) regulate the basal and cAMP-induced transcription of the human 11 β -hydroxysteroid dehydrogenase encoding gene in adipose cells. *Biochimie* 88: 1115–1124.
- Arai N, Masuzaki H, Tanaka T, Ishii T, Yasue S, et al. (2007) Ceramide and adenosine 5'-monophosphate-activated protein kinase are two novel regulators of 11 β -hydroxysteroid dehydrogenase type 1 expression and activity in cultured preadipocytes. *Endocrinology* 148: 5268–5277.
- Yang Z, Guo C, Zhu P, Li W, Myatt L, et al. (2007) Role of glucocorticoid receptor and CCAAT/enhancer-binding protein α in the feed-forward induction of 11 β -hydroxysteroid dehydrogenase type 1 expression by cortisol in human amnion fibroblasts. *J Endocrinol* 195: 241–253.
- Sai S, Esteves CL, Kelly V, Michailidou Z, Anderson K, et al. (2008) Glucocorticoid regulation of the promoter of 11 β -hydroxysteroid dehydrogenase type 1 is indirect and requires CCAAT/enhancer-binding protein- β . *Mol Endocrinol* 22: 2049–2060.
- Cao Z, Umek RM, McKnight SL (1991) Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes Dev* 5: 1538–1552.
- Wang ND, Finegold MJ, Bradley A, Ou CN, Abdelsayed SV, et al. (1995) Impaired energy homeostasis in C/EBP α knockout mice. *Science* 269: 1108–1112.
- Tanaka T, Yoshida N, Kishimoto T, Akira S (1997) Defective adipocyte differentiation in mice lacking the C/EBP β and/or C/EBP δ gene. *EMBO J* 16: 7432–7443.
- Otto TC, Lane MD (2005) Adipose development: from stem cell to adipocyte. *Crit Rev Biochem Mol Biol* 40: 229–242.

Acknowledgments

We thank members of the Endocrinology Unit for helpful discussions. We also thank BRR technical staff for expert animal care.

Author Contributions

Conceived and designed the experiments: CLE KEC. Performed the experiments: CLE VK VB TYM. Analyzed the data: CLE JRS KEC. Contributed reagents/materials/analysis tools: NMM AL. Wrote the paper: CLE JRS KEC.

21. Ariyama Y, Shimizu H, Satoh T, Tsuchiya T, Okada S, et al. (2007) Chop-deficient mice showed increased adiposity but no glucose intolerance. *Obesity* 15: 1647–1656.
22. Ossipov V, Descombes P, Schibler U (1993) CCAAT/enhancer-binding protein mRNA is translated into multiple proteins with different transcription activation potentials. *Proc Natl Acad Sci U S A* 90: 8219–8223.
23. Welm AL, Timchenko NA, Darlington GJ (1999) C/EBP α regulates generation of C/EBP β isoforms through activation of specific proteolytic cleavage. *Mol Cell Biol* 19: 1695–1704.
24. Calkhoven CF, Muller C, Leutz A (2000) Translational control of C/EBP α and C/EBP β isoform expression. *Genes Dev* 14: 1920–1932.
25. Lin FT, MacDougald OA, Dichl AM, Lane MD (1993) A 30-kDa alternative translation product of the CCAAT/enhancer binding protein α message: transcriptional activator lacking antimitotic activity. *Proc Natl Acad Sci U S A* 90: 9606–9610.
26. Duong DT, Waltner-Law ME, Sears R, Sealy L, Granner DK (2002) Insulin inhibits hepatocellular glucose production by utilizing liver-enriched transcriptional inhibitory protein to disrupt the association of CREB-binding protein and RNA polymerase II with the phosphoenolpyruvate carboxykinase gene promoter. *J Biol Chem* 277: 32234–32242.
27. Masaki T, Matsuura T, Ohkawa K, Miyamura T, Okazaki I, et al. (2006) All-trans retinoic acid down-regulates human albumin gene expression through the induction of C/EBP β -LIP. *Biochem J* 397: 345–353.
28. Siu F, Chen C, Zhong C, Kilberg MS (2001) CCAAT/enhancer-binding protein- β is a mediator of the nutrient-sensing response pathway that activates the human asparagine synthetase gene. *J Biol Chem* 276: 48100–48107.
29. Payne VA, Au WS, Gray SL, Nora ED, Rahman SM, et al. (2007) Sequential regulation of diacylglycerol acyltransferase 2 expression by CAAT/enhancer-binding protein β (C/EBP β) and C/EBP α during adipogenesis. *J Biol Chem* 282: 21005–21014.
30. Yang Z, Zhu X, Guo C, Sun K (2009) Stimulation of 11 β -HSD1 expression by IL-1 β via a C/EBP binding site in human fetal lung fibroblasts. *Endocrine* 36: 404–411.
31. Smink JJ, Begay V, Schoenmaker T, Sterneck E, de Vries TJ, et al. (2009) Transcription factor C/EBP β isoform ratio regulates osteoclastogenesis through MafB. *EMBO J* 28: 1769–1781.
32. Wethmar K, Begay V, Smink JJ, Zaragoza K, Wiesenthal V, et al. C/EBP β Δ uORF mice—a genetic model for uORF-mediated translational control in mammals. *Genes Dev* 24: 15–20.
33. Napolitano A, Voice MW, Edwards CR, Seckl JR, Chapman KE (1998) 11 β -hydroxysteroid dehydrogenase 1 in adipocytes: expression is differentiation-dependent and hormonally regulated. *J Steroid Biochem Mol Biol* 64: 251–260.
34. Ozcan U, Cao Q, Yilmaz E, Lee AH, Iwakoshi NN, et al. (2004) Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. *Science* 306: 457–461.
35. Li Y, Bevilacqua E, Chiribau CB, Majumder M, Wang C, et al. (2008) Differential control of the CCAAT/enhancer-binding protein β (C/EBP β) products liver-enriched transcriptional activating protein (LAP) and liver-enriched transcriptional inhibitory protein (LIP) and the regulation of gene expression during the response to endoplasmic reticulum stress. *J Biol Chem* 283: 22443–22456.
36. Ron D, Habener JF (1992) CHOP, a novel developmentally regulated nuclear protein that dimerizes with transcription factors C/EBP and LAP and functions as a dominant-negative inhibitor of gene transcription. *Genes Dev* 6: 439–453.
37. Cucinotta M, Visalli M, Aguenouz M, Valenti A, Loddo S, et al. (2008) Regulation of Interleukin-8 Gene at a Distinct Site of Its Promoter by CCAAT Enhancer-binding Protein Homologous Protein in Prostaglandin E2-treated Human T Cells. *J Biol Chem* 283: 29760–29769.
38. Lee ES, Yoon CH, Kim YS, Bae YS (2007) The double-strand RNA-dependent protein kinase PKR plays a significant role in a sustained ER stress-induced apoptosis. *FEBS Lett* 581: 4325–4332.
39. Gregor MF, Hotamisligil GS (2007) Thematic review series: Adipocyte Biology. Adipocyte stress: the endoplasmic reticulum and metabolic disease. *J Lipid Res* 48: 1905–1914.
40. Arizmendi C, Liu S, Croniger C, Poli V, Friedman JE (1999) The transcription factor CCAAT/enhancer-binding protein β regulates gluconeogenesis and phosphoenolpyruvate carboxykinase (GTP) gene transcription during diabetes. *J Biol Chem* 274: 13033–13040.
41. Liu S, Croniger C, Arizmendi C, Harada-Shiba M, Ren J, et al. (1999) Hypoglycemia and impaired hepatic glucose production in mice with a deletion of the C/EBP β gene. *J Clin Invest* 103: 207–213.
42. Millward CA, Heaney JD, Sinasac DS, Chu EC, Bederman IR, et al. (2007) Mice with a deletion in the gene for CCAAT/enhancer-binding protein β are protected against diet-induced obesity. *Diabetes* 56: 161–167.
43. Rahman SM, Schroeder-Gloekler JM, Janssen RC, Jiang H, Qadri I, et al. (2007) CCAAT/enhancing binding protein β deletion in mice attenuates inflammation, endoplasmic reticulum stress, and lipid accumulation in diet-induced nonalcoholic steatohepatitis. *Hepatology* 45: 1108–1117.
44. Christy RJ, Kaestner KH, Geiman DE, Lane MD (1991) CCAAT/enhancer binding protein gene promoter: binding of nuclear factors during differentiation of 3T3-L1 preadipocytes. *Proc Natl Acad Sci U S A* 88: 2593–2597.