

Nuclear factor-kappa B1 inhibits early apoptosis of glioma cells by promoting the expression of Bcl-2

Tian-quan Yang^{1,*}
 Min Chen^{1,*}
 Yong-qiang Wang¹
 Wei Xu¹
 Yong Han¹
 Jin Xu¹
 Yong-jun Xiang¹
 Bin Yuan¹
 Hang-zhou Wang¹
 You-xin Zhou²

¹Department of Neurosurgery, Children's Hospital of Soochow University, ²Department of Neurosurgery and Brain and Nerve Research Laboratory, The First Affiliated Hospital of Soochow University, Suzhou, People's Republic of China

*These authors contributed equally to this work

Abstract: Glioma is one of the most common types of adult primary brain tumors, and the underlying molecular mechanisms still remain unclear. Nuclear factor-kappa B1 (NF-κB1) is involved in a variety of malignancies and is widely expressed in malignant tumors. However, the expression of NF-κB1 in different grades of glioma, the correlation between NF-κB1 and Bcl-2 expressions in gliomas, and the research between NF-κB1 and early apoptosis of glioma cells have not been reported so far. In this study, the expression level of NF-κB1 in 31 human glioma tissues and six nonneoplastic brain tissues was determined using quantitative real-time polymerase chain reaction. Results showed that the expression of NF-κB1 in human glioma tissues and glioma cell lines, SHG44 and U87, was significantly higher compared to noncancerous brain tissues and that the expression increased with increasing degrees of tumor malignancy. Similar results were demonstrated with the expression of Bcl-2 in the same human glioma specimens. Flow cytometry results showed that inhibition of NF-κB1 expression significantly promoted apoptosis of SHG44 and U87 in human glioma cells. Western blot analysis further confirmed decreased expression of Bcl-2 protein after inhibition of NF-κB1 protein expression. Taken together, NF-κB1 overexpression inhibits early apoptosis of glioma cells and high expression of NF-κB1 promotes the expression of antiapoptotic gene *Bcl-2*. Therefore, our study results provide a theoretical basis for antiapoptotic mechanism of tumor cells in association with NF-κB1.

Keywords: NF-κB1, Bcl-2, glioma, apoptosis

Introduction

Glioma is one of the most common types of primary brain tumors in adults, accounting for nearly 40% of all central nervous system malignancies,^{1,2} and is one of the most aggressive and lethal human cancer types.³ They have the ability of higher proliferation, invasion, migration, and lower apoptosis. In spite of advances in surgery, radiation therapy, and chemotherapy, the prognosis of patients with gliomas has not significantly improved, and the median survival time of patients is expected to be 12–15 months.^{4,5} The unrestricted growth of glioma is not only the result of abnormal cell differentiation and proliferation but also closely related to the inhibition of apoptosis.⁶ During the occurrence and development process of glioma, abnormal expression or mutation of apoptosis-related genes not only block apoptosis but also promote the occurrence and development of tumor.⁷ Therefore, studying the underlying mechanism of glioma apoptosis is of great significance.

Nuclear factor-kappa B1 (NF-κB1) is a member of the Rel gene family and involved in the control of tumor cell growth, immune response, apoptosis, and transcriptional regulation.⁸ Studies have shown the association of NF-κB1 gene with thyroid cancer, oral squamous cell carcinoma, and colorectal cancer.^{9–11} However, there is little research on the association of NF-κB1 and gliomas. According to our

Correspondence: Hang-zhou Wang
 Department of Neurosurgery, Children's Hospital of Soochow University, 92 Zhongnan Street, Suzhou, Jiangsu 215000, People's Republic of China
 Tel +86 139 1408 3728
 Email wanghangzhoudr@126.com

You-xin Zhou
 Department of Neurosurgery and Brain and Nerve Research Laboratory, The First Affiliated Hospital of Soochow University, 188 Shizi Street, Suzhou, Jiangsu 215006, People's Republic of China
 Tel +86 139 1552 2828
 Email zhoyxyq2008@163.com

previous study, NF- κ B1 was highly expressed in gliomas.¹² However, the expression of NF- κ B1 in different grades of gliomas, the correlation between NF- κ B1 and Bcl-2 expressions in gliomas, and the research between NF- κ B1 and early apoptosis of glioma cells have not been reported so far. Hence, in this study, the expression of NF- κ B1 protein in U87 and SHG44 glioma cells was downregulated by transfecting the cells to NF- κ B1 shRNA plasmid vector and, then, observing the changes of early apoptosis of glioma cells. Also, the correlation between NF- κ B1 and Bcl-2 was explored by *in vitro* experiments, which indicated that NF- κ B1 inhibited the early apoptosis of glioma cells by promoting the expression of Bcl-2.

Materials and methods

Human tissue samples

For the study, 31 glioma samples were obtained from 31 Chinese patients between March 2000 and September 2016 from the Children's Hospital of Soochow University and The First Affiliated Hospital of Soochow University (Suzhou, China). Patients included 19 men and 12 women. There were eight cases of grade I glioma (pilocytic astrocytoma [PA]), nine cases of grade II (diffuse astrocytoma [DA]), seven cases of grade III (anaplastic astrocytoma [AA]), and seven cases of grade IV (primary brain glioblastoma [GBM]), according to the 2007 World Health Organization classification system. Six samples of nonneoplastic brain tissue samples were obtained from adult patients with craniocerebral injuries, for whom a partial resection of brain tissue was required as decompression treatments to reduce intracranial pressure. This research was permitted and approved by the ethics committees of the Children's Hospital of Soochow University and The First Affiliated Hospital of Soochow University. All human samples were used in accordance with the policies of the ethics committee. Informed written consents were obtained from the children's parents prior to participation in the study.

Cell source and culture

Human glioma cell lines U87 and SHG44 were purchased from the Cell Bank Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). Cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM; U87; Hyclone, Thermo Fisher Scientific, Waltham, MA, USA) or RPMI-1640 medium (SHG44; Hyclone, Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS) and 100 U/mL penicillin/streptomycin. All cultures were maintained in a humidified atmosphere of 5% CO₂ at 37°C.

RNA extraction and quantitative real-time PCR

Total RNA from tissues (100 mg) and cells (1×10^7 cells) were isolated using the TRIzol reagent (Thermo Fisher Scientific) for both mRNA and miRNA analyses. Relative levels of NF- κ B1 and Bcl-2 mRNA were examined using SYBR green quantitative real-time polymerase chain reaction (qRT-PCR) (LightCycler[®] 480; Hoffman-La Roche Ltd., Basel, Switzerland) and were normalized to levels of GAPDH mRNA. The following primers were used: NF- κ B1 forward, 5'-CCTCTCTCTAATCAGCCCTCTG-3'; NF- κ B1 reverse, 5'-GAGGACCTGGGAGTAGATGAG-3'; GAPDH forward, 5'-AGAAGGCTGGGGCTCATTTG-3'; GAPDH reverse, 5'-AGGGGCCATCCACAGTCTTC-3'. Each sample was performed in triplicate, and the program was set at 95°C 1 min, 95°C 10 s, 59°C 20 s, and 72°C 10 s, in Roche 480 quantitative PCR instrument for 45 cycles. The results were analyzed with $2^{-\Delta\Delta CT}$ values.

Plasmid transfection and experimental grouping

U87 and SHG44 cells in the logarithmic growth phase were inoculated into six-well plates 1 day before transfection. The cells were divided into blank control group (mock), no-load plasmid group (negative), NF- κ B1 silencing group (NF- κ B1 shRNA), NF- κ B1 overexpression group (NF- κ B1), Bcl-2 silencing group (Bcl-2 shRNA), and NF- κ B1 overexpression + Bcl-2 silencing group (NF- κ B1 + Bcl-2 shRNA), and three replicates in each group. When the cells reached a confluence of 40%–50%, Lipofectamine 2000 was used to mediate no-load plasmid, NF- κ B1 shRNA plasmid, NF- κ B1 overexpression plasmid, and Bcl-2 shRNA plasmid transfections. In the human glioma cell lines, U87 and SHG44, Lipofectamine 2000 was used to mediate NF- κ B1 overexpression plasmid and, then, the stable expression of the cell line was screened. The Bcl-2 shRNA plasmid was further transfected with the Lipofectamine 2000 reagent (Thermo Fisher Scientific) according to the manufacturer's instructions. Grouped cells were collected at 24, 48, and 72 h after transfection.

Analysis of apoptosis

Cells were allowed to attach and proliferate in six-well plates for 24 h before transfection with no-load plasmid and NF- κ B1 shRNA plasmid. Human glioma cell lines U87 and SHG44 were collected at 24, 48, and 72 h after transfection. Then, the proportion of cells in early-stage apoptosis was detected using the annexin V-PE Apoptosis Detection Kit (eBioscience, San Diego, CA, USA) and flow cytometry (FACSCanto; BD Biosciences, San Jose, CA, USA). Cells were washed once in phosphate

buffered solution ([PBS]; Gibco, Thermo Fisher Scientific USA) and once in 1 \times binding buffer. Cells were resuspended in 1 \times binding buffer at 1–5 $\times 10^6$ /mL. Fluorochrome-conjugated annexin V (5 μ L) was added to 100 μ L cell suspension. Cells were incubated for 10–15 min at room temperature. Cells were washed in 1 \times binding buffer and resuspended in 200 μ L of 1 \times binding buffer. The 7-AAD Viability Staining Solution (5 μ L) was added, and the cells were analyzed by flow cytometry. All analyses were carried out in triplicate.

Western blot analysis

After 72 h of transfection, the total protein of the abovementioned experimental group was extracted, quantified by Bradford method, and 20 μ L was added to each sample (protein amount 30 μ g). Protein samples were separated with 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred onto nitrocellulose membranes. Membranes were incubated with primary antibodies overnight at 4°C. Membranes were washed and incubated for 2 h with horseradish peroxidase-conjugated antirabbit secondary antibodies (ProSci, Poway, CA, USA), followed by detection and visualization using electrochemiluminescence Western blotting detection reagents (Pierce antibodies; Thermo Fisher Scientific). The primary antibodies used were anti-NF- κ B1 (Abcam, Tokyo, Japan) and anti-Bcl-2 (Abcam).

Statistical methods

The data were analyzed using the SPSS 17.0 statistical software (SPSS Inc., Chicago, IL, USA). Data were expressed as mean \pm standard deviation ($\chi \pm s$). Statistical significance of the two means was analyzed by single factor analysis of variance (ANOVA) and least square difference (LSD) *t*-test. $P < 0.05$ was considered statistically significant.

Results

NF- κ B1 and Bcl-2 were highly expressed in glioma, and the expression trend was consistent

To assess NF- κ B1 and Bcl-2 expressions in noncancerous brain tissues and in different grades of glioma, we detected mRNA expression levels of NF- κ B1 and Bcl-2 in six nonneoplastic brain tissues and in 31 human glioma tissue samples. qRT-PCR results revealed that the expression of NF- κ B1 and Bcl-2 in the tissue specimens of five groups was statistically significant ($P < 0.05$; Figure 1A and B). Data showed that the expression of NF- κ B1 in glioma was significantly higher than that in nonneoplastic brain tissues ($P < 0.01$; Figure 1A), and the expression of NF- κ B1 in I, II, III, and IV grades of glioma was gradually increased; the difference was statistically significant ($P < 0.05$; Figure 1A). At the same time, Bcl-2 expression

in the same brain glioma specimens was significantly higher than that of the nontumor brain tissues ($P < 0.01$; Figure 1B), and the expression of Bcl-2 in I, II, III, and IV grades of glioma was gradually increased; the difference was statistically significant ($P < 0.05$; Figure 1B). In addition, we found that the expression of NF- κ B1 in glioma cell lines, U87 and SHG44, was higher compared to noncancerous brain tissues ($P < 0.01$; Figure 1C). Therefore, the expression trend of NF- κ B1 and Bcl-2 in the same glioma specimens was consistent.

NF- κ B1 inhibits early apoptosis of glioma cells

The experiment was divided into blank group (mock), no-load plasmid group (negative), and NF- κ B1 shRNA plasmid group (NF- κ B1 shRNA). NF- κ B1 shRNA plasmids were transfected into glioma cell lines (U87 and SHG44). The cells were collected from each group at 24, 48, and 72 h after transfection. The experimental procedure was performed and detected the proportion of cells in early-stage apoptosis by flow cytometry (Figure 2A–F). Results showed that the apoptotic rate of glioma cell lines (U87 and SHG44) was significantly increased after silencing of NF- κ B1 expression compared to blank and no-load plasmid groups. The difference was statistically significant ($P < 0.01$; Figure 3A and B).

NF- κ B1 affects the expression of Bcl-2 mRNA

The relative expression levels of NF- κ B1 and Bcl-2 mRNA in each group (blank control group, no-load plasmid group, NF- κ B1 overexpression group, NF- κ B1 silencing group, and BCL2 silencing group) were analyzed by qRT-PCR. The relative quantitative analysis of NF- κ B1 mRNA expression showed that compared to the blank and no-load plasmid groups $2^{-\Delta\Delta Ct}$ value, NF- κ B1 shRNA plasmid group $2^{-\Delta\Delta Ct}$ value was significantly decreased and the difference was statistically significant (compared with the blank group and no-load plasmid group, the expression of Bcl-2 in U87 and SHG44, $P < 0.05$; Table 1). In addition, the expression of NF- κ B1 mRNA showed no increase after silencing the expression of Bcl-2 and no statistical significance was observed (compared with the blank group and no-load plasmid group, the expression of NF- κ B1 in U87 and SHG44, $P > 0.05$; Table 1). The relative quantitative analysis of Bcl-2 mRNA expression showed that the expression of Bcl-2 mRNA was significantly decreased after silencing NF- κ B1 gene expression compared to blank and no-load plasmid groups, and the difference was statistically significant (compared with the blank group and no-load plasmid group, the expression of Bcl-2 in U87 and SHG44, $P < 0.05$; Table 1).




Figure 1 Expression of NF-κB1 and Bcl-2 detection in glioma tissues by qRT-PCR.

Notes: The expression of NF-κB1 in glioma was significantly higher than that in nonneoplastic brain tissues ($P < 0.01$; **A**), and the expression of NF-κB1 in I, II, III, and IV grades of glioma was gradually increased, and the difference was statistically significant ($P < 0.05$; **A**). At the same time, the Bcl-2 expression in the same brain glioma specimens was significantly higher than that of the nonneoplastic brain tissues ($P < 0.01$; **B**), and the expression of Bcl-2 in I, II, III, and IV grades of glioma was gradually increased, and the difference was statistically significant ($P < 0.05$; **B**). In addition, we found that the expression of NF-κB1 in glioma cell lines, U87 and SHG44, was higher than the noncancerous brain tissues ($P < 0.01$; **C**). The same entry represents a class of tissue specimens, such as: Norm, nonneoplastic brain tissues; PA, Grade I; DA, Grade II; AA, Grade III; GBM, primary brain GBM, Grade IV. Gliomas were divided into four grades (Grade I, Grade II, Grade III, and Grade IV) according to the 2007 WHO classification system.

Abbreviations: AA, anaplastic astrocytoma; DA, diffuse astrocytoma; GBM, glioblastoma; NF-κB1, nuclear factor-κB1; PA, pilocytic astrocytoma; qRT-PCR, quantitative real-time polymerase chain reaction; WHO, World Health Organization.

On the contrary, the expression of Bcl-2 mRNA was increased after overexpression of NF-κB1, and the difference was statistically significant (compared with the blank group and no-load plasmid group, the expression of Bcl-2 in U87 and SHG44, $P < 0.05$; Table 1).

qRT-PCR was performed to evaluate the mRNA levels of blank group, no-load plasmid group, and NF-κB1 overexpression + Bcl-2 shRNA plasmid groups. Results showed that compared to blank and no-load plasmid groups, the expression of Bcl-2 mRNA was not significantly reduced after NF-κB1 overexpression and silencing of Bcl-2 expression, and the difference was not statistically significant (compared with the blank group and no-load plasmid group, the expression of Bcl-2 in U87 and SHG44, $P > 0.05$; Table 1). These results confirmed that NF-κB1 affected the expression of Bcl-2 mRNA.

NF-κB1 affects the expression of Bcl-2 protein

After 72 h of transfection, the total protein was extracted from each group (blank control group, no-load plasmid

group, NF-κB1 silencing group, NF-κB1 overexpression group, Bcl-2 silencing group, and NF-κB1 overexpression + Bcl-2 silencing group). The expression levels of NF-κB1 and Bcl-2 proteins were analyzed by Western blotting (Figure 4) and the gray value (Figure 5). Results showed that the expression of NF-κB1 and Bcl-2 proteins in NF-κB1 shRNA plasmid group was significantly decreased compared to the blank group and no-load plasmid group (Figure 4A and B), and the difference was statistically significant ($P < 0.01$; Figure 5A and B). The expression of NF-κB1 and Bcl-2 proteins was increased significantly in NF-κB1 overexpression plasmid group (Figure 4C and D), and the difference was statistically significant ($P < 0.05$; Figure 5C and D). The expression of Bcl-2 protein was not significantly reduced in NF-κB1 overexpression + Bcl-2 silencing group (Figure 4E and F), and the difference was not statistically significant ($P > 0.05$; Figure 5E and F). In addition, the expression of NF-κB1 protein was not significantly increased after silencing of Bcl-2 gene expression (Figure 4C and D), and the difference was not statistically significant ($P > 0.05$; Figure 5C and D).




Figure 2 Early apoptotic rate detection in glioma cells by flow cytometry.

Notes: The early apoptotic rate of glioma cells (U87 and SHG44) was detected by flow cytometry at 72 h. (A–C) U87 cells, (D–F) SHG44 cells, (A and D) mock, (B and E) negative, and (C and F) NF-κB1 shRNA. Mock, blank group; negative, no-load plasmid group; NF-κB1 shRNA, NF-κB1 silencing group.

Abbreviation: NF-κB1, nuclear factor-kappa B1.

Discussion

NF-κB1 gene is located on human chromosome 4q24 and is a key transcription factor of the NF-κB signaling pathway. In recent years, NF-κB1 has been shown to be involved in a variety of malignancies and is widely expressed in several types of malignant tumors.^{13,14} Previous studies have shown that NF-κB1 can promote invasion of glioma cell lines by

MMP9.¹² However, the correlation between NF-κB1 and Bcl-2 expressions in gliomas and the underlying mechanism between NF-κB1 and early apoptosis of glioma cells have not been reported so far.

Glioma is one of the most common types of adult primary brain tumors.¹⁵ The unrestricted growth of tumors is not only due to abnormal cell differentiation and




Figure 3 Early apoptotic rate detection of glioma cells by histograms.

Notes: The apoptotic rate of glioma cell lines (U87 and SHG44) was significantly increased after silencing of NF-κB1 expression compared to mock and negative. The difference was statistically significant ($P < 0.01$). (A) U87 cells and (B) SHG44 cells; ** $P < 0.01$. Mock, blank group; negative, no-load plasmid group; NF-κB1 shRNA, NF-κB1 silencing group.

Abbreviation: NF-κB1, nuclear factor-kappa B1.

Table 1 qRT-PCR for detecting NF-κB1 and Bcl-2 expressions in the blank control group, no-load plasmid group, NF-κB1 shRNA group, NF-κB1 overexpression group, Bcl-2 shRNA group, and NF-κB1 overexpression + Bcl-2 shRNA group (mean ± SD)

| Group | Cases | U87 | | SHG44 | |
|---|-------|---------------------------|--------------------------|---------------------------|--------------------------|
| | | NF-κB1 | Bcl-2 | NF-κB1 | Bcl-2 |
| Blank control | 3 | 1.314±0.027 | 2.027±0.198 | 3.948±0.028 | 4.676±0.184 |
| No-load plasmid | 3 | 1.145±0.125 | 1.706±0.618 | 2.293±1.156 | 3.893±2.506 |
| NF-κB1 silencing | 3 | 0.485±0.049 ^a | 0.302±0.031 ^b | 0.858±0.091 ^a | 1.881±0.438 ^b |
| NF-κB1 overexpression | 3 | 16.625±1.334 ^a | 4.171±0.097 ^b | 14.023±0.591 ^a | 6.002±1.063 ^b |
| Bcl-2 silencing | 3 | 1.220±0.072 ^a | 0.700±0.072 ^b | 2.322±0.540 ^a | 1.666±0.017 ^b |
| NF-κB1 overexpression + Bcl-2 silencing | 3 | 15.818±0.292 | 1.848±0.342 ^b | 12.190±1.433 | 4.121±0.174 ^b |
| F value | | 91.249 | 0.433 | 16.025 | 0.230 |
| P-value | | 0.000 | 0.667 | 0.000 | 0.801 |

Note: Compared to the blank control group and no-load plasmid group: ^athe expression of NF-κB1 in U87 and SHG44, $P < 0.05$; ^bcompared with the blank group and no-load plasmid group, the expression of Bcl-2 in U87 and SHG44, $P < 0.05$.

Abbreviations: NF-κB1, nuclear factor-kappa B1; qRT-PCR, quantitative real-time polymerase chain reaction.

proliferation but is also closely associated with inhibition of cell apoptosis. Therefore, early apoptosis of glioma cells is of great significance. NF-κB1 acts as a DNA-binding subunit in NF-κB complex.¹⁶ After being stimulated by cytokines, such as TNF-α or DNA damage, the activated NF-κB then enters the nucleus and initiates the expression of downstream target genes, thereby activating a series of cellular responses.¹⁷ The study has reported that NF-κB pathway can enhance the expression of manganese superoxide dismutase to remove reactive oxygen species and

thus induces cell apoptosis.¹⁸ In addition, NF-κB can also inhibit cell apoptosis by negatively regulating the activation of c-Jun N-terminal kinase which is also known as stress-activated protein kinase and is a mitogen-activated protein kinase signaling pathway of another subclass.¹⁹ Finally, NF-κB can activate Akt and promote cell survival and proliferation by downregulating the expression of phosphatase and tensin homolog deleted on chromosome ten.²⁰ This study attempts to identify the expression of NF-κB1 in glioma cells. We found that NF-κB1 was

OncoTargets and Therapy downloaded from https://www.dovepress.com/ by 54.70.40.11 on 28-May-2018 For personal use only.




Figure 4 Analysis of NF-κB1 and Bcl-2 protein expressions in glioma cells by Western blotting.

Notes: Numbers 1–3 are blank group, no-load plasmid group, and NF-κB1 shRNA plasmid group; 4–6 are blank group, no-load plasmid group, and NF-κB1 overexpression + Bcl-2 shRNA plasmid group; 7–9 are no-load plasmid group, NF-κB1 overexpression plasmid group, and Bcl-2 shRNA plasmid group; (A, C, and E) U87 cells and (B, D, and F) SHG44 cells; relative molecular weight: NF-κB1: 50,000, Bcl-2: 26,000, and GAPDH: 36,000. NF-κB1 shRNA, NF-κB1 silencing; Bcl-2 shRNA, Bcl-2 silencing.

Abbreviation: NF-κB1, nuclear factor-kappa B1.




Figure 5 Effect of NF- κ B1 on Bcl-2 protein expressions.

Notes: (A, C, and E) The grayscale values of Western blotting bands of U87 cells. (B, D, and F) The grayscale values of Western blotting bands of SHG44 cells. Mock, blank control group; negative, no-load plasmid group; NF- κ B1 shRNA, NF- κ B1 silencing group; NF- κ B1, NF- κ B1 overexpression group; Bcl-2 shRNA, Bcl-2 silencing group; NF- κ B1 + Bcl-2 shRNA, NF- κ B1 overexpression + Bcl-2 silencing group. * $P < 0.05$; ** $P < 0.01$.

Abbreviation: NF- κ B1, nuclear factor-kappa B1.

highly expressed in most of the glioma cells, especially in high-grade glioma cells. Preliminary experiment results revealed that the expression of NF- κ B1 in human glioma was significantly higher than that in normal brain tissues, and the expression of NF- κ B1 in I, II, III, and IV grades of glioma was increased gradually; the difference was statistically significant. In this study, we found that the early apoptotic rate of U87 and SHG44 cells was significantly increased after NF- κ B1 shRNA transfection. It was found that the expression level of NF- κ B1 was related to the early apoptosis of glioma cells.

Antiapoptotic gene Bcl-2 plays a crucial role in carcinogenesis. It exists in mitochondria and has antiapoptotic protein effect.²¹ Bcl-2 is different from the general oncogene and tumor suppressor genes. The apoptotic gene controls cell survival and does not affect cell proliferation.²² The

antiapoptotic mechanisms of Bcl-2 include inhibition of ion channel opening that is activated by IP3R, which thereby inhibits the release of Ca²⁺ from the endoplasmic reticulum and produces antiapoptotic effect.²³ The coexpression of Bcl-2 and P53 genes could delay the role of P53 gene-induced apoptosis. Meanwhile, the synergistic effect of Bcl-2 and c-myc could block the entry of P53 gene into the nucleus, thus blocking the effect of P53 on apoptosis and growth arrest.²⁴ In addition, Bcl-2 can inhibit apoptosis by participating in signal transmission.²⁵ Few studies have demonstrated a close association of Bcl-2 in glioma development.^{26,27} The results of this study showed that Bcl-2 expression in human glioma specimens was significantly higher than normal brain tissues, and the expression of Bcl-2 in I, II, III, and IV grades of glioma was increased gradually; the difference was statistically significant. Hussein et al²⁸

demonstrated that the expression of Bcl-2 protein was increased gradually with the increasing malignancy of human gliomas, and our study results were consistent with these. It was found that the expression trend of Bcl-2 and NF- κ B1 was similar in the same glioma specimens, which indicated that NF- κ B1 was closely related with Bcl-2. In vitro experiments revealed that silencing of NF- κ B1 expression could inhibit the expression of Bcl-2, while overexpression of NF- κ B1 could promote the expression of Bcl-2. Furthermore, overexpression of NF- κ B1 silenced the expression of Bcl-2 by salvage strategy of the experiment. Results showed that the expression level of Bcl-2 was not significantly decreased. These data suggest that NF- κ B1 can promote the expression of Bcl-2.

Conclusion

NF- κ B1 overexpression can inhibit the early apoptosis of glioma cells, and high expression of NF- κ B1 can promote the expression of antiapoptotic gene Bcl-2, which is closely related to each other. In glioma cells, antiapoptotic mechanism of NF- κ B1 may be achieved by promoting Bcl-2 expression and provides a theoretical basis for studying the antiapoptotic mechanism of tumor cells in association with NF- κ B1.

Acknowledgments

We thank our central laboratory for providing technical instruction and assistance. This work was partly supported by the Fund Projects: National Natural Science Foundation of China (NSFC 81372689), the Jiangsu Province Health and Family Planning Commission (Q201606), and the Suzhou City Health Bureau Science and Education project (kjxw2016020).

Disclosure

The authors report no conflicts of interest in this work.

References

- Furnari FB, Fenton T, Bachoo RM, et al. Malignant astrocytic glioma: genetics, biology, and paths to treatment. *Genes Dev.* 2007;21(21):2683–2710.
- Tang SL, Gao YL, Hu WZ. PAQR3 inhibits the proliferation, migration and invasion in human glioma cells. *Biomed Pharmacother.* 2017;18(92):24–32.
- Chen B, Lei Y, Wang H, et al. Repression of the expression of TET2 by ZEB1 contributes to invasion and growth in glioma cells. *Mol Med Rep.* 2017;15(5):2625–2632.
- Bao Z, Xu X, Liu Y, et al. CBX7 negatively regulates migration and invasion in glioma via Wnt/ β -catenin pathway inactivation. *Oncotarget.* 2017;5(28):1–16.
- Gabriely G, Wurdinger T, Kesari S, et al. MicroRNA 21 promotes glioma invasion by targeting matrix metalloproteinase regulators. *Mol Cell Biol.* 2008;28(17):5369–5380.
- Mao K, You C, Lei D, Zhang H. Potential regulation of glioma through the induction of apoptosis signaling via Egl-9 family hypoxia-inducible factor 3. *Oncol Lett.* 2017;13(2):893–897.
- Sato A, Mizobuchi Y, Nakajima K, et al. Blocking COX-2 induces apoptosis and inhibits cell proliferation via the Akt/survivin- and Akt/ID3 pathway in low-grade-glioma. *J Neurooncol.* 2017;132(2):231–238.
- Shuang T, Wang M, Zhou Y, Shi C, Wang D. NF- κ B1, c-Rel, and ELK1 inhibit miR-134 expression leading to TAB1 upregulation in paclitaxel-resistant human ovarian cancer. *Oncotarget.* 2017;8(15):24853–24868.
- Wang X, Peng H, Liang Y, et al. A functional insertion/deletion polymorphism in the promoter region of the NFKB1 gene increases the risk of papillary thyroid carcinoma. *Genet Test Mol Biomarkers.* 2015;19(3):167–171.
- Lin SC, Liu CJ, Yeh WI, Lui MT, Chang KW, Chang CS. Functional polymorphism in NFKB1 promoter is related to the risks of oral squamous cell carcinoma occurring on older male areca (betel) chewers. *Cancer Lett.* 2006;243(1):47–54.
- Mohd Suzairi MS, Tan SC, Ahmad Aizat AA, et al. The functional-94 insertion/deletion ATTG polymorphism in the promoter region of NFKB1 gene increases the risk of sporadic colorectal cancer. *Cancer Epidemiol.* 2013;7(5):634–638.
- Yang TQ, Lu XJ, Wu TF, et al. MicroRNA-16 inhibits glioma cell growth and invasion through suppression of BCL2 and the nuclear factor- κ B1/MMP9 signaling pathway. *Cancer Sci.* 2014;105(3):265–271.
- Shuang T, Wang M, Zhou Y, Shi C. Over-expression of nuclear NF- κ B1 and c-Rel correlates with chemoresistance and prognosis of serous epithelial ovarian cancer. *Exp Mol Pathol.* 2016;100(1):139–144.
- Kravtsova-Ivantsiv Y, Shomer I, Cohen-Kaplan V, et al. KPC1-mediated ubiquitination and proteasomal processing of NF- κ B1 p105 to p50 restricts tumor growth. *Cell.* 2015;161(2):333–347.
- Wiese M, Walther N, Diederichs C, et al. The β -catenin/CBP-antagonist ICG-001 inhibits pediatric glioma tumorigenicity in a Wnt-independent manner. *Oncotarget.* 2017;8(16):27300–27313.
- Jacque E, Schweighoffer E, Visekruna A, et al. IKK-induced NF- κ B1 p105 proteolysis is critical for B cell antibody responses to T cell-dependent antigen. *J Exp Med.* 2014;211(10):2085–2101.
- Huang H, Ma L, Li J, et al. NF- κ B1 inhibits c-Myc protein degradation through suppression of FBW7 expression. *Oncotarget.* 2014;5(2):493–505.
- Hussain AR, Ahmed SO, Ahmed M, et al. Cross-talk between NF κ B and the PI3-kinase/AKT pathway can be targeted in primary effusion lymphoma (PEL) cell lines for efficient apoptosis. *PLoS One.* 2012;7(6):1–12.
- Xuan Y, Chi L, Tian H, et al. The activation of the NF- κ B-JNK pathway is independent of the PI3K-Rac1-JNK pathway involved in the bFGF-regulated human fibroblast cell migration. *J Dermatol Sci.* 2016;82(1):28–37.
- Ma J, Liu J, Wang Z, et al. NF-kappaB-dependent microRNA-425 upregulation promotes gastric cancer cell growth by targeting PTEN upon IL-1 β induction. *Mol Cancer.* 2014;26(13):1–11.
- Wang JH, Bi XW, Li PF, et al. Overexpression of MYC and BCL2 predicts poor prognosis in patients with extranodal NK/T-cell lymphoma, nasal type. *J Cancer.* 2017;8(5):793–800.
- Conti A, Majorini MT, Fontanella E, et al. Lemur tyrosine kinase 2 (LMTK2) is a determinant of cell sensitivity to apoptosis by regulating the levels of the BCL2 family members. *Cancer Lett.* 2017;28(389):59–69.
- Singh K, Briggs JM. Functional implications of the spectrum of BCL2 mutations in lymphoma. *Mutat Res Rev Mutat Res.* 2016;6(16):1–18.
- Patel JH, McMahon SB. BCL2 is a downstream effector of MIZ-1 essential for blocking c-MYC-induced apoptosis. *J Biol Chem.* 2007;282(1):5–13.

25. Song W, Song C, Chen Y, et al. Polysaccharide-induced apoptosis in H22 cells through G2/M arrest and BCL2/BAX caspase-activated Fas pathway. *Cell Mol Biol (Noisy-le-grand)*. 2015;61(7):88–95.
26. Harmalkar M, Upraity S, Kazi S, Shirsat NV. Tamoxifen-induced cell death of malignant glioma cells is brought about by oxidative-stress-mediated alterations in the expression of BCL2 family members and is enhanced on miR-21 inhibition. *J Mol Neurosci*. 2015;57(2):197–202.
27. Li W, Qian C, Wang L, et al. Association of BCL2-938C>A genetic polymorphism with glioma risk in Chinese Han population. *Tumour Biol*. 2014;35(3):2259–2264.
28. Hussein MR, El-Ghorori RM, El-Rahman YG. Alterations of p53, BCL-2, and hMSH2 protein expression in the normal brain tissues, gliosis, and gliomas. *Int J Exp Pathol*. 2006;87(4):297–306.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress