

Submit a Manuscript: <https://www.f6publishing.com>*World J Clin Cases* 2021 April 16; 9(11): 2619-2626DOI: [10.12998/wjcc.v9.i11.2619](https://doi.org/10.12998/wjcc.v9.i11.2619)

ISSN 2307-8960 (online)

CASE REPORT

Cryptococcus infection with asymptomatic diffuse pulmonary disease in an immunocompetent patient: A case report

Yong Li, Lei Fang, Fang-Qun Chang, Fang-Zhou Xu, Yan-Bei Zhang

ORCID number: Yong Li [0000-0002-2091-3793](https://orcid.org/0000-0002-2091-3793); Lei Fang [0000-0001-6798-2932](https://orcid.org/0000-0001-6798-2932); Fang-Qun Chang [0000-0002-7951-8640](https://orcid.org/0000-0002-7951-8640); Fang-Zhou Xu [0000-0002-3101-8727](https://orcid.org/0000-0002-3101-8727); Yan-Bei Zhang [0000-0003-0551-4054](https://orcid.org/0000-0003-0551-4054).

Author contributions: Li Y, Chang FQ, and Xu FZ collected the clinical data and wrote the paper; Zhang YB and Fang L contributed to the design and revision of the paper.

Supported by The Colleges and Universities in Anhui Province Natural Science Research Projects, No. KJ2018A0208.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflicts of interest to report.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in

Yong Li, Lei Fang, Fang-Qun Chang, Fang-Zhou Xu, Yan-Bei Zhang, Department of Geriatric Respiratory and Critical Care, The First Affiliated Hospital of Anhui Medical University, Hefei 230000, Anhui Province, China

Yong Li, Department of Tuberculosis, Anhui Medical University Clinical College of Chest & Anhui Chest Hospital, Hefei 230000, Anhui Province, China

Corresponding author: Yan-Bei Zhang, MAMS, Chief Doctor, Professor, Department of Geriatric Respiratory and Critical Care, The First Affiliated Hospital of Anhui Medical University, No. 218 Jixi Street, Hefei 230000, Anhui Province, China.

zhangyanbei1963@126.com

Abstract

BACKGROUND

Cryptococcus presenting as an opportunistic pathogen mainly affects immunocompromised patients, but the disseminated form of infection is rare among immunocompetent populations. The partial radiographic characteristics of pulmonary cryptococcosis mimic lung carcinoma, leading to unnecessary open chest exploratory surgery, and the lack of a gold-standard noninvasive diagnostic increases the risk of misdiagnosis. Positron emission tomography/computed tomography (PET/CT), a sensitive method for distinguishing malignant tumors, coupled with cryptococcal latex agglutination test showing a high positive rate may overcome these issues.

CASE

A 36-year-old man presented for general examination, without health complaints. Routine CT showed multiple pulmonary nodules and a mass with high maximum standardized uptake value. Initially, we suspected primary malignancy with hematogenous metastasis. Although his routine fungal analysis had been negative, subsequent CT-guided percutaneous core needle biopsy and histopathology examination indicated a diagnosis of pulmonary cryptococcosis. Fluconazole (200 mg/d) antifungal drug treatment was initiated, and 1 mo later the pulmonary mass had reduced in size markedly (on chest CT scan) without any complications.

CONCLUSION

Serologic and PET/CT examinations may not rule out cryptococcosis, and percutaneous lung puncture is critical under all circumstances.

accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Respiratory system

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

Received: December 7, 2020

Peer-review started: December 7, 2020

First decision: December 21, 2020

Revised: January 9, 2021

Accepted: February 10, 2021

Article in press: February 10, 2021

Published online: April 16, 2021

P-Reviewer: Tripathy SK

S-Editor: Zhang H

L-Editor: Wang TQ

P-Editor: Li JH



Key Words: *Cryptococcus*; Immunocompetent; Positron emission tomography; Computed tomography; Cryptococcal latex agglutination test; Lung diseases; Fungal; Case report

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Cryptococcosis is a systemic fungal infection, with presentation ranging from asymptomatic pulmonary involvement to meningitis and disseminated disease. Generally, cryptococcosis is considered an opportunistic infection in immunocompromised individuals, affecting persons with human immunodeficiency virus infection in particular. Mass-like and disseminated cryptococcosis lesions are rare in patients with normal immunity, and a negative result on the cryptococcal latex agglutination test may be related to low-grade virulence of the fungus. We describe here a case of pulmonary cryptococcosis that was likely caused by a low-virulence strain, presenting bilaterally distributed lesions on imaging and mimicking hematogenous metastasis.

Citation: Li Y, Fang L, Chang FQ, Xu FZ, Zhang YB. *Cryptococcus* infection with asymptomatic diffuse pulmonary disease in an immunocompetent patient: A case report. *World J Clin Cases* 2021; 9(11): 2619-2626

URL: <https://www.wjgnet.com/2307-8960/full/v9/i11/2619.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v9.i11.2619>

INTRODUCTION

Cryptococcosis is a systemic infection caused by two species of the encapsulated yeast-like fungus *Cryptococcus*: *Cryptococcus gattii* (*C. gattii*) and *Cryptococcus neoformans* (*C. neoformans*)^[1]. These opportunistic fungal pathogens usually cause cryptococcosis in immunocompromised patients, with the respiratory system being the usual route of infection. Infections in immunocompetent persons are much rarer.

Positron emission tomography/computed tomography (PET/CT) is a useful tool for differentiating benign and malignant tissues. Higher uptake values of fluorodeoxyglucose on PET/CT reflect the characteristics of a malignant disease. Unfortunately, since the radiographic characteristics of pulmonary cryptococcosis mimic lung carcinoma, clinicians may order open chest exploratory surgery, with patients bearing the burden of undergoing an unnecessary invasive procedure^[2].

Cases of pulmonary cryptococcosis, therefore, require further noninvasive testing for a definitive diagnosis; yet, no gold-standard noninvasive diagnostic exists and the risk of misdiagnosis remains large. Prior to a biopsy and histological assessment [by Grocott's methenamine silver (GMS) staining], the most common suspected diagnoses are lung cancer, inflammatory granuloma, metastases from an extrathoracic malignancy, and pulmonary tuberculosis.

As described below for our case, CT-guided percutaneous core needle biopsy of a presumed lung carcinoma can be critical for a definitive diagnosis of *Cryptococcus* infection.

CASE PRESENTATION

Chief complaints

A 34-year-old apparently immunocompetent Chinese man presented for general health examination. On routine CT, multiple pulmonary nodules and a mass were incidentally found.

History of present illness

The patient reported having had no symptoms of discomfort prior to presentation, including cough, sputum, chest pain, chills, or fever. He also denied tiring easily, having night sweats, and rapid weight loss.

History of past illness

The patient reported occupation as an engineer, smoking ten cigarettes a day for the past 3 years, and having no pets. He also had no history of allergies, hepatitis B, or diabetes mellitus, or any other significant medical history.

Physical examination

Physical examination upon admission found bilateral lungs to be clear, with no wheezing, rales, or rhonchi detected by auscultation. Temperature was 36.5 °C. Heart rate was 95 beats/min. Respiratory rate was 20 breaths/min. Blood pressure was 104/71 mmHg. There were no other remarkable findings.

Laboratory examinations

Test findings for liver function, renal function, and electrolytes were normal. The values of fasting blood glucose were within the normal range twice during hospitalization. All inflammatory biomarkers were also within the normal range. The result of human immunodeficiency virus (HIV) serology test was negative. Tumor markers in blood, including carcinoembryonic antigen, progastrin-releasing peptide, neuron-specific enolase, and cytokeratin 19-fractions (CYFRA 21.1), were all normal. Serology test for *Cryptococcus* antigen also gave negative results.

Imaging examinations

Due to the pulmonary lesions of unknown cause, PET/CT was arranged. The imaging showed multiple nodules in both lungs, enlarged right hilar lymph nodes, and an elliptic mass lesion without clear margin in the right lower lobe (Figure 1). The mass showed higher uptake values of ¹⁸F-fluorodeoxyglucose (FDG) and SUVmax of 16.4 in normal scans and 19.1 in delayed scans. The SUVmax for FDG uptake of those nodules was 6.8 in normal scans and 10.2 in delayed scans. The two enlarged lymph nodes in the right hilum gave 4.8 in normal scans and 6.9 in delayed scans.

Further diagnostic work-up

Subsequently, the patient underwent CT-guided percutaneous core needle biopsy of the right lower-lobe mass. No fungi or remarkable bacteria were found in the puncture fluid smear.

MULTIDISCIPLINARY EXPERT CONSULTATION

Department of Pathology, Nanjing Gulou Hospital Affiliated to Medical School of Nanjing University

The pathologic consultation found granulomatous inflammation. GMS was positive and periodic acid-Schiff staining was negative (Figure 2). No evidence of malignancy was noted. In summary, a diagnosis of cryptococcal pneumonia was established.

FINAL DIAGNOSIS

A diagnosis of pulmonary cryptococciosis was made according to the radiologic evidence of pulmonary lesions and positive histopathology, although the serology test for *Cryptococcus* antigen was negative.

TREATMENT

Fluconazole (200 mg) per day orally.

OUTCOME AND FOLLOW-UP

After 1 mo and 3 mo of treatment, the pulmonary lesions were found to be markedly reduced in size on chest CT (Figures 3 and 4).




Figure 1 Positron emission tomography/computed tomography showing bilateral lesions. An elliptic mass without clear margin was shown in the right lower lobe. The size of the mass was 3.46 cm × 2.39 cm. The enlarged right hilar lymph nodes showed a high value of SUV_{max}. There were also scattered nodules found in both lungs.




Figure 2 Pathological examination revealed *Cryptococcus* infection. A: Grocott's methenamine silver staining showed *Cryptococcus* spores by black staining (arrow); B: Periodic acid-Schiff staining was negative. Original magnification: × 400.

DISCUSSION

Pulmonary cryptoccosis is prevalent in patients with compromised cell-mediated immunity, such as those with HIV infection or who have been subject to chemotherapy or other immunosuppressive therapy^[3,4]. It is known that patients with HIV infection are more likely to have central nervous system involvement with the cryptococcosis disease than those without. In that group, the disease itself is associated with a high mortality rate. Cryptococcosis can also occur in individuals with normal immunity and causes severe nervous system infections^[5], but such cases are rare. Table 1 illustrates the special features of our report.

For our case, the patient was otherwise healthy and had no neurological symptoms. However, he presented bilaterally disseminated lesions in the lung, which is extremely rare for an asymptomatic immunocompetent person, and this increased the chance of mistaken and missed diagnosis. Interestingly, his cryptococcal latex agglutination test (CLAT) was also negative. In other reports in the literature, the positive rate for CLAT has been high, even exceeding 90%^[6,7]. CLAT is also used to evaluate response to therapy for pulmonary cryptoccosis in patients without HIV infection^[7].

Most cryptococcal strains are encapsulated by polysaccharides^[8] and capsule-deficient *C. neoformans* may have reduced virulence^[9]. A cryptococcal strain that has lost its capsular material can, however, trigger strong inflammatory reactions, which tends to lead to the formation of granulomas composed of histiocytes, giant cells, and lymphocytes. This is consistent with our pathological outcome. We speculated that our patient may have been infected with a low-virulence *C. neoformans* strain, thus causing

Table 1 *Cryptococcus* infection in immunocompetent patients

Case	Sex, age	Medical history	Diagnostic modalities	Clinical presentation	Imaging	Immuno-assay	Histopathology	Treatment
1, this report	M36	Good condition	CT-guided percutaneous core needle biopsy	None	Pulmonary nodules and a mass in both lungs; PET-CT showed the SUVmax of the cryptococcal lesions fluctuated from 4.8 to 19.1	Negative	Granulomatous inflammation; GMS was positive and PSA was negative	Fluconazole
2, Bavishi et al ^[20]	F67	Hypertension and cholelithiasis	CT-guided percutaneous core needle biopsy	Recurrent dry cough for 4 yr	Multiple pulmonary nodules in both the lower lobes	Serology cryptococcal antigen titer of 1: 32	FMS staining was strongly positive	Intravenous amphotericin B for 2 d and then changed to fluconazole
3, Zhou et al ^[21]	M44	Good condition	Bronchoscopy	3-mo history of cough, hemoptysis	Pulmonary nodules in both lungs; PET-CT showed the SUVmax of the cryptococcal lesions fluctuated from 9.86 to 10.99	Titer of more than 1: 1, 280	GMS stain was positive; Culture of bronchoscopy with brush was positive	Amphotericin B
4, Marroni et al ^[22]	F21	Good condition	CT-guided percutaneous core needle biopsy	Rigors, fever, dyspnoea, dry cough, and chest pain	A round mass in the lung	Positive at a titre of 1: 256	GMS and PSA staining was positive	Fluconazole
5, Oliveira et al ^[3]	M64	Arterial hypertension	Fiberoptic bronchoscopy with bronchoalveolar lavage	Fever, weakness, anorexia, headache, dyspnea, cough, purulent sputum production, and disorientation	Pulmonary spherical mass lesion, 5 cm in diameter	The CSF cryptococcal antigen titer was 1: 4096 with a serum titer of 1:2048	MGG staininh was positive	Amphotericin B
6, Ruan et al ^[12]	M68	Good condition	Surgical drainage	Progressive multiple abscesses, fever, lower extremity weakness, and urinary retention	Pulmonary abscess formation and multiple destruction of vertebral bodies	Negative	Culture revealed; <i>Cryptococcus neoformans</i> ; India ink staining	Itraconazole; Fluorocytosine; Fluconazole

FMS: Fontana-Masson silver; CT: Computed tomography; PET: Positron emission tomography; GMS: Grocott's methenamine-silver staining; CSF: Cerebrospinal fluid; COPD: Chronic obstructive pulmonary disease; PAS: Periodic acid-Schiff staining; MGG: May-Grunwald Giemsa.

the false negative CLAT result^[10]. Certainly, other unknown possibilities cannot be excluded.

Of the two *Cryptococcus* species, *C. neoformans* is widespread, especially in pigeon feces or in soil contaminated with these feces, and it tends to cause disease in patients who are immunocompromised. In Taiwan, there is evidence to show a significant prevalence of *C. gattii* infection associated with the eucalyptus tree^[11]. *C. gattii* usually causes disease in immunocompetent persons and behaves as a primary pathogen. The percentage of *C. gattii* infection in immunocompetent hosts is significantly higher than that of *C. neoformans* infection^[12].

In 2018, the World Health Organization recommended that without the results of cerebrospinal fluid, rapid serum, plasma, or whole-blood cryptococcal antigen assays are the preferred diagnostic methods (either lateral flow assay or latex agglutination assay). Some studies have reported that CLAT has an insufficient sensitivity for *C. gattii* compared to the lateral flow assay, increasing the risk of delayed diagnosis and probability of misdiagnosis^[3]. As such, in order to diagnose cryptococcosis, multiple detection methods should be applied to improve the accuracy of diagnosis.

C. gattii infection is more common in tropical areas^[11], which is inconsistent with the geography of our patient's residential area. During the subsequent visit, the patient told us that one of his neighbors kept pigeons, which supported our speculation that the patient was likely to have been infected with a low-virulence strain of *C. neoformans*.




Figure 3 Chest computed tomography scan after 1 mo of antifungal treatment showing resolution of the bilateral lesions. The mass in the right lung was reduced markedly. A: The 25th floor scan; B: The 26th floor scan.




Figure 4 Chest computed tomography scan after 3 mo of antifungal treatment showing near complete disappearance of the nodules and infiltration distributed around the lesions. Only a small number of pulmonary cavities remained at this time. A: The 29th floor scan; B: The 30th floor scan.

The most common radiographic characteristics of pulmonary cryptococcosis consist of solitary pulmonary mass or nodules, combined patchy and nodular shadows, cavitation, and enlarged mediastinal lymph nodes^[14]. However, the most common imaging finding of immunocompetent patients is single nodule. Mass-like and bronchopneumonic patterns are common in immunocompromised patients and rare in the patients with normal immunity^[15]. PET/CT is a sensitive method for distinguishing malignant tumors from benign tissue, and it has gradually become a useful imaging technology for the differential diagnosis, staging, and follow-up of cancer^[16]. The related SUVmax quantitatively measures the FDG uptake of a tumor.

Generally, the SUVmax value of 2.5 is used as the cut-off for differentiating benign lesions from malignant ones, as it is known to have the best sensitivity and specificity for such. However, the diagnostic accuracy of this value is not high in pulmonary infectious diseases, especially for those due to granulomatous diseases^[17] such as in our patient. The SUVmax values for cases of pulmonary cryptococcosis may vary widely, indicating anywhere from mild to marked uptake. In previous reports, the SUVmax of the cryptococcal lesions fluctuate from 0.93 to 13.0^[15,18], with most cases exceeding the 2.5 threshold. Nevertheless, few of the SUVmax values of pulmonary lesions reach 19.

The mechanism of increased FDG uptake in cryptococcal lesions may be related to the “respiratory burst”, which increases cellular glucose metabolism. Based on this principle, PET/CT can play a role in assessing the treatment of patients with fungal infection^[19]. Therefore, diagnosis of malignant tumor by PET/CT should exclude the possibility of fungal infection. Our case was remarkable since we could not find evidence of immunocompromise in our patient; additional reports of such cases and bench-to-bedside studies will help to explain the development of masses and diffuse lesions in patients who are infected with a low-virulence strain (as we suspect our

patient was).

CONCLUSION

We describe here an immunocompetent patient with pulmonary *Cryptococcus* infection, who manifested a mass and diffuse lesions in his lung while being asymptomatic. Although it is very rare, cryptococcosis must be included in the differential diagnosis of pulmonary lesions in patients without immunosuppression. It is important to recognize that pulmonary fungal infection can mimic metastases, even when the patient has been diagnosed with lung cancer by PET/CT. Furthermore, the potential for false negative results of fungal smears and serologic testing for cryptococcosis is another risk for misdiagnosis of this infectious disease. Thus, percutaneous lung puncture is critical under all circumstances in order to promote more timely initiation of appropriate treatment and avoidance of unnecessary surgery.

REFERENCES

- 1 Li SS, Mody CH. Cryptococcus. *Proc Am Thorac Soc* 2010; **7**: 186-196 [PMID: 20463247 DOI: 10.1513/pats.200907-063AL]
- 2 Sun LM, Chen TY, Chen WJ, Hsieh MJ, Liu JW, Huang CC, Wang CJ. Cryptococcus infection in a patient with nasopharyngeal carcinoma: imaging findings mimicking pulmonary metastases. *Br J Radiol* 2002; **75**: 275-278 [PMID: 11932223 DOI: 10.1259/bjr.75.891.750275]
- 3 Ou KW, Hsu KF, Cheng YL, Hsu GC, Hsu HM, Yu JC. Asymptomatic pulmonary nodules in a patient with early-stage breast cancer: Cryptococcus infection. *Int J Infect Dis* 2010; **14**: e77-e80 [PMID: 19477671 DOI: 10.1016/j.ijid.2009.03.007]
- 4 Tamagawa S, Hotomi M, Yuasa J, Tuchihashi S, Yamauchi K, Togawa A, Yamanaka N. Primary laryngeal cryptococcosis resembling laryngeal carcinoma. *Auris Nasus Larynx* 2015; **42**: 337-340 [PMID: 25701260 DOI: 10.1016/j.anl.2015.01.003]
- 5 Oliveira Fde M, Severo CB, Guazzelli LS, Severo LC. Cryptococcus gattii fungemia: report of a case with lung and brain lesions mimicking radiological features of malignancy. *Rev Inst Med Trop Sao Paulo* 2007; **49**: 263-265 [PMID: 17823759 DOI: 10.1590/s0036-46652007000400014]
- 6 Berlin L, Pincus JH. Cryptococcal meningitis. False-negative antigen test results and cultures in nonimmunosuppressed patients. *Arch Neurol* 1989; **46**: 1312-1316 [PMID: 2686594 DOI: 10.1001/archneur.1989.00520480054020]
- 7 Lin TY, Yeh KM, Lin JC, Wang NC, Peng MY, Chang FY. Cryptococcal disease in patients with or without human immunodeficiency virus: clinical presentation and monitoring of serum cryptococcal antigen titers. *J Microbiol Immunol Infect* 2009; **42**: 220-226 [PMID: 19812855]
- 8 Zeng Y, Wu X, Yi X, Luo B, Zhu M, Rui W, Zhu X, Li X, Li H. Clinicopathologic and ultrastructural study of non-HIV-related primary pulmonary cryptococcosis in China: report of 43 cases. *Ultrastruct Pathol* 2011; **35**: 19-25 [PMID: 21214404 DOI: 10.3109/01913123.2010.521293]
- 9 Kronstad JW, Attarian R, Cadieux B, Choi J, D'Souza CA, Griffiths EJ, Geddes JM, Hu G, Jung WH, Kretschmer M, Saikia S, Wang J. Expanding fungal pathogenesis: Cryptococcus breaks out of the opportunistic box. *Nat Rev Microbiol* 2011; **9**: 193-203 [PMID: 21326274 DOI: 10.1038/nrmicro2522]
- 10 Gazzoni AF, Severo CB, Salles EF, Severo LC. Histopathology, serology and cultures in the diagnosis of cryptococcosis. *Rev Inst Med Trop Sao Paulo* 2009; **51**: 255-259 [PMID: 19893977 DOI: 10.1590/s0036-46652009000500004]
- 11 Chen YC, Chang SC, Shih CC, Hung CC, Luhbd KT, Pan YS, Hsieh WC. Clinical features and in vitro susceptibilities of two varieties of *Cryptococcus neoformans* in Taiwan. *Diagn Microbiol Infect Dis* 2000; **36**: 175-183 [PMID: 10729660 DOI: 10.1016/s0732-8893(99)00137-6]
- 12 Ruan Q, Zhu Y, Chen S, Zhu L, Zhang S, Zhang W. Disseminated cryptococcosis with recurrent multiple abscesses in an immunocompetent patient: a case report and literature review. *BMC Infect Dis* 2017; **17**: 369 [PMID: 28558705 DOI: 10.1186/s12879-017-2459-9]
- 13 Tintelnot K, Hagen F, Han CO, Seibold M, Rickerts V, Boekhout T. Pitfalls in Serological Diagnosis of *Cryptococcus gattii* Infections. *Med Mycol* 2015; **53**: 874-879 [PMID: 26337090 DOI: 10.1093/mmy/mv061]
- 14 Ye F, Xie JX, Zeng QS, Chen GQ, Zhong SQ, Zhong NS. Retrospective analysis of 76 immunocompetent patients with primary pulmonary cryptococcosis. *Lung* 2012; **190**: 339-346 [PMID: 22246551 DOI: 10.1007/s00408-011-9362-8]
- 15 Wang SY, Chen G, Luo DL, Shao D, Liu ET, Sun T, Wang SX. ¹⁸F-FDG PET/CT and contrast-enhanced CT findings of pulmonary cryptococcosis. *Eur J Radiol* 2017; **89**: 140-148 [PMID: 28267531 DOI: 10.1016/j.ejrad.2017.02.008]
- 16 Hung GU, Shiau YC, Tsai SC, Ho YJ, Kao CH, Yen RF. Differentiation of radiographically indeterminate solitary pulmonary nodules with. *Jpn J Clin Oncol* 2001; **31**: 51-54 [PMID: 11302341 DOI: 10.1093/jjco/hye011]

- 17 **Cengiz A**, Aydin F, Sipahi M, Dertsiz L, Özbilim G, Bozkurt S, Güngör F, Boz A, Erkiliç M. The role of F-18 FDG PET/CT in differentiating benign from malignant pulmonary masses and accompanying lymph nodes. *Tuberk Toraks* 2018; **66**: 130-135 [PMID: 30246656 DOI: 10.5578/tt.10809]
- 18 **Huang CJ**, You DL, Lee PI, Hsu LH, Liu CC, Shih CS, Shih CC, Tseng HC. Characteristics of integrated 18F-FDG PET/CT in Pulmonary Cryptococcosis. *Acta Radiol* 2009; **50**: 374-378 [PMID: 19242830 DOI: 10.1080/02841850902756532]
- 19 **Sharma P**, Mukherjee A, Karunanithi S, Bal C, Kumar R. Potential role of 18F-FDG PET/CT in patients with fungal infections. *AJR Am J Roentgenol* 2014; **203**: 180-189 [PMID: 24951213 DOI: 10.2214/AJR.13.11712]
- 20 **Bavishi AV**, McGarry TM. A case of pulmonary cryptococcosis caused by capsule-deficient cryptococcus neoformans in an immunocompetent patient. *Respir Care* 2010; **55**: 937-941 [PMID: 20587108]
- 21 **Zhou Q**, Hu B, Shao C, Zhou C, Zhang X, Yang D, Li C. A case report of pulmonary cryptococcosis presenting as endobronchial obstruction. *J Thorac Dis* 2013; **5**: E170-E173 [PMID: 23991332 DOI: 10.3978/j.issn.2072-1439.2013.08.40]
- 22 **Marroni M**, Pericolini E, Cenci E, Bistoni F, Vecchiarelli A. Functional defect of natural immune system in an apparent immunocompetent patient with pulmonary cryptococcosis. *J Infect* 2007; **54**: e5-e8 [PMID: 16678269 DOI: 10.1016/j.jinf.2006.03.018]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoftice@wjnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjnet.com>

