

## Effect of dietary change upon the amylase and trypsin activities of the rat pancreas

BY FRANCES HOWARD AND JOHN YUDKIN

*Department of Nutrition, Queen Elizabeth College, University of London*

(Received 18 September 1962—Revised 16 January 1963)

Most of the reports on enzyme adaptation, by ourselves and other workers, have been concerned with the enzymes of micro-organisms. We have, however, also made a few observations on rats, some of which have been concerned with digestive enzymes (e.g. Lawrie & Yudkin, 1949; Roberts & Yudkin, 1961). The work to be described was carried out as a confirmation and extension of that reported by Ivy and his colleagues (Grossman, Greengard & Ivy, 1942-3, 1944). These workers measured the activity of three pancreatic enzymes: amylase, trypsin and lipase. Our own work is concerned with the first two.

Grossman *et al.* (1942-3) in their main experiment fed rats for 21 days on one of four purified diets, in which the chief differences were in the proportion of carbohydrate as starch, protein as casein and fat as lard. The control diet contained 47% starch, 18% casein and 18% fat. When the starch was increased to 67% of the diet, there was a 40% increase in the amylase activity of the pancreas and a 50% fall in the trypsin activity. When the casein was increased to 67% of the diet, there was a 20% fall in amylase activity and a threefold increase in trypsin activity. When the lard was increased to 54% of the diet, there was a 65% fall in amylase activity and no change in trypsin activity. It should perhaps be added that the four diets differed in the proportion of all three major constituents, so that, for example, the high-carbohydrate diet contained 67% starch, 15% casein and 27% lard.

In later experiments, the same workers showed that a substitution of the starch in the control diet by glucose increased amylase activity by about 20%, but did not affect trypsin activity (Grossman *et al.* 1944). A substitution of the casein by hydrolysed casein decreased trypsin activity by about 20%, but did not affect amylase activity. The daily injection of protamine zinc insulin into rats given the control diet resulted in a 30% fall in amylase activity, but did not affect trypsin activity.

### EXPERIMENTAL AND RESULTS

#### *Animals*

Male albino rats were used in all the experiments. They were weaned at 23 days, and normally housed during the experiments in large cages with grids. The exceptions were the experiments with purified diets containing starch; the rats were then housed in individual cages so that they could be given vitamin supplements separately.

### *Procedure*

Rats were transferred at weaning either directly to the experimental diet or for a few days to the stock diet. Hokin (1951) has described considerable individual variation in amylase activity of the pancreas in pigeons. We found a similar variation between rats, and at different times of the day. Much of it could have been due to variations in secretions of pancreatic juice, stimulated by ingestion of food. We attempted to reduce the variation by removing the food from the cages at 9.30 a.m. and killing the rats for enzyme measurement about 4 h later.

The rats were killed by a blow on the head, and the pancreas was removed immediately. It was washed in ice-cold distilled water, and, after as much as possible of the remaining fat and mesentery had been removed, it was blotted dry with filter-paper and weighed. It was then chopped with scissors, and transferred with 1.5 ml ice-cold water to a homogenizer consisting of a glass tube and a Perspex pestle. The homogenizer was surrounded by ice, and homogenization carried out at 2000 rev/min for 2 min. The homogenate was filtered through a square of fine gauze and the filtrate made up to 20 ml with water. Dry weight was determined on a 5 ml portion by heating to constant weight at 110°. Activation of trypsin was carried out on another 5 ml portion. The activated sample and the remainder of the homogenate were stored overnight in the refrigerator for measurement the next day of trypsin and of amylase. All measurements were carried out in duplicate.

### *Diets*

The animals had free access to the diets and to water at all times, except that the food pots were removed from the cages about 4 h before the animals were killed for measurement of enzymes. The stock diet consisted of cubes, supplemented by green vegetables and fresh milk daily, and by one drop weekly of vitamins A and D concentrate (Adexolin; Glaxo Laboratories Ltd) (see Wiesner & Yudkin, 1951). The purified diets, except those containing starch, were based on the standard purified diet of this laboratory. It consists of sucrose 60, low-vitamin casein 20, arachis oil 15, salts 5, choline 0.1, B-vitamin mixture 0.05 g, and cyanocobalamin 10 µg. The concentration of vitamins in mg/kg diet was: inositol 220, nicotinic acid 100, calcium-D-pantothenate 100, *p*-aminobenzoic acid 75, riboflavin 30, pyridoxine 8, thiamine hydrochloride 5, folic acid 1, biotin 0.2 and cyanocobalamin 0.1. In addition, the rats were given 1 mg  $\alpha$ -tocopherol on one day a week, and on another day vitamin A 120 i.u., vitamin D 20 i.u., and menaphthone 500 µg. The modification of this purified diet consisted usually in changing the quantity or type of carbohydrate and casein, the two together continuing to amount to 80 g/100 g diet.

The diets with starch contained, in 100 g, either 60 g maize starch and 20 g casein, or 30 g maize starch and 50 g casein. They were prepared by mixing together the starch and casein with 5 g salt. To this mixture were added 10 ml water at 70°; after mixing to a smooth paste, 50 g arachis oil at 70° were added. The whole was then stirred for 2–3 h at 70° and, after cooling, the solid cake was cut into cubes. These could be kept for up to 2 weeks in a closed vessel in the refrigerator. It was not found

possible to make satisfactory cubes from a diet with 20% starch; it was for this reason that our low-starch diet contained 30%. Supplements of fat-soluble vitamins were given as for the purified diets without starch. Cyanocobalamin was given in a dose of 1  $\mu$ g weekly. The remaining B vitamins were given daily as 1 ml of solution containing choline chloride 5, inositol 1.1, nicotinic acid 0.5, calcium-D-pantothenate 0.5, *p*-aminobenzoic acid 3, riboflavin 0.15, pyridoxine 0.04, thiamine hydrochloride 0.025, folic acid 0.002 and biotin 0.001 mg.

#### *Measurement of enzyme activity*

*Amylase.* The method of Smith & Roe (1949) for amylase in blood has been modified for the much higher amylase activity in our pancreatic homogenate. The method is based on the decrease in the intensity of blue colour given by the reaction of starch and iodine as the starch is hydrolysed. Instead of 1.2 g starch/100 ml, and an incubation time of 30 min, we used 2 g starch/100 ml and an incubation time of 10 min. The starch solution was prepared by making a paste of 2 g Lintner's soluble starch with a little cold water and pouring it into a boiling mixture of phosphate buffer 33 ml, N-NaCl 5.5 ml, and water 45 ml. After boiling for 3 min, the solution was cooled and made up to 100 ml. The starch solution was made afresh for each experiment, since it deteriorated even when kept overnight in the refrigerator. The phosphate buffer, pH 7.2, was made by dissolving 7.62 g anhydrous  $\text{KH}_2\text{PO}_4$  and 51.5 g  $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$  in 1 l. water. The iodine reagent contained 6 g iodine and 60 g KI in 1 l. water.

A boiling tube containing 10 ml starch solution was placed in a water-bath at 37° for about 10 min, 1 ml pancreatic homogenate suitably diluted was added, and after exactly 10 min the reaction was stopped by the addition of 2 ml N- $\text{H}_2\text{SO}_4$ . Into a 250 ml volumetric flask containing about 200 ml water were measured 5 ml N- $\text{H}_2\text{SO}_4$ , 1 ml iodine reagent and 2 ml incubation mixture, and water was then added to 250 ml. The standard was prepared by placing 5 ml starch solution and 5 ml water in a boiling tube and adding 2 ml N- $\text{H}_2\text{SO}_4$  and 1 ml pancreatic homogenate. Portions of 2 ml were taken and the colour was developed in 250 ml volumetric flasks as above. Comparisons of colour were made in a Hilger Spekker absorptiometer with a filter with maximum absorption at 620  $m\mu$  and cells of 1 cm width.

Calibration curves were made for each experiment with 5, 7 and 10 ml of starch solution. There was strict proportionality between enzyme concentration and degree of hydrolysis between 100 and 200 mg starch hydrolysed, and the initial dilution of pancreatic homogenate was made so as to give an activity within this range. The original method of Smith & Roe (1949) showed proportionality only between 0 and 40 mg starch hydrolysed.

We have used an empirical unit of amylase activity, which we define as the amount in 1 mg dry weight of tissue that will hydrolyse 20 mg starch in 10 min to a stage giving no colour with iodine as measured at 620  $m\mu$ .

*Proteolytic activity.* The method was adapted from those of a number of authors, including Kunitz (1946-7) and Gad (1948). After treatment of the pancreatic extract with enteropeptidase, total proteolytic activity at pH 8 was measured by digestion of

casein and measurement of liberated phenolic groups in the trichloroacetic-acid-soluble fraction. No attempt was made to separate the activities of trypsin and chymotrypsin; we have nevertheless followed customary usage in referring to the activity we have measured as 'trypsin'.

*Preparation of enteropeptidase.* The enzyme was prepared by the method of Hawk, Oser & Summerson (1954). The duodenal mucosa from four freshly killed pigs was scraped with a glass slide. The material was shaken with 3 vol. of acetone and left to stand for 2 h. The residue was filtered off, washed with acetone, then with a mixture of acetone and diethyl ether, and finally twice with diethyl ether. The residue was dried in air and pulverized. It was kept in the refrigerator and parts of it were taken for the whole series of experiments described in this paper.

*Activation of trypsin.* The activation was carried out with 0.1 g enteropeptidase powder, which was ground with 5 ml phosphate buffer, 0.05 M at pH 6.3. After centrifugation, 1 ml of supernatant liquid was added to 1 ml pancreatic extract in a 10 ml volumetric flask. The mixture was incubated for 15 min at 37°, and then made up to 10 ml. The activity of the sample was measured after it had been kept at 4° overnight, during which period the activity had not changed.

*Substrate.* Hammarsten's casein, 0.3 g, was suspended in 90 ml 0.1 M-phosphate buffer pH 8.04, and heated in a boiling water-bath for 15 min. The resulting solution was cooled and made up to 100 ml with buffer. It kept without deterioration for 2 weeks at 4°.

*Measurement of tryptic activity.* Casein solution, 5 ml, was placed in a boiling tube in a water-bath at 37° and 1 ml enzyme solution added. After exactly 10 min the reaction was stopped by the addition of 4 ml 10% (w/v) trichloroacetic acid. A blank solution was prepared by adding 4 ml acid to 5 ml casein solution, and then 1 ml enzyme solution. After 13 min the mixture was filtered through Whatman no. 2 paper. To 5 ml N-NaOH solution in a 25 ml flask were added 5 ml filtrate and 1 ml phenol reagent of Folin and Ciocalteu (British Drug Houses Ltd). The volume was made up to 25 ml and, after 10 min for the colour to develop, the blue colour was read in the Hilger Spekker absorptiometer with Ilford filter 700 m $\mu$  (no. 608) and cells 1 cm in width. Standard curves were produced by developing the colour of known amounts of tyrosine from 0.05 to 0.25 mg. We define a unit of tryptic activity as the amount in 1 mg dry weight of tissue which in 10 min gives a colour equal to that given by 0.05 mg pure L-tyrosine.

#### *Preliminary experiments*

*Expt 1.* From 3 to 4 days after weaning, six rats were fed for 2 weeks on the diet containing 20% sucrose and 60% casein, instead of the usual 60% sucrose and 20% casein. They showed a reduction by about one-half in amylase activity, and an increase by about one-half in trypsin activity.

*Expt 2.* Three rats were fed for 3 weeks on either of the diets used in Expt 1 (first period). Each group was then changed over to the other diet for 2 weeks (second period). At the end of this time, the enzyme activities were found to be similar to those of rats kept for the whole of the time on the diets of the second period. Thus,

the enzyme changes induced by the high- or low-sucrose diets of the first period were reversed by the changed diets during the second period.

*Expt 3.* From 4 days after weaning groups of three rats were fed for 16 days on normal or high-protein diets, but with 60 or 20% glucose or fructose instead of sucrose. Trypsin activity was the same with all these sugars. Amylase activity was also the same with all three sugars at dietary levels of 20%. At 60% sugar, fructose caused the same amylase activity as did sucrose, but glucose caused amylase activity about 50% higher than did sucrose or fructose. In other words, whereas an increase in dietary sucrose or fructose from 20 to 60% doubled amylase activity, an increase in dietary glucose by the same amount trebled amylase activity.




Fig. 1. Change with age in amylase and trypsin activities in the pancreas of rats fed on a stock diet. ●—●, amylase; ○—○, trypsin. \* See p. 283. † See p. 284.

#### *Change in enzyme activities with increasing age: stock diet*

*Expt 4.* Rats were fed from weaning on the stock diet as described earlier. Enzyme activities were measured in four to seven rats at intervals from the day of weaning at 23 days to 12 weeks of age.

It will be seen (Fig. 1) that amylase activity increased more than threefold between the ages of 4 and 7 weeks, and then did not change. Tryptic activity also increased, mostly between 4 and 6 weeks, but by only about 60%.

#### *Effect of change from high- to low-sucrose diets, or from low- to high-sucrose diets*

*Expt 5a.* Twenty-four rats were weaned on to the basal purified diet with 60% sucrose and 20% casein. After 1 week, twelve of the rats were put on to the diet with 20% sucrose and 60% casein. Enzyme measurements were carried out on about half of the animals in each group after a further 14 days and on the remainder after 22 days.

After 2 weeks, rats on the low-sucrose diet had significantly less amylase activity and significantly more trypsin activity than rats on the high-sucrose diet (Table 1). During the next week there was a significant increase in amylase activity with the 60% sucrose diet; the other changes were not significant.

*Expt 5b.* To determine how rapidly the change in enzyme activity occurs, enzyme measurements were carried out on the day the diet was changed, and 1 day and 7 or 8 days thereafter. Thirty rats were weaned on to diets containing either 60% sucrose or 20% sucrose. After 1 week, half of each group was changed over to the other diet. Owing to the inadvertent loss of some of the samples of pancreas before dry weights could be determined, all the results in this experiment have been quoted per mg wet weight of tissue.

Table 1. *Effect of change from a high-sucrose to a low-sucrose diet on amylase and trypsin activities in the rat pancreas*

(Rats fed for 1 week from weaning on diet with 60% sucrose and 20% casein. Then half of them changed to diet with 20% sucrose and 60% casein, and half kept on same diet. Days of experiment counted from change. Mean values with standard deviations)

Sucrose in diet (%)	No. of rats/group	Duration of experiment (days)	Amylase (units)*	Trypsin (units)†
60	7	14	25.5 ± 6.7	11.7 ± 2.1
20	6	14	13.8 ± 4.8	17.3 ± 3.7
60	5	22	32.9 ± 4.6	10.1 ± 0.7
20	6	22	15.0 ± 3.1	14.8 ± 2.3

\* See p. 283. † See p. 284.

It will be seen from Fig. 2 that the differences in the preliminary dietary treatment again produced differences in the amount of enzyme activity. The diet with 60% sucrose caused more amylase activity and less trypsin activity than did the diet with 20% sucrose. The curves show that the effects of this diet on both amylase and trypsin activities are reversible. The increase and decrease of trypsin activity with the decrease and increase of dietary sucrose occurred within 1 day. The corresponding changes in the amylase activity occurred more slowly, mostly after the 1st day.

#### *Changes in enzyme activity with increasing age: purified diets*

*Expt 6a.* The experiments with casein-sucrose diets having extended for no longer than 3 weeks, a similar but more prolonged experiment was undertaken. Thirty-four rats were fed from weaning on a diet with 60% sucrose and 20% casein. After 1 week, half of them were changed to the diet with 20% sucrose and 60% casein, and enzyme measurements were carried out at intervals for 8 further weeks. During this time the growth rates of the rats in the different groups were not significantly different, and were the same as of rats kept on the laboratory stock diet.

As before, there was more amylase activity when the diet contained 60% sucrose than when it contained 20% (Table 2). In this experiment, the differences after 1 week on the different diets were small. Though the amount of amylase activity continued to increase with age on both diets, the increase was much greater between

the 1st and 2nd weeks on the diet with 60% sucrose. From 2 to 8 weeks, amylase activity remained about twice that found when the diet with less sucrose was used.

Also in line with the previous results was the fact that there was much more trypsin activity in rats fed on the 60% casein diet after 1 week. Unlike with amylase, however, the trypsin activity did not increase with either diet between 2 and 8 weeks. This finding was also in contrast to that in the experiment with the stock diet, which produced an increase of about 60% in tryptic activity between 4 and 6 weeks.




Fig. 2. Change in amylase and trypsin activities in the pancreas of rats changed from high- to low-sucrose diets, or from low- to high-sucrose diets. Rats were weaned on to diets containing either 60% sucrose (60) or 20% sucrose (20). After 1 week, half of each group was changed on to the other diet (60 → 20 or 20 → 60). —, rats on 60% sucrose diet; ---, rats on 20% sucrose diet. \* See p. 283. † See p. 284.

*Expt 6b.* An experiment similar to Expt 6a was carried out at the same time, but with starch instead of sucrose. One diet contained 60% starch with 20% casein; the second diet contained 30% starch with 50% casein. This experiment lasted for 6 weeks after the rats had received for 1 week from weaning the standard diet of 60% sucrose

and 20% casein. Twenty-eight rats were used, and the rates of growth were again not significantly different from those of rats on the other diet.

The effects of the diets high and low in starch were, on the whole, the same as of those containing sucrose (Table 3). The only difference was that the effect of starch

Table 2. *Effect of age on amylase and trypsin activities in the pancreas of growing rats on a diet with 60 or 20% sucrose*

(Rats fed on diet containing 60% sucrose and 20% casein, or 20% sucrose and 60% casein, from weaning at 23 days. Mean values with standard deviations)

Weeks after weaning	No. of rats/group	Diet with 60% sucrose		Diet with 20% sucrose		
		Amylase (units)*	Trypsin (units)†	Amylase (units)*	Trypsin (units)†	
1	3	9.9 ± 0.5	8.1 ± 0.6	3	8.2 ± 0.3	15.7 ± 1.2
2	3	21.5 ± 3.8	10.4 ± 1.1	4	9.2 ± 1.2	14.7 ± 3.0
4	3	30.7 ± 3.9	10.7 ± 0.9	6	12.7 ± 1.3	13.7 ± 1.1
6	3	39.8 ± 5.1	10.7 ± 0.5	3	18.6 ± 3.2	15.9 ± 1.7
8	3	51.1 ± 6.2	9.8 ± 1.2	3	23.8 ± 4.7	15.6 ± 1.9

\* See p. 283. † See p. 284.

Table 3. *Effect of age on the amylase and trypsin activities in the pancreas of growing rats on a diet with 60 or 30% starch*

(Rats fed on diet with 60% starch and 20% casein, or with 30% starch and 50% casein, from weaning at 23 days. Mean values with standard deviations)

Weeks after weaning	No. of rats/group	Diet with 60% starch		Diet with 30% starch		
		Amylase (units)*	Trypsin (units)†	Amylase (units)*	Trypsin (units)†	
1	3	19.0 ± 3.2	9.1 ± 2.1	4	10.9 ± 1.9	18.0 ± 1.9
3	4	37.6 ± 5.2	13.4 ± 0.7	3	10.6 ± 2.8	13.5 ± 0.9
4	4	32.8 ± 4.8	10.4 ± 2.7	3	18.4 ± 3.1	14.9 ± 0.5
6	4	50.0 ± 6.3	10.4 ± 2.1	3	25.7 ± 5.2	17.9 ± 2.4

\* See p. 283. † See p. 284.

Table 4. *Effect of kind of dietary carbohydrate on amylase and trypsin activities in the rat pancreas*

(Rats fed for 1 week from weaning on diet with 60% sucrose, then on diet with 60 or 20% glucose or fructose. Mean values with standard deviations)

Days on glucose or fructose diet	Carbohydrate	Diet with 60% sugar			Diet with 20% sugar		
		No. of rats/group	Amylase (units)*	Trypsin (units)†	No. of rats/group	Amylase (units)*	Trypsin (units)†
14	Glucose	3	28.1 ± 3.0	11.9 ± 1.9	3	9.5 ± 1.8	15.0 ± 2.1
14	Fructose	—	—	—	3	10.0 ± 1.1	17.2 ± 1.8
21	Glucose	3	40.2 ± 4.7	10.5 ± 0.2	3	12.5 ± 0.6	14.6 ± 2.4
21	Fructose	3	26.3 ± 3.8	10.1 ± 1.1	3	13.3 ± 2.0	14.5 ± 1.6

\* See p. 283. † See p. 284.

on amylase activity was greater than the effect of sucrose. The fact that the diet with 30% starch caused more amylase activity than that with 20% sucrose may have been due to the difference in the amount of carbohydrate. But there was also the same effect when either starch or sucrose was present at the same level of 60% in the diet.

*Effect of glucose and fructose on enzyme activity*

*Expt 7.* Twenty-one rats were fed for 1 week from weaning on a diet with 60% sucrose. They were then transferred to one of four diets, containing 60 or 20% glucose, or 60 or 20% fructose. Enzyme activities were measured after 14 days and 21 days. As with sucrose, the level of tryptic activity was greater with 20% carbohydrate than with 60% carbohydrate (Table 4). However, no difference was found with the different sugars. Again, there was more amylase activity with the 60% level of carbohydrate than with the 20% level. At the level of 20%, the kind of sugar made no difference to amylase activity. However, after 21 days, and with the sugars at a level of 60%, fructose produced the same activity as sucrose, whereas the activity produced by glucose—40.2 units—was significantly greater ( $P < 0.01$ ) than that produced by sucrose. It was also greater than the value of 32.9 produced by sucrose after 22 days in *Expt 5a* (Table 1) or that of 30.7 after 4 weeks from weaning in *Expt 6a* (Table 2).

Table 5. *Effect of glycerol or sorbitol in diet on amylase and trypsin activities in the rat pancreas*

(Rats fed for 1 week from weaning on diet with 60% sucrose, then 1 week on diet with 10% glycerol or sorbitol, then 2 weeks on diet with 20% glycerol or sorbitol. Four rats in each group. Mean values with standard deviations)

Carbohydrate substitute	Amylase (units)*	Trypsin (units)†
Glycerol	10.8 ± 2.3	14.8 ± 2.9
Sorbitol	7.2 ± 1.2	17.4 ± 3.3

\* See p. 283. † See p. 284.

*Effect of glycerol and sorbitol on enzyme activity*

*Expt 8.* Glycerol could not be given at levels higher than 20%. Sorbitol was given for 7 days at 10% and a further 7 days at 20%; immediate introduction of 20% sorbitol into the diet of the rat causes severe diarrhoea (Morgan & Yudkin, 1957).

Four weanling rats were fed on a diet with 10% glycerol and 70% casein for 1 week, and on a diet with 20% glycerol and 60% casein for a further 2 weeks. Another four rats were given similar diets, but with sorbitol instead of glycerol. The diets with glycerol gave the same values for both enzymes as the diets with sucrose, glucose or fructose (Table 5; cf. Tables 1 and 4). The diets with sorbitol appeared to give rather less amylase activity, and rather more trypsin activity, but the differences were not significant.

*Effect of carbohydrate-free diet on enzyme activity*

Our experiments have shown that lowering the proportion of sugar from 60% of the diet to 20% decreased amylase activity; the concomitant increase of casein from 20 to 60% increased the level of trypsin. The effect was now studied of diets free from sugar, which meant also raising the level of casein to 80% of the diet.

*Expt 9.* Three days after weaning, eight rats were put on a carbohydrate-free diet with 80% casein. Enzyme activity was measured after 28 days. The amylase activity was  $5.3 \pm 0.1$  units and trypsin activity  $19.3 \pm 2.1$  units. The value for amylase was thus about 50% less, and that for trypsin about 40% more, than the values in animals in *Expt 6a* receiving the diet with 20% sucrose (see Table 2). These differences assessed by the within-experiment error variability were significant at the 1% level.

*Effect of hydrolysed casein diet on enzyme activity*

Amylase activity was about the same after rats were fed on diets with either the substrate of the enzyme, starch, or its hydrolysis product, glucose, at low or high levels. It was of interest, therefore, to see whether a tryptic digest of casein would give values for pancreatic trypsin similar to those given by casein itself.

Table 6. *Effect of hydrolysis of dietary casein on amylase and trypsin activities in the rat pancreas*

(Rats fed for 1 week from weaning on diet with 60% sucrose and 20% casein, then for 2 weeks on diet with 60% sucrose and 20% hydrolysed casein, or 20% sucrose and 60% hydrolysed casein. Three rats in each group. Mean values with standard deviations)

Hydrolysed casein in diet (%)	Amylase (units)*	Trypsin (units)†
20	$26.0 \pm 3.2$	$5.7 \pm 0.9$
60	$11.4 \pm 2.4$	$9.2 \pm 1.2$

\* See p. 283. † See p. 284.

*Expt 10.* It was possible to obtain only a small quantity of a tryptic digest of dried skim milk, with a reduced proportion of lactose. Its percentage composition, as given by the manufacturers (Benger's Laboratories Ltd) was: hydrolysed protein 70, fat 1.0, lactose 10.5, ash 8.5, moisture 10; ratio, free amino N:total amino N 60%. Two diets were constructed, containing either 20 or 60% 'protein' on the dry basis; they were thus equivalent to the casein-containing diet used in earlier experiments. There was enough of the hydrolysed casein to feed only three rats on each diet for 14 days. They were fed for 1 week after weaning on the basal diet with 20% casein before being placed on the experimental diet. The rats on the diet with 20% hydrolysed casein gained about 40 g weekly, somewhat less than the 50 g or so in the previous experiments with casein; those on the 60% hydrolysed casein diet gained only about 20 g weekly.

Amylase activity was not significantly different from that found with the casein diet

(Table 6; cf. Table 1). There was more trypsin activity with the higher level of hydrolysed casein than with the lower level, but there was less trypsin activity at both levels than found with diets containing the same amount of casein.

*Effect of change from high- to low-sucrose diets in older rats*

The experiments so far described were carried out on rats 4 weeks old at the beginning of each experiment. We now examined the effect of changing from the 60% sucrose diet to the 20% sucrose diet in animals 12–18 weeks old.

*Expt 11.* Twenty-seven rats were fed on the laboratory stock diet for from 9 to 11 weeks from weaning. They were then given the basal purified diet with 60% sucrose for 1 week. After this time, the diet of twelve rats was changed to that with 20% sucrose and 60% casein. Enzyme measurements were carried out for a further 1, 7 and 14 days.

Table 7 shows that, with 60% sucrose, there was more amylase activity in these older animals than in the younger animals, but there was no difference in trypsin activity (cf. Expts 5*a*, *b*). As before, the change from the high to the low level of sucrose led to a slow fall in amylase activity and a rapid rise in trypsin activity. The change in amylase activity was seen after 1 week and continued during the 2nd week. The change in trypsin activity was seen after 1 day.

Table 7. *Effect of change from high-sucrose to low-sucrose diet on amylase and trypsin activities in the pancreas of older rats*

(Rats fed for 9–11 weeks from weaning on laboratory stock diet, then 1 week on diet with 60% sucrose and 20% casein. Then half of them changed to diet with 20% sucrose and 60% casein, and half kept on same diet. Days of experiment counted from change. Mean values with standard deviations)

Duration of experiment (days)	Diet with 60% sucrose			Diet with 20% sucrose		
	No. of rats/group	Amylase (units)*	Trypsin (units)†	No. of rats/group	Amylase (units)*	Trypsin (units)†
0	3	54.2 ± 4.3	10.1 ± 1.5	—	—	—
1	3	51.0 ± 4.9	11.8 ± 0.5	3	60.2 ± 9.8	15.8 ± 0.2
7	3	45.0 ± 1.0	9.9 ± 1.2	3	25.0 ± 4.2	13.8 ± 1.7
14	6	45.2 ± 6.4	10.6 ± 1.0	6	16.3 ± 3.4	16.7 ± 2.3

\* See p. 283. † See p. 284.

#### DISCUSSION

The results reported here demonstrate several features, both qualitative and quantitative. They include, for example, the much greater increase with age in amylase activity than in that of trypsin, and the rapid change in trypsin activity with dietary change compared with the much slower change in amylase activity. In our present stage of knowledge, however, it seems profitless to discuss some of these findings.

Nevertheless, one feature is worth some comment, and that is the adaptive response of these enzymes to the components of the diet. The diets in our first experiments were varied by changing the proportion of sucrose and casein from the 60:20 of our

standard basal diet. It might then have been said that, for example, the effect of changing this proportion to sucrose 20 and casein 60, which decreased the level of amylase and increased the level of trypsin, could have been due either to the decrease in the proportion of sucrose, or to the increase in the proportion of casein. Our later experiments, however, revealed that by keeping the nature as well as the proportion of the protein component constant, but varying the nature of the carbohydrate, we were able to affect the amylase but not the trypsin activity. Thus, with 20% casein, tryptic activity was unaltered by substitution for sucrose of either starch, glucose, fructose, sorbitol or glycerol. But amylase activity was significantly lower with the last three than with starch or glucose. The converse was also true, so that changing the protein from casein to hydrolysed casein did not change amylase activity but reduced tryptic activity. We conclude, therefore, that the enzyme activities are affected independently, by change in the proportion and the nature of the carbohydrate or of the protein in the diet.

If we define adaptation as purposeful change in the organism, brought about by a change in the environment, then the increase or decrease in amylase or trypsin activity is adaptive in that it follows an increase or a decrease in dietary starch or dietary protein. Since, however, neither of these substances is absorbed as such, the factor which stimulates the pancreas to produce the appropriate enzyme can only be the product of hydrolysis. With amylase, we can say that our observations fit this suggestion, at least in part. Both starch and glucose, when present in large amounts in the diet, produced more amylase activity than did other substances. In our experiments, too, starch and glucose were equally effective in this respect, so that we have not been able to confirm the findings of Grossman *et al.* (1942-3, 1944) that glucose produces more amylase activity than does starch. Yet it is clear that fructose and sucrose also stimulate amylase production, since when the level of these sugars was raised from 20 to 60% in the diet, amylase activity was doubled.

One might have expected that sucrose, giving on hydrolysis an equal mixture of glucose and fructose, would cause a level of amylase activity somewhere between the levels produced by fructose and glucose. The fact that it did not do so may have something to do with the speed at which the hydrolysis products are absorbed into the blood and so reach the pancreas.

Since glucose caused as much amylase activity as the substrate starch, it seemed possible that a tryptic hydrolysate of casein would cause as much tryptic activity as the substrate casein. As we have seen, however, it did not do so. This may well have been due to the fact that the digestion of casein and other proteins *in vivo* leads to the release of amino acids in ratios which are not the same as in the final hydrolysed product. Thus the substances reaching the pancreas when unhydrolysed casein is given are not in the same proportion as those reaching it when the completely hydrolysed casein is given. If one were to state this situation teleologically, one would say that the pancreas is stimulated to produce more enzyme when it is called upon to digest more protein, and that the stimulus comes from the products of the digestion as they are produced in the body.

## SUMMARY

1. The effects of alteration in diet upon the amylase and trypsin activities of the pancreas were studied in male albino rats.

2. In rats fed on the laboratory stock diet, both amylase and trypsin activities increased with age, amylase activity increasing by more than threefold during the 2 months after weaning, and trypsin activity by about 60% in a somewhat shorter time. Rats fed on a purified diet with 60% sucrose and 20% casein showed a similar increase of about fourfold in amylase activity but a smaller and quicker increase in trypsin activity.

3. When the purified diet contained 20% sucrose and 60% casein, the amylase activity was about 50% lower, and the trypsin activity about 50% higher, than when it contained 60% sucrose and 20% casein.

4. Rats whose diets were changed from high-sucrose, low-casein to low-sucrose, high-casein showed a decrease in amylase activity and an increase in tryptic activity. Rats whose diets were changed in the opposite direction showed the opposite change in enzyme activities.

5. The change in amylase activity occurred between 1 and 7 days after the change in the dietary sucrose:casein ratio. The change in tryptic activity occurred in less than 24 h.

6. The effect was studied of diets with two levels of starch, glucose, or fructose, and one level of sorbitol or glycerol, instead of sucrose. Tryptic activity was unaffected by changes at the level of 60 or 20% in the diet. Amylase activity was also unaffected by changes at the level of 20%. At 60%, starch and glucose produced about 50% more amylase activity than did sucrose or fructose.

7. The substitution of hydrolysed casein for 60% or 20% casein in the diet did not affect amylase activity. It reduced tryptic activity by nearly half.

8. The lowest values for amylase activity were found when rats were fed on a carbohydrate-free diet with 80% casein. These were one-sixth of the highest values, found when rats were fed on a diet with 60% starch or glucose.

9. The lowest values for trypsin activity were found when rats were fed on a diet with 20% hydrolysed casein. These were one-third of the highest values, found when rats were fed on the diet with 80% casein.

10. Thus, amylase and trypsin activities are affected independently by changing the proportion or nature of the carbohydrate or protein in the diet.

We take this opportunity of recording our gratitude to Genatosan Ltd and to the Energen Foods Company for their generous support of the research programme of this laboratory, and the British Empire Cancer Campaign for a personal grant to one of us (F.H.).

## REFERENCES

- Gad, I. (1948). *Dansk Tidsskr. Farm.* **22**, 277.
- Grossman, M. I., Greengard, H. & Ivy, A. C. (1942-3). *Amer. J. Physiol.* **138**, 676.
- Grossman, M. I., Greengard, H. & Ivy, A. C. (1944). *Amer. J. Physiol.* **141**, 38.
- Hawk, P. B., Oser, B. L. & Summerson, W. H. (1954). *Practical Physiological Chemistry*, 13th ed. London: Churchill.
- Hokin, L. E. (1951). *Biochem. J.* **48**, 320.
- Kunitz, M. (1946-7). *J. gen. Physiol.* **30**, 291.
- Lawrie, N. R. & Yudkin, J. (1949). *Biochem. J.* **45**, 438.
- Morgan, T. B. & Yudkin, J. (1957). *Nature, Lond.*, **180**, 543.
- Roberts, A. H. & Yudkin, J. (1961). *Brit. J. Nutr.* **15**, 457.
- Smith, B. W. & Roe, J. H. (1949). *J. biol. Chem.* **179**, 53.
- Wiesner, B. P. & Yudkin, J. (1951). *Nature, Lond.*, **167**, 979.

*Printed in Great Britain*