

## Soybean oil and linseed oil supplementation affect profiles of ruminal microorganisms in dairy cows

S. L. Yang<sup>1ab</sup>, D. P. Bu<sup>1b</sup>, J. Q. Wang<sup>1†</sup>, Z. Y. Hu<sup>1</sup>, D. Li<sup>1</sup>, H. Y. Wei<sup>1</sup>, L. Y. Zhou<sup>1</sup> and J. J. Loo<sup>2</sup>

<sup>1</sup>State Key Laboratory of Animal Nutrition, Institute of Animal Science, Chinese Academy of Agricultural Sciences, Beijing 100193, PR China; <sup>2</sup>Department of Animal Sciences, University of Illinois, Urbana 61801, USA

(Received 11 June 2008; Accepted 11 June 2009; First published online 10 July 2009)

*The objective of this study was to evaluate changes in ruminal microorganisms and fermentation parameters due to dietary supplementation of soybean and linseed oil alone or in combination. Four dietary treatments were tested in a Latin square designed experiment using four primiparous rumen-cannulated dairy cows. Treatments were control (C, 60:40 forage to concentrate) or C with 4% soybean oil (S), 4% linseed oil (L) or 2% soybean oil plus 2% linseed oil (SL) in a 4 × 4 Latin square with four periods of 21 days. Forage and concentrate mixtures were fed at 0800 and 2000 h daily. Ruminal fluid was collected every 2 h over a 12-h period on day 19 of each experimental period and pH was measured immediately. Samples were prepared for analyses of concentrations of volatile fatty acids (VFA) by GLC and ammonia. Counts of total and individual bacterial groups (cellulolytic, proteolytic, amylolytic bacteria and total viable bacteria) were performed using the roll-tube technique, and protozoa counts were measured via microscopy in ruminal fluid collected at 0, 4 and 8 h after the morning feeding. Content of ruminal digesta was obtained via the rumen cannula before the morning feeding and used immediately for DNA extraction and quantity of specific bacterial species was obtained using real-time PCR. Ruminal pH did not differ but total VFA (110 v. 105 mmol/l) were lower (P < 0.05) with oil supplementation compared with C. Concentration of ruminal NH<sub>3</sub>-N (4.4 v. 5.6 mmol/l) was greater (P < 0.05) due to oil compared with C. Compared with C, oil supplementation resulted in lower (P < 0.05) cellulolytic bacteria (3.25 × 10<sup>8</sup> v. 4.66 × 10<sup>8</sup> colony-forming units (CFU)/ml) and protozoa (9.04 × 10<sup>4</sup> v. 12.92 × 10<sup>4</sup> cell/ml) colony counts. Proteolytic bacteria (7.01 × 10<sup>8</sup> v. 6.08 × 10<sup>8</sup> CFU/ml) counts, however, were greater in response to oil compared with C (P < 0.05). Among oil treatments, the amount of *Butyrivibrio fibrisolvens*, *Fibrobacter succinogenes* and *Ruminococcus flavefaciens* in ruminal fluid was substantially lower (P < 0.05) when L was included. Compared to C, the amount of *Ruminococcus albus* decreased by an average of 40% regardless of oil level or type. Overall, the results indicate that some ruminal microorganisms, except proteolytic bacteria, are highly susceptible to dietary unsaturated fatty acids supplementation, particularly when linolenic acid rich oils were fed. Dietary oil effects on ruminal fermentation parameters seemed associated with the profile of ruminal microorganisms.*

**Keywords:** soybean oil, linseed oil, rumen fermentation, microorganisms

### Implications

Level of dietary oil supplementation has been shown to reduce ruminal fermentation parameters in dairy cows. However, relatively little data exists pertaining to the effects of unsaturated oils on species of ruminal microorganisms. Lactating cows were fed linseed oil or soybean oil alone

(4% of dry matter (DM)) or in combination (2% of DM from each oil) to examine effects on microbial populations, bacterial species and fermentation parameters. Oils reduced cellulolytic bacteria and protozoal counts, along with total ruminal volatile fatty acids compared to controls. Furthermore, linseed oil resulted in the most pronounced decrease in amount of *Butyrivibrio fibrisolvens*, *Fibrobacter succinogenes* and *Ruminococcus flavefaciens* in ruminal fluid assessed by real-time reverse transcriptase-PCR. The negative effect of polyunsaturated fatty acids on ruminal fermentation parameters is explained at least in part by reductions in cellulolytic bacteria.

<sup>a</sup> Present address: Yunnan Provincial Laboratory of Animal Nutrition and Feed Science, Faculty of Animal Science and technology, Yunnan Agricultural University, Kunming 650201, PR China.

<sup>b</sup> These authors contribute equally.

<sup>†</sup> E-mail: wang-jia-qj@263.net

## Introduction

Supplemental oil in the diet of dairy cows changes the composition of milk fat and enhances the concentration of conjugated linoleic acid (CLA), which has been shown to have potential health benefits, including anticarcinogenic properties (Piperova *et al.*, 2000; Roche *et al.*, 2001; Loor *et al.*, 2005a). Consumption of n-3 fatty acids also has been associated with positive effects on human health. Addition of soybean oil (SBO; high linoleic acid content) to lactating dairy cow diets is expected to increase n-6 and CLA, whereas linseed oil (LSO; high linolenic acid content) would increase n-3 and CLA in milk fat (Demeyer and Doreau, 1999; Dhiman *et al.*, 2000; Chilliard *et al.*, 2001). Further, oils higher in linoleic acid than in linolenic acid were more effective in enhancing CLA in milk fat (Bu *et al.*, 2007).

In ruminants, feeding large amounts of vegetable oil has the potential to inhibit ruminal fermentation (Jenkins, 1993; Machmüller *et al.*, 1998). High dietary oil levels have been shown to decrease the counts of ruminal bacteria and protozoa (Matsumoto *et al.*, 1991; Dohme *et al.*, 2001). It was found that when feeding 2.5% fish oil to lactating dairy cows, ruminal protozoa counts were higher as compared to 5% LSO and 5% sunflower oil (Loor *et al.*, 2005b).

Negative effects on DM intake (DMI) and ruminal fermentation are often reported when unsaturated oils are fed, but some researchers have hypothesized that the negative effect of the unsaturated fatty acids on ruminal digestion would be minimized if the diet contained a higher proportion of forage due to the ability of forage to promote normal ruminal function for maximum biohydrogenation (Palmquist, 1988). As suggested previously, both increased biohydrogenation of polyunsaturated fatty acids (PUFA) as well as competition between bacteria and feed particles for adsorption of fatty acids can explain the lower toxicity of PUFA with high-forage diets. A recent study with lactating cows observed greater ruminal fiber digestibility but lower duodenal microbial-N flow when a high-forage (F : C = 65 : 35, DM basis) diet was supplemented with LSO at 3% of DM than a control (Ueda *et al.*, 2003).

Linoleic and linolenic acid are the most abundant PUFA in fresh forage or feedstuffs (e.g. cottonseeds, linseeds) typically fed to dairy cows. It is not clear whether the presence of two (linoleic acid) v. three (linolenic acid) double bonds affects the profiles of ruminal microorganisms to different extents. Maia *et al.* (2007) evaluated the effect of different fatty acids on growth of pure strains of ruminal bacteria. Linolenic acid was more toxic than linoleic acid, whereas eicosapentaenoic acid and docosahexaenoic acid were more toxic than linolenic acid. Although a few studies have examined the negative effect of unsaturated fatty acids on some ruminal microbiota (Ferlay *et al.*, 1993; Oldick and Firkins, 2000), none has evaluated relative changes in the profiles of ruminal microbiota in response to dietary PUFA. Thus, the main objective of this study was to determine the effect of SBO and LSO supplementation of a high-forage diet (60% forage) on profiles of ruminal microorganisms.

## Material and methods

### Animals and diets

Four primiparous Holstein cows equipped with a rumen cannula (# 1C; Bar Diamond, Inc., Boise, Idaho, USA) were used in a 4 × 4 Latin square design. The average body weight and days in milk at the beginning of the experiment were 521 ± 38 kg and 45 ± 10, respectively. Dietary treatments were: control diet (C), C with 4% supplemental (wt/wt basis) SBO (S), C with 4% LSO (L), and C with 2% SBO and 2% LSO (SL). The C diet contained 60% forage and 40% concentrate (Table 1). Each period was 21 days, including 14 days to allow the cows to adjust to the diet, followed by 7 days for sampling. Cows were housed in a tie-stall barn and fed individually. The total mixed ration (TMR) diets were mixed once a day and fed twice daily in equal portions at 0800 and 2000 h. The oils were stored in a cold room at 4°C and were mixed evenly with the concentrate each morning. Supplements were stored in the shade in a cool environment until fed. Orts were restricted

**Table 1** Ingredient and chemical composition of total mixed diets in cows fed a control diet or C supplemented with soybean oil, linseed oil and soybean oil plus linseed oil

Ingredients <sup>†</sup>	C	S	L	SL
	g/kg diet DM			
Alfalfa hay	100	100	100	100
Chinese wildrye (hay, ripe)	280	280	280	280
Maize silage	220	220	220	220
Maize	157	117	117	117
Wheat bran	30	20	20	20
Soybean meal	135	145	145	145
Cotton seed meal	50	50	50	50
Calcium carbonate	4	4	4	4
Calcium phosphate (dibasic)	14	14	14	14
NaCl	5	5	5	5
Premix <sup>‡</sup>	5	5	5	5
Soybean oil	0	40	0	20
Linseed oil			40	20
Nutrient composition (DM basis)				
NE <sub>L</sub> (MJ/kg DM) <sup>§</sup>	6.0	6.6	6.6	6.6
Residual OM (OM minus CP, NDF and EE)	390	360	360	360
CP	161	161	161	161
NDF	423	414	414	414
ADF	272	270	270	270
EE	26	65	65	65
Ca	9	9	9	9
Available P	6	6	6	6

DM = dry matter; OM = organic matter; EE = ether extract.

<sup>†</sup>Cows were fed a basal diet (C) or basal diet supplemented with either 4.0% soybean oil (S), 4.0% linseed oil (L) or 2.0% soybean oil + 2.0% linseed oil (SL).

<sup>‡</sup>Contained (per kilogram of DM) a minimum 1 000 000 IU of vitamin A; 65 000 IU of vitamin D; 5000 mg of vitamin E; Fe 2000 mg; Cu 1750 mg; Zn 5500 mg; Mn 2550 mg; Se 75 mg; I 70 mg; and Co 40 mg.

<sup>§</sup>The NE<sub>L</sub> values (MJ/kg DM) used for alfalfa hay, Chinese wildrye, maize silage, maize, wheat bran, soybean meal, cottonseed meal, soybean oil and linseed oil were 4.5, 4.7, 5.0, 8.1, 6.9, 9.2, 8.2, 23.7 and 23.7, respectively (China NY/t34, 2004).

to 5% to 10% of intake on an as-fed basis. Procedures used for animal care and handling were approved and conducted under established standards of the Institute of Animal Science, Chinese Academy of Agricultural Sciences, Beijing, China.

#### *Sampling and analysis*

Samples of TMR were collected daily, frozen and stored at  $-20^{\circ}\text{C}$ . Orts were sampled twice a week from each cow, composited for each treatment, frozen and stored at  $-20^{\circ}\text{C}$  until analyses were performed. Weekly representative samples of TMR and Orts from each treatment were analyzed for DM content. The DM content of feed ingredients was determined by oven-drying at  $60^{\circ}\text{C}$  for 48 h. Dried feed samples from each week during the experimental period were ground in a Cyclotec1093 mill (Tecator, Hoganas, Sweden) and analyzed for concentrations of DM, CP, ether extract, calcium and phosphorus (AOAC, 1990) and of NDF and ADF (Van Soest *et al.*, 1991). The NDF content was analyzed by the neutral detergent solution method after pretreatment with alpha-amylase. Samples were further dried at  $105^{\circ}\text{C}$  for 8 h to determine absolute DM and all chemical analyses are expressed on DM basis.

Ruminal fluid samples for volatile fatty acids (VFA),  $\text{NH}_3\text{-N}$  and pH were collected through the cannula every 2 h over 12 h on day 19 of each experimental period beginning just before the morning feeding. Whole ruminal contents were hand-collected from the anterior, dorsal and mid-ventral regions of the rumen and were squeezed through four layers of cheesecloth. The first 100 ml of strained ruminal fluid was discarded. Residual rumen fluid was immediately used to measure pH (PB-10 Standardize). Ruminal fluid samples were analyzed for  $\text{NH}_3\text{-N}$  (Broderick and Kang, 1980). The portion of ruminal fluid (20 ml) obtained for measurement of VFA was acidified with 5 ml metaphosphoric acid (25%, w/v). Acidified ruminal fluid was centrifuged at  $10\,000 \times g$  for 10 min at  $4^{\circ}\text{C}$ . The supernatant was stored at  $-20^{\circ}\text{C}$  pending determination of VFA by GLC (Model Agilent6890N; Agilent Co., Wilmington, Delaware, USA). Total and individual bacterial groups (cellulolytic, proteolytic, amylolytic bacteria and total viable bacteria) numbers were assessed using the roll-tube technique (Hungate, 1969). Protozoa counts were assessed by microscope (Olympus DP70; Olympus, Tokyo, Japan) (Boyne *et al.*, 1957; Dehority, 1984; Lu and Xie, 1991) in ruminal fluid samples collected at 0, 4 and 8 h after the morning feeding. The ruminal samples for DNA extraction were obtained via the cannula before the morning feeding. Whole ruminal contents from the anterior, dorsal and mid-ventral regions of the rumen were squeezed by hand through four layers of cheesecloth. The strained ruminal fluid was centrifuged at  $15\,000 \times g$  for 15 min at room temperature. The supernatant was discarded. The residue was immediately added to sterile normal saline and stored at  $-80^{\circ}\text{C}$ . Samples for bacterial groups and DNA extraction were taken on the same day as the samples that were collected for VFA analysis.

#### *DNA extraction*

Total DNA from the ruminal fluid was extracted as described by Tajima *et al.* (2001), with some modifications, using

a Mini Bead-Beater-1 (Biospec Products, Inc., Bartlesville Oklahoma, OK, USA) for cell lysis. Briefly, 0.5 ml ruminal fluid mixed with  $900 \mu\text{l}$  cetyl trimethyl ammonium bromide (CTAB) solution (100 mM Tris-HCl, 100 mM EDTA, 100 mM  $\text{Na}_3\text{PO}_4$ , 1.5 M NaCl, 1% CTAB, pH 8.0) in a 2 ml tube containing 1 g glass beads ( $150 \mu\text{m}$  in diameter) was shaken for 1 min at 4200 r.p.m. in a Mini Bead Beater. Tubes were then centrifuged at  $12\,000 \times g$  for 10 min. The supernatant was transferred to a fresh tube and extracted with buffered phenol, chloroform and isoamyl alcohol (volume ratio, 25:24:1). The resultant supernatant was precipitated with two volumes of cold ethanol for 1 to 2 h. After centrifugation at  $12\,000 \times g$  for 10 min at room temperature, the eluted DNA was washed with 70% ethanol, dried, dissolved in sterile Tris-EDTA (TE) buffer (1 M Tris-HCl, 0.5 M EDTA, pH 8.0), and treated with DNase-free RNase A. Samples were re-extracted with buffered phenol, precipitated with ethanol and dissolved in TE buffer. DNA was purified with the DNA fragment purification kit (TaKaRa Biotechnology Co., Ltd, Dalian, China) according to the manufacturer's recommendations. Eluted DNA was dissolved in sterile TE buffer, and stored at  $-20^{\circ}\text{C}$ . DNA concentrations were measured at 260 nm with a Beckman DU800 spectrophotometer (Beckman Co., Fullerton, CA, USA). On average, the DNA used for these experiments possessed an A260/A280 ratio of 1.8.

#### *Design and synthesis of PCR primers*

Primers designed to detect the target species are listed in Table 2. Although primers to detect the chosen bacteria are available (Asanuma *et al.*, 2001; Koike and Kobayashi, 2001), we elected to re-design primers to obtain products of 80 to 300 bp (base pairs) as typically recommended (<http://www.takara.com.cn/newproducts/015.htm>). The 16S rDNA sequences of interest were downloaded from GenBank. Sequence regions specific for a given species (with 97% similarity) were searched online against the GenBank database using the Basic Local Alignment Search Tool (BLAST) family of programs to ensure the specificity of primers. The primers satisfied requirements of species conservation within a given bacteria, and specificity from other clusters of species in the phylogenetic tree. Primer sequences were then tested against online nucleotide databases to ensure their specificity. These primers also were tested for the requirements imposed by real-time quantitative PCR (see further below). Primers were designed and sequences were edited using DNASTar 5.0 (DNASTAR Inc., Madison, WI, USA). Oligonucleotides were synthesized by Sangon (Sangon Biological Engineering Technology and Service Co., Ltd, Shanghai, China).

#### *Conventional PCR*

PCR was performed with the Takara *Taq*<sup>TM</sup> PCR kit (TaKaRa Biotechnology Co., Ltd) using PE9700 thermal cycler (PerkinElmer, Waltham, MA, USA). Amplification conditions were: one cycle at  $95^{\circ}\text{C}$  for 3 min of denaturation, 35 cycles at  $95^{\circ}\text{C}$  for 30 s, annealing at  $60^{\circ}\text{C}$  for 30 s, and extension at  $72^{\circ}\text{C}$  for 1 min. A total of  $25 \mu\text{l}$  of PCR mixture contained:

**Table 2** PCR primers and template DNA for detection of ruminal bacteria

Target bacterium (Strain)	Primer	Primer concentration ( $\mu\text{M}$ )	Purified template DNA (ng)	Product size (bp)
<i>Fibrobacter succinogenes</i> (ATCC 19169 <sup>T</sup> )	Forward 5'-GCGGGATTGAATGTACCTTGAGA-3'	10	30	204
	Reverse 5'-TCCGCCTGCCCTGAACTATC-3'			
<i>Ruminococcus flavefaciens</i>	Forward 5'-GATGCCGCGTGGAGGAAGAAG-3'	7	20	286
	Reverse 5'-CATTTCACCGCTACACCAGGAA-3'			
<i>Butyrivibrio fibrisolvens</i>	Forward 5'-TAACATGAGAGTTTGATCCTGGCTC-3'	10	25	136
	Reverse 5'-CGTACTACCCGTCGCC-3'			
<i>Ruminococcus albus</i>	Forward 5'-GTTTTAGGATTGTAACCTCTGTCTT-3'	7	25	270
	Reverse 5'-CCTAATATCTACGCATTTACCGC-3'			


2.5  $\mu\text{l}$  10 $\times$  PCR buffer (100 mM Tris-HCl (pH 8.3), 500 mM KCl, 15 mM MgCl<sub>2</sub>), 10  $\mu\text{M}$  each primer and 20 ng of purified template DNA concentration, 2  $\mu\text{l}$  dNTP mixture (each deoxynucleoside triphosphate 2.5 mM), 2.5 U *Taq*<sup>TM</sup> DNA polymerase. PCR products were separated by electrophoresis on agarose gels and stained with ethidium bromide.

#### Sequence of amplicons

The specificity of amplification was confirmed by sequencing (Sangon Biological Engineering Technology and Service Co., Ltd). After detection of species-specific PCR amplifications in total rumen DNA, the PCR products were cloned into the TA cloning kit (TaKaRa Biotechnology Co., Ltd), and the transformants were picked randomly. Recombinant plasmids were extracted using the TaKaRa MiniBEST Plasmid Purification Kit Version 2.0 (TaKaRa Biotechnology Co., Ltd). After PCR amplification with the DNA plasmid (amplification conditions were as described above), the species-specific amplification was confirmed by sequencing (Sangon Biological Engineering Technology and Service Co., Ltd). Dilutions of purified plasmid DNA from each strain were used to construct species-specific calibration curves. These calibration curves were used for calculation of the copy of targeted species in total rumen DNA. The value of the slope of the species-specific calibration curves ranged between  $-3.2$  and  $-3.6$ .

#### Real-time PCR

Quantification of each bacterial species was performed with an Applied Biosystems 7500 System Real-Time PCR (Model 7500; Applied Biosystems, Foster city, CA, USA). Optimal amplification conditions for each primer pair were achieved. The reaction mixture in 25  $\mu\text{l}$  of the final volume contained 12.5  $\mu\text{l}$  of SYBR green PCR 2 $\times$  Mastermix (OBT Instruments Co., Ltd, Beijing, China), primer and template DNA (described in Table 2). Amplification involved one cycle at 95°C for 10 min for initial denaturation, and then 40 cycles at 95°C for 15 s, followed by annealing at 60°C for 30 s, and then 1 min at 72°C. Detection of fluorescent product was set at the last step of each cycle. Specificity of amplification was performed by analysis of melting curves after each amplification. The melting curve was obtained by slow heating at 0.1°C/s increments from 65°C to 95°C, with



**Figure 1** Amplification of the targeted species in total ruminal fluid DNA. DNA size marker (2000 bp) is in the far right lane.

fluorescence recording at 0.1°C intervals. Additional specificity analyses included product size verification by gel electrophoresis of samples after the PCR run, and melting point determination analysis (Figure 1).

#### Nucleotide sequence accession number

The 16S rDNA sequences described in this paper have been deposited in the GenBank database under accession no. AY169412 (*Butyrivibrio fibrisolvens*), X85098 (*Ruminococcus albus*), X85097 (*Ruminococcus flavefaciens*) and AJ496032 (*Fibrobacter succinogenes*).

#### Data analysis

All data were analyzed as a 4 $\times$ 4 Latin square using the MIXED procedure of SAS 8.2 (SAS Institute Inc., Cary, NC, USA). The statistical model included cow as random effect, and period and treatment as fixed effects. For the statistical analysis of ruminal fluid characteristics (pH, VFA, NH<sub>3</sub>-N) and microorganism counts, sampling time and sampling time  $\times$  treatment were added to the model and analyzed using repeated measures. Orthogonal contrasts included: C v. oil-supplemented diets to test the effect of oil supplementation; S v. L to test the effect of high S v. high L intake; and SL v. S + L to test the additive effect of S and L. The significance level was declared at  $P < 0.05$  unless otherwise noted. Trends for significance were declared at  $P = 0.05$  to 0.10.

**Results**

*Chemical composition of the experimental diets*

Ingredients and chemical composition of the diets are shown in Table 1. Each treatment contained 60% forage and 40% concentrate. Oils used for diets S, L and SL were substituted for corn grain on a weight basis to achieve the desired level of supplementation. Replacing corn grain with oil supplements increased the energy content of diets in the S, L and SL diets compared with the control (6.61 v. 5.98 MJ/kg of feed DM). Because the oil content of corn grain is typically <5% of dry matter (e.g. Dhiman *et al.*, 1999; Looor *et al.*, 2002), it is unlikely that it provided a quantitatively significant amount of PUFA for biohydrogenation. Milk yield and DMI during the study averaged 14.1 ± 0.3 (*P* = 0.75) and 12.6 ± 0.6 (*P* = 0.28) kg/day and no differences (*P* > 0.05) were observed between diets.

*Ruminal fermentation parameters*

Average ruminal pH and concentrations of total VFA and NH<sub>3</sub>-N are shown in Table 3. Effect of time and time × treatment was not significant. Ruminal pH did not differ

among treatments. Ruminal NH<sub>3</sub>-N concentration was greater (4.4 v. 5.6 mmol/l), and the concentration of total VFA (110 v. 105 mmol/l) was lower due to oil supplementation. Molar proportion of acetate was not affected significantly, whereas that of propionate was significantly increased and butyrate significantly decreased for oil supplementation. Feeding the blend of S and L (SL) increased (*P* < 0.05) NH<sub>3</sub>-N concentration to a greater extent than feeding them separately.

*Microorganism counts*

Cows fed diets with supplemental oil had lower (*P* < 0.05) cellulolytic bacteria (3.25 × 10<sup>8</sup> v. 4.66 × 10<sup>8</sup> colony-forming units (CFU)/ml) and protozoa counts (9.04 × 10<sup>4</sup> v. 12.92 × 10<sup>4</sup> cells/ml), but higher proteolytic bacteria (7.01 × 10<sup>8</sup> v. 6.08 × 10<sup>8</sup> CFU/ml) counts compared with cows fed the control diet (*P* < 0.05) (Table 4). Cows fed supplemental oil had numerically lower total viable bacteria than cows fed the control diet. Counts of total viable bacteria, proteolytic bacteria, cellulolytic bacteria and protozoa in cows fed S or L were similar.

**Table 3** Ruminal pH, VFA and ammonia-N concentration in cows fed a control diet or C supplemented with soybean oil, linseed oil or soybean oil plus linseed oil

	Diet <sup>†</sup>				s.e.	Contrast		
	C	S	L	SL		C v. oil <sup>‡</sup>	S v. L	SL v. S + L
	Mean					<i>P</i> =		
pH	6.43	6.41	6.34	6.38	0.03	0.09	0.06	0.63
Total VFA (mmol/l)	110	104	105	106	4.26	*	0.81	0.61
Acetate (mol/100 mol)	69.4	70.2	69.7	69.6	0.69	0.07	0.57	0.73
Propionate (mol/100 mol)	19.5	19.6	20.4	20.2	0.28	***	***	0.15
Butyrate (mol/100 mol)	9.4	9.2	9.0	8.9	0.25	*	0.29	0.47
Valerate (mol/100 mol)	1.1	1.0	1.0	1.0	0.03	0.17	0.33	0.89
NH <sub>3</sub> -N (mmol/l)	4.4	5.5	5.5	5.8	0.09	***	0.71	*

VFA = volatile fatty acids.

<sup>†</sup>Cows were fed a basal diet (C) or basal diet supplemented with either 4.0% soybean oil (S), 4.0% linseed oil (L) or 2.0% soybean oil + 2.0% linseed oil (SL).

<sup>‡</sup>C v. oil = C versus oil (S, L, SL).

\**P* < 0.05; \*\*\**P* < 0.001.

**Table 4** Colony counts of ruminal microorganisms from cows fed a control diet and C supplemented with soybean oil, linseed oil, and soybean oil plus linseed oil

	Diet <sup>†</sup>				s.e.	Contrast		
	C	S	L	SL		C v. oil <sup>‡</sup>	S v. L	SL v. S + L
	Mean					<i>P</i> =		
Amylolytic bacteria (×10 <sup>7</sup> CFU/ml)	8.28	10.60	7.62	11.00	1.04	0.22	*	0.13
Total viable bacteria (×10 <sup>9</sup> CFU/ml)	6.19	5.89	5.06	5.69	0.77	0.52	0.50	0.84
Proteolytic bacteria (×10 <sup>8</sup> CFU/ml)	6.08	6.81	6.80	7.41	0.40	*	0.98	0.16
Cellulolytic bacteria (×10 <sup>8</sup> CFU/ml)	4.66	3.30	2.82	3.63	0.30	***	0.27	0.14
Protozoa (×10 <sup>4</sup> cells/ml)	12.92	9.92	8.17	9.04	1.11	**	0.23	0.99

CFU = colony-forming units.

<sup>†</sup>Cows were fed a basal diet (C) or basal diet supplemented with either 4.0% soybean oil (S), 4.0% linseed oil (L) or 2.0% soybean oil + 2.0% linseed oil (SL).

<sup>‡</sup>C v. oil = C versus oil (S, L, SL).

\**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001.

**Table 5** Quantification of ruminal bacteria by real time PCR in cows fed a control diet and C supplemented with soybean oil, linseed oil and soybean oil plus linseed oil

	Diet <sup>†</sup>				s.e.	Contrast		
	C	S	L	SL		C v. oil <sup>‡</sup>	S v. L	SL v. S + L
Copies per template DNA (ng)	Mean					P =		
<i>Butyrivibrio fibrisolvens</i> ×10 <sup>8</sup>	3.31	3.12	2.04	3.03	0.227	*	***	*
<i>Ruminococcus albus</i> ×10 <sup>7</sup>	9.51	5.05	5.69	5.89	0.90	***	0.59	0.61
<i>Fibrobacter succinogenes</i> ×10 <sup>5</sup>	7.95	4.83	0.48	2.64	2.06	*	0.13	0.99
<i>Ruminococcus flavefaciens</i> ×10 <sup>9</sup>	2.08	1.99	1.50	1.87	0.16	0.11	*	0.50

<sup>†</sup>Cows were fed a basal diet (C) or basal diet supplemented with either 4.0% soybean oil (S), 4.0% linseed oil (L) or 2.0% soybean oil + 2.0% linseed oil (SL).

<sup>‡</sup>C v. oil = C versus oil (S, L, SL).

\*P < 0.05; \*\*\*P < 0.001.

### Quantification of ruminal bacteria

The resulting primer set (Table 2) produced PCR products of the expected size with test strains. All PCR products were of the expected size (Figure 1). The results of quantification are shown in Table 5. Overall, the quantity of *B. fibrisolvens*, *R. albus* and *F. succinogenes* DNA decreased ( $P < 0.05$ ) by 18%, 42% and 67%, respectively, when cows were fed supplemental oil compared with control. Among oil treatments, DNA quantity of *B. fibrisolvens* and *R. flavefaciens* decreased ( $P < 0.05$ ) by 35% and 25% in cows fed L v. S.

## Discussion

### Ruminal fermentation

Oil supplementation in our study led to lower total VFA concentration. It is suggested that unsaturated FA from oil could have interfered with ruminal fermentation resulting in greater gut fill. This is supported by the observed 3% reduction in 'residual organic matter' digestion (which could be mainly starch) (Table 1), which accounted for the reduction in VFA concentration (ca. 4% reduction). Additionally, the shift in carbohydrate sources (i.e. relatively more (hemi) cellulose due to less corn grain) because of the substitution of grain for oil in the diets coupled with the reduction in cell wall fermentation due to reduced bacterial activity, partly explained the reduction in total VFA concentrations.

Dietary oil supplementation effects on ruminal NH<sub>3</sub>-N concentration have been variable. For example, in studies with sheep fed hay and concentrate, a decrease in ruminal NH<sub>3</sub>-N was observed with LSO supplementation (Ikwuegbu and Sutton, 1982; Broudiscou *et al.*, 1994). In contrast, an increase in ruminal NH<sub>3</sub>-N was observed in steers fed grass silage and concentrate (Scollan *et al.*, 2001) or fed supplemental SBO or LSO (Brokaw *et al.*, 2001; Ueda *et al.*, 2003). Our results were consistent with those found previously in bovine. In this study, protozoal counts decreased and proteolytic bacteria counts increased with supplemental oil, which likely accounted for the increase in ruminal NH<sub>3</sub>-N concentration. Supplemental oil resulted in greater proteolytic bacteria counts, possibly due to a decrease in protozoa counts and predation. The greater content of proteolytic bacteria likely accelerated protein degradation,

causing ruminal NH<sub>3</sub>-N concentration to increase. Furthermore, increased ruminal NH<sub>3</sub>-N concentration could possibly be associated with a decrease in microbial protein synthesis by ruminal microbes.

### Ruminal microorganisms

Over the years, it has been well recognized that supplemental oil affects certain aspects of microbial metabolic activity and/or modifies the microbial population in the rumen that are involved with cellulose digestion (White *et al.*, 1958; Ørskov *et al.*, 1978). Changes in the ruminal microbiota, as well as increased availability of free fatty acids in the rumen, are major factors affecting ruminal digestion. Some researchers have explained these effects based on (i) direct disruption of microbial cell membrane and cellular function by the unsaturated fatty acids, (ii) lipid coating of bacteria and feed particles, (iii) toxicity of cellulolytic bacteria by long-chain fatty acids and (iv) affecting bacterial population distribution due to antimicrobial effects of fatty acids (Jenkins, 1993). In the study of Ferlay *et al.* (1993), PUFA had a more negative effect on the metabolism of cellulolytic bacteria than saturated fatty acids. Few experiments, however, have evaluated changes in the profiles of ruminal microorganisms in response to dietary PUFA.

In the present study, cows fed supplemental oil had considerably lower numbers of cellulolytic bacteria, which likely accounted for the lower ruminal digestibility of NDF (56% v. 51%) and ADF (53% v. 50%) we observed (Hu *et al.*, 2007). Unsaturated fatty acids have been reported to have direct inhibitory effects on ruminal protozoa (Oldick and Firkins, 2000). Linoleic acid is toxic to ruminal protozoa and previous research has shown a consistent decrease in protozoal counts *in vivo* (Sutton *et al.*, 1983; Hristov *et al.*, 2004) and *in vitro* (Newbold and Chamberlain, 1988). Hristov *et al.* (2005) reported *in vitro* data showing 48%, 88% and 100% eradication of ruminal protozoa with inclusion of 0.25%, 0.5% and 1% linoleic acid, respectively, in the incubation media. Another study showed that oleic acid at 0.25%, 0.5% and 1% decreased protozoal counts by 26%, 45% and 78%, respectively (Hristov *et al.*, 2004). Further, Oldick and Firkins (2000) observed a linear decrease in ruminal protozoa with increasing degree of

unsaturation of dietary fats. In our study, a marked decrease in protozoa and cellulolytic bacteria numbers were observed when oils were supplemented. These effects were possibly associated with direct inhibition and/or coating action of the unsaturated fatty acids on microorganisms.

There is a paucity of data regarding the effects of supplemental oil on other bacterial (except for cellulolytic bacteria) populations in cattle. In the present study, total viable bacteria and amylolytic bacteria counts were not affected by oil. However, oil resulted in greater proteolytic bacteria counts possibly resulting from a decrease in protozoa as observed in cows fed L or S compared with fish oil (Loo *et al.*, 2005b).

The extreme complexity of the ruminal microbiota has been uncovered in numerous publications that employed isolation of pure cultures. However, culture-based approaches are very time- and labor-consuming, and the results, which are based on phenotypic characteristics, are not precise or conclusive. Modern molecular techniques based on sequence comparison of nucleic acids (DNA and RNA) offer certain advantages. These techniques can be used to provide molecular characterization and a classification scheme that predicts natural evolutionary relationships without cultivating the microorganisms (Mackie *et al.*, 2000), but also can be suitable for quantifying microorganisms (Reilly and Attwood, 1998). Advances in molecular microbial technique allowed us to quantify four ruminal bacteria that play key roles in aspects of ruminal biohydrogenation (e.g. lipid metabolism in ruminants). We specifically set out to design and validate PCR primers for detection of the chosen ruminal bacteria. These primers were designed to satisfy the stringent criteria required in real-time PCR quantification of a target in a microbial community DNA mixture. *Butyrivibrio fibrisolvens* is the most active bacterial species in the biohydrogenation of PUFA in the rumen (Jenkins, 1993). *Ruminococcus albus* also plays a role in the biohydrogenation of PUFA in the rumen. They remove unsaturated double bonds in order to detoxify the PUFA and enable growth of the bacterium (Devillard *et al.*, 2006).

In the present study, we observed that cows fed S or L or SL had lower DNA quantity of ruminal bacteria associated with biohydrogenation (*B. fibrisolvens* and *R. albus*) and also fibrolytic bacteria (*F. succinogenes* and *R. flavefaciens*). Kepler *et al.* (1966) reported that PUFA were particularly toxic, and other long-chain fatty acids were toxic to some of the cellulolytic bacteria also found in the rumen (Maczulak *et al.*, 1981). Furthermore, it appeared that L had a more potent inhibitory effect on ruminal microbiota. In the present study, cows fed L had lower DNA quantity of *B. fibrisolvens* and *R. flavefaciens* compared with cows fed S. Our results (except for *R. albus*) were consistent with those from a recent study by Maia *et al.* (2007), who reported that linolenic acid was more toxic than linoleic acid. In fact, the inhibitory effect of unsaturated oils on ruminal bacteria appears to be associated with the degree of unsaturation of the fatty acids, e.g. two unsaturated double bonds (linoleic acid) *v.* three double

bonds (linolenic acid). Oldick and Firkins (2000) observed a linear decrease in ruminal protozoa with an increasing degree of unsaturation of dietary fats when diets (48.5% maize silage and 51.5% concentrate) added 4.85% fat from prills of partially hydrogenated tallow, tallow or animal-vegetable fat.

## Conclusions

Similar to previous studies, we observed that oil supplementation (4% of diet DM) to dairy cows decreases ruminal fermentation leading to lower VFA concentrations. Populations of ruminal microorganisms also were affected by LSO and SBO, with total protozoa and cellulolytic bacteria being reduced, and total proteolytic bacteria being increased by oils. Our data are some of the first to use molecular techniques to quantify detrimental effects of dietary PUFA on key ruminal bacteria with major roles in fiber digestion and biohydrogenation. In this context, our results clearly showed that, at the same level of supplementation linolenic acid was more potent than linoleic acid in reducing the amount of *B. fibrisolvens*, *F. succinogenes* and *R. flavefaciens*. Detrimental effects of oils on ruminal microorganisms could partly explain the observed effects on ruminal fermentation parameters.

## Acknowledgements

This work was partially funded by the Ministry of Science and Technology and Ministry of Agricultural grants (Nos. 2006BAD12B08, 2006BAD04A08 and 2006DFB32160). The authors sincerely thank the staff of the ruminant nutrition Lab, Institute of Animal Science, Chinese Academy of Agricultural Science for all their full practical assistance and technical support. The authors also appreciate Dr Weidong Deng for providing help with bacterial cultures.

## References

- Asanuma N, Kawato M and Hino T 2001. Presence of *Butyrivibrio fibrisolvens* in the digestive tract of dogs and cats, and its contribution to butyrate production. *The Journal of General and Applied Microbiology* 47, 313–319.
- Association of Official Analytical Chemists 1990. Official methods of analysis, 14th edition. AOAC, Arlington, VA, USA.
- Boyne AW, Eadie JM and Raitt K 1957. The development and testing of a method of counting rumen ciliate protozoa. *Journal of General Microbiology* 17, 414–423.
- Broderick GA and Kang JH 1980. Automated simultaneous determination of ammonia and total amino acids in ruminal fluid and *in vitro* media. *Journal of Dairy Science* 63, 64–75.
- Brokaw L, Hess BW and Rule DC 2001. Supplemental soybean oil or corn for beef heifers grazing summer pasture: effects on forage intake, ruminal fermentation, and site and extent of digestion. *Journal of Animal Science* 79, 2704–2712.
- Broudicou L, Pochet S and Poncet C 1994. Effect of linseed oil supplementation on feed degradation and microbial synthesis in the rumen of ciliate-free and renaunated sheep. *Animal Feed Science and Technology* 49, 189–202.
- Bu DP, Wang JQ, Dhiman TR and Liu SJ 2007. Effectiveness of oils rich in linoleic and linolenic acids to enhance conjugated linoleic acid in milk from dairy cows. *Journal of Dairy Science* 90, 998–1007.
- Chilliard Y, Ferlay A and Doreau M 2001. Effect of different types of forages, animal fat or marine oils in cow's diet on milk fat secretion and composition,

- especially conjugated linoleic acid (CLA) and polyunsaturated fatty acids. *Livestock Production Science* 70, 31–48.
- China NY/t34 2004. Feeding standard of dairy cattle. China nongye hangye biao zhun/Tuijian-34. China Agricultural Publisher, Beijing, China (in Chinese).
- Dehority BA 1984. Evaluation of subsampling and fixation procedures used for counting rumen protozoa. *Applied and Environmental Microbiology* 48, 182–185.
- Demeyer D and Doreau M 1999. Targets and procedures for altering ruminant meat and milk lipids. *The Proceedings of Nutrition Society* 58, 593–607.
- Devillard E, Andant N and Wallace RJ 2006. Increased expression of a molecular chaperone GroEL in response to unsaturated fatty acids by the biohydrogenating ruminal bacterium, *Butyrivibrio fibrisolvens*. *FEMS Microbiology Letters* 262, 244–248.
- Dhiman TR, Helmink ED, McMahon DJ, Ffie RL and Pariza MW 1999. Conjugated linoleic acid content of milk and cheese from cows fed extruded oilseeds. *Journal of Dairy Science* 82, 412–419.
- Dhiman TR, Satter LD, Pariza MW, Galli MP, Albright K and Tolosa MX 2000. Conjugated linoleic acid (CLA) content of milk from cows offered diets rich in linoleic and linolenic acid. *Journal of Dairy Science* 83, 1016–1027.
- Dohme F, Machmüller A, Wasserfallen A and Kreuzer M 2001. Ruminal methanogenesis as influenced by individual fatty acids supplemented to complete ruminant diets. *Letters in Applied Microbiology* 32, 47–51.
- Ferlay A, Chabrot J, Elmeddah Y and Doreau M 1993. Ruminal lipid balance and intestinal digestion by dairy cows fed calcium salts of rapeseed oil fatty acids or rapeseed oil. *Journal of Animal Science* 71, 2237–2245.
- Hristov AN, Ivan M and McAllister TA 2004. *In vitro* effects of individual fatty acids on protozoal numbers and on fermentation products in ruminal fluid from cattle fed a high-concentrate, barley-based diet. *Journal of Animal Science* 82, 2693–2704.
- Hristov AN, Kennington LR, McGuire MA and Hunt CW 2005. Effect of diets containing linoleic acid- or oleic acid-rich oils on ruminal fermentation and nutrient digestibility, and performance and fatty acid composition of adipose and muscle tissues of finishing cattle. *Journal of Animal Science* 83, 1312–1321.
- Hu ZY, Wang JQ, Bu DP, Yang SL, Deng LF, Wei HY and Zhou LY 2007. Effect of supplementation of soybean oil and linseed oil on digestibility of organic matters and fibers in dairy cows. *China Animal Husbandry and Veterinary Medicine* 6, 5–8 (in Chinese).
- Hungate RE 1969. A roll tube method for cultivation of strict anaerobes. In *Methods in microbiology* (ed. JR Norris and DW Ribbons), vol. 3B, pp. 117–132. Academic Press, New York.
- Ikwuegbu OA and Sutton JD 1982. The effect of varying the amount of linseed oil supplementation on rumen metabolism in sheep. *The British Journal of Nutrition* 48, 365–375.
- Jenkins TC 1993. Lipid metabolism in the rumen. *Journal of Dairy Science* 76, 3851–3863.
- Kepler CR, Hirons KP, McNeill JJ and Tove SB 1966. Intermediates and products of the biohydrogenation of linoleic acid by *Butyrivibrio fibrisolvens*. *The Journal of Biological Chemistry* 241, 1350–1354.
- Koike S and Kobayashi Y 2001. Development and use of competitive PCR assays for the rumen cellulolytic bacteria: *Fibrobacter succinogenes*, *Ruminococcus albus* and *Ruminococcus flavefaciens*. *FEMS Microbiology Letters* 204, 361–366.
- Loor JJ, Ferlay A, Ollier A, Ueda K, Doreau M and Chilliard Y 2005a. High-concentrate diets and polyunsaturated oils alter *trans* and conjugated isomers in bovine rumen, blood, and milk. *Journal of Dairy Science* 88, 3986–3999.
- Loor JJ, Herbein JH and Polan CE 2002. *Trans*18:1 and 18:2 isomers in blood plasma and milk fat of grazing cows fed a grain supplement containing solvent-extracted or mechanically-extracted soybean meal. *Journal of Dairy Science* 85, 1197–1207.
- Loor JJ, Ueda K, Ferlay A, Chilliard Y and Doreau M 2005b. Intestinal flow and digestibility of *trans* fatty acids and conjugated linoleic acids (CLA) in dairy cows fed a high-concentrate diet supplemented with fish oil, linseed oil, or sunflower oil. *Animal Feed Science and Technology* 119, 203–225.
- Lu DX and Xie CW 1991. Modern research methods and technologies in ruminant nutrition. Agricultural Press, Beijing (in Chinese).
- Machmüller A, Ossowski DA, Wanner M and Kreuzer M 1998. Potential of various fatty feeds to reduce methane release from rumen fermentation *in vitro* (Rusitec). *Animal Feed Science and Technology* 71, 117–130.
- Mackie RI, Aminov RI, White BA and McSweeney CS 2000. Molecular ecology and diversity in gut microbial ecosystems. In *Ruminant physiology: digestion, metabolism, growth and reproduction* (ed. PB Cronjé), pp. 61–77. CABI Publishing, Oxford, United Kingdom.
- Maczulak AE, Dehority BA and Palmquist DL 1981. Effects of long-chain fatty acids on growth of rumen bacteria. *Applied and Environmental Microbiology* 42, 856–862.
- Maia MRG, Chaudhary LC, Figueres L and Wallace RJ 2007. Metabolism of polyunsaturated fatty acids and their toxicity to the microflora of the rumen. *Antonie van Leeuwenhoek* 91, 303–314.
- Matsumoto M, Kobayashi T, Takenaka A and Itabashi H 1991. Defaunation effects of medium-chain fatty acids and their derivatives on goat rumen protozoa. *The Journal of General and Applied Microbiology* 37, 439–445.
- Newbold CJ and Chamberlain DG 1988. Lipids as rumen-defaunating agents. *The Proceedings of the Nutrition Society* 47, 154A.
- Oldick BS and Firkins JL 2000. Effects of degree of fat saturation on fiber digestion and microbial protein synthesis when diets are fed twelve times daily. *Journal of Animal Science* 78, 2412–2420.
- Ørskov ER, Hine RS and Grubb DA 1978. The effect of urea on digestion and voluntary intake by sheep of diets supplemented with fat. *Animal Production* 27, 241–245.
- Palmquist DL 1988. The feeding value of fats. In *Feed science* (ed. ER Ørskov), pp. 293–311. Elsevier Science Publishers, Amsterdam, Netherlands.
- Piperova LS, Teter BB, Bruckental I, Sampugna J, Mills SE, Yurawecz MP, Fritsche J, Ku K and Erdman RA 2000. Mammary lipogenic enzyme activity, *trans* fatty acids and conjugated linoleic acids are altered in lactating dairy cows fed a milk fat-depressing diet. *The Journal of Nutrition* 130, 2568–2574.
- Reilly K and Attwood GT 1998. Detection of clostridium proteoclasticum and closely related strains in the rumen by competitive PCR. *Applied and Environmental Microbiology* 64, 907–913.
- Roche HM, Noone E and Gibney AN 2001. Conjugated linoleic acid: a novel therapeutic nutrient? *Nutrition Research Reviews* 14, 173–187.
- Scollan ND, Dhanoa MS, Choi NJ, Maeng WJ, Enser M and Wood JD 2001. Biohydrogenation and digestion of long chain fatty acids in steers fed on different sources of lipid. *The Journal of Agricultural Science* 136, 345–355.
- Sutton JD, Knight R, McAllan AB and Smith RH 1983. Digestion and synthesis in the rumen of sheep given diets supplemented with free and protected oils. *The British Journal of Nutrition* 49, 419–432.
- Tajima K, Aminov RI, Nagamine T, Matsui H, Nakamura M and Benno Y 2001. Diet-dependent shift in the bacterial population of the rumen revealed with real-time PCR. *Applied and Environmental Microbiology* 67, 2766–2774.
- Ueda K, Ferlay A, Chabrot J, Loor JJ, Chilliard Y and Doreau M 2003. Effect of linseed oil supplementation on ruminal digestion in dairy cows fed diets with different forage: concentrate ratios. *Journal of Dairy Science* 86, 3999–4007.
- Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3579–3583.
- White TW, Grainger RB, Baker FH and Stroud JW 1958. Effect of supplemental fat on digestion and the ruminal calcium requirement of sheep. *Journal of Animal Science* 17, 797–803.