

Placenta growth factor in sickle cell disease: association with hemolysis and inflammation

*Julia E. Brittain,¹ Ben Hulkower,² Susan K. Jones,² Dell Strayhorn,² Laura De Castro,³ Marilyn J. Telen,³ Eugene P. Orringer,² Alan Hinderliter,⁴ and *Kenneth I. Ataga²

¹Departments of Biochemistry & Biophysics and Obstetrics & Gynecology and ²Division of Hematology/Oncology, University of North Carolina at Chapel Hill;

³Division of Hematology, Duke University, Durham, NC; and ⁴Division of Cardiology, University of North Carolina at Chapel Hill

Placenta growth factor (PIGF) is released by immature erythrocytes and is elevated in sickle cell disease (SCD). Previous data generated in vitro suggest that PIGF may play a role in the pathophysiology of SCD-associated pulmonary hypertension (PHT) by inducing the release of the vasoconstrictor, endothelin-1. In this cross-sectional study of 74 patients with SCD, we confirm that PIGF is significantly el-

evated in SCD compared with healthy control subjects. We found significantly higher levels of PIGF in SCD patients with PHT but observed no association of PIGF with the frequency of acute pain episodes or history of acute chest syndrome. The observed correlation between PIGF and various measures of red cell destruction suggests that hemolysis, and the resultant erythropoietic response, results in

the up-regulation of PIGF. Although relatively specific, PIGF, as well as N-terminal pro-brain natriuretic peptide and soluble vascular cell adhesion molecule, has low predictive accuracy for the presence of PHT. Prospective studies are required to conclusively define the contribution of PIGF to the pathogenesis of PHT and other hemolytic complications in SCD. (Blood. 2010;115:2014-2020)

Introduction

Patients with sickle cell disease (SCD) exhibit elevated leukocyte counts and abnormal activation of granulocytes, monocytes, and endothelial cells.¹⁻³ They also manifest increased thrombin and fibrin generation,^{4,5} increased tissue factor procoagulant activity,^{6,7} and increased platelet activation even when they are in the noncrisis, steady state.^{5,8,9} Furthermore, increased levels of multiple inflammatory mediators are observed in patients with SCD.^{10,11} Indeed, the baseline leukocyte count is a strong independent risk factor for disease severity. Leukocytosis is a risk factor for increased mortality,¹² acute chest syndrome, hemorrhagic stroke,¹³ and vasoocclusive crises.^{14,15} As a result, SCD is increasingly referred to as a chronic inflammatory state.¹⁶

Placenta growth factor (PIGF), an angiogenic growth factor belonging to the vascular endothelial growth factor (VEGF) family,^{17,18} is produced not only by placental trophoblasts and umbilical vein endothelial cells during pregnancy but also by maturing erythroblasts.¹⁹ Plasma levels of PIGF are higher in patients with SCD than in healthy control subjects and have been reported to correlate with the frequency of acute pain episodes.²⁰ Plasma levels of PIGF may also increase during acute pain episodes in SCD.²¹ The higher PIGF levels in patients with SCD may be due to hypoxia,²² increased erythropoiesis,¹⁹ and increased erythropoietin concentrations that follow anemia in these patients.²⁰ Although PIGF binds to the VEGF-1 class of receptors,^{23,24} this cytokine elicits its own unique proinflammatory and arteriogenic effects.²⁵ Perelman et al²⁰ have demonstrated *ex vivo* that PIGF activates monocytes and promotes the release of interleukin-1 β , interleukin-8, monocyte chemoattractant protein-1, and VEGF from these

cells. PIGF also induces leukotriene formation in SCD.²⁶ These data suggest a clinical role for PIGF in inflammation in SCD.

PIGF may play a role in pulmonary hypertension (PHT) in SCD. Data generated in vitro suggest that PIGF induces the release of the vasoconstrictor, endothelin-1 (ET-1) from pulmonary microvascular endothelial cells.²⁷ In addition, treatment of these cells with PIGF induced expression of the endothelin B receptor, suggesting that PIGF may contribute to the pathogenesis of SCD-associated PHT. Furthermore, the arteriogenic effects of PIGF suggests a role for this cytokine in the plexiform lesions noted in PHT, the formation of which may be monocyte-dependent.^{25,28}

In this study, we sought to evaluate the association of PIGF with measures of both inflammation and hemolysis in patients with SCD and to examine the association of PIGF with specific clinical complications in a cohort of patients followed at an adult sickle cell clinic.

Methods

Patients and study design

The study patients represent a cohort followed at the Sickle Cell Clinic at the University of North Carolina, Chapel Hill. Consecutive patients seen in the clinic for routine follow-up were evaluated. Seventy-four patients with SCD and an additional 19 healthy, race-matched control subjects were included in the analyses. Patients were assessed while in the noncrisis, steady state; had not experienced an episode of acute chest syndrome in the 4 weeks preceding enrollment; and had no clinical evidence of congestive

Submitted April 29, 2009; accepted December 1, 2009. Prepublished online as *Blood* First Edition paper, December 29, 2009; DOI 10.1182/blood-2009-04-217950.

*J.E.B. and K.I.A. contributed equally to this manuscript.

Parts of this work were presented in abstract form at the 50th annual meeting of the American Society of Hematology, San Francisco, CA, December 7, 2008.⁴³

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2010 by The American Society of Hematology

heart failure. None of the study patients were on chronic red blood cell transfusion. This study was approved by the Institutional Review Board at University of North Carolina, Chapel Hill, and all subjects gave written informed consent to participate in accordance with the Declaration of Helsinki.

Echocardiography and PHT determination

Transthoracic Doppler echocardiography was performed in all study subjects with the use of a Philips Sonos 5500 ultrasound system as previously described.²⁹ All the echocardiograms were interpreted by a single cardiologist blinded to all patient data. The pulmonary artery systolic pressure (PASP) was calculated using the modified Bernoulli equation ($PASP = 4V^2 + \text{right atrial pressure}$), with the right atrial pressure assumed to be 10 mm Hg. Patients with no detectable tricuspid regurgitant jet were assumed to have a normal PASP if they had no other findings suggestive of PHT and were assigned a tricuspid regurgitant jet velocity that was lower than any measured in the study (1.8 m/s). The diagnosis of PHT in our study was based on PASP values adjusted for age, sex, and body mass index.³⁰

SCD-related clinical complications

The presence (or history) of clinical complications was ascertained from a history obtained at the time of evaluation, combined with a detailed review of the patients' medical records. Acute pain episodes (or pain crises), acute chest syndrome, stroke, and other complications were defined with the use of standard definitions in patients with SCD.^{13,31-33}

Measurement of laboratory variables

Blood samples were obtained by venipuncture and drawn into citrate-containing tubes. To minimize any artifactual platelet activation and release during processing, 1 μ M indomethacin and 0.05 μ g/mL prostaglandin I₂ (final concentration) were mixed gently with citrated whole blood. The sample was incubated for 15 minutes at 37°C and centrifuged at 150g for 15 minutes. Indomethacin and prostaglandin I₂ were added to the extracted plasma sample, followed by centrifugation at 400g for 15 minutes to prepare platelet-poor plasma. The plasma samples were then divided into aliquots and frozen immediately at -80°C for subsequent analysis. Quantification of PIGF, ET-1, and human soluble vascular cell adhesion molecule-1 (VCAM) was accomplished with the use of commercially

available enzyme-linked immunoabsorbent assay kits from R&D Systems. Samples were assayed in duplicate and according to manufacturer's instructions. Measurements of N-terminal pro-brain natriuretic peptide (NT-proBNP) and other routine laboratory studies were performed by the McClendon Clinical Laboratory at the University of North Carolina Hospitals. Values of NT-proBNP that were below the measurable limit (< 50 pg/dL) were assigned a value of 49 pg/dL.

Statistical analysis

Nonparametric Mann-Whitney 2-sided *t* tests were used to compare continuous variables, and categorical variables were compared with the use of a χ^2 test. Spearman correlations were used to identify associations between PIGF and specified variables ($\alpha = 0.05$). Bonferroni adjustments were made for multiple comparisons (PIGF and measures of hemolysis, such as lactate dehydrogenase level, reticulocyte count, and hemoglobin level; and PIGF and measures of inflammation such as soluble VCAM, absolute neutrophil count, and absolute monocyte count). Logistic regression analysis was used to explore whether the observed bivariable relationship between PIGF and PHT might be confounded by other specified variables (ie, reticulocyte count, lactate dehydrogenase level, and hemoglobin level). Receiver-operator characteristic curves (ROCs) for PIGF, NT-proBNP, soluble VCAM, and PHT were generated with the use of Analyze It for Microsoft Excel. Other statistical analyses were conducted by Prism software (Version 5.00 for Windows; GraphPad Software Inc).

Results

Demographic and laboratory characteristics

The demographic and laboratory characteristics of all the study subjects are shown in Table 1. Sixty-two patients (83.7%) had sickle cell anemia (HbSS), 8 had HbSC disease (10.8%), 3 had sickle β^0 thalassemia (4.1%), and 1 had sickle β^+ thalassemia (1.4%). All the healthy control subjects (18 HbAA and 1 HbAC) had normal PASP values. As expected, patients with SCD had higher white blood cell counts ($9.4 \times 10^9/L$ vs $7.2 \times 10^9/L$; $P = .001$), platelet counts ($426 \times 10^9/L$ vs $284 \times 10^9/L$; $P < .001$),

Table 1. Demographic and laboratory characteristics of study subjects

Characteristic	SCD (n = 74)	Control (n = 19)	P
Median age, y (IQR)	41 (30-48.3)	33 (22-50)	NS
Female sex, n (%)	47 (64)	15 (79)	NS
Genotype, n			—
SS	62	—	
SC	8	—	
S β^0	3	—	
S β^+	1	—	
PASP, mm Hg, median (IQR)	31 (23-39)	26 (23-28)	< .001
TRjet, m/s, median (IQR)	2.3 (1.8-2.7)	2.0 (1.8-2.1)	.009
White blood cell count, $\times 10^9/L$, median (IQR)	9.4 (7.4-11.5)	7.2 (6.1-9.7)	.001
Hemoglobin level, g/dL, median (IQR)	8.9 (7.6-10.2)	12.4 (11.9-13.5)	< .001
Platelet count, $\times 10^9/L$, median (IQR)	426 (325.5-512)	284 (265-322)	< .001
Reticulocyte count, %, median (IQR percentile)	6.9 (4.5-10.5)	1.5 (1.3-1.9)	< .001
Absolute neutrophil count, $\times 10^9/L$, median (IQR)	5.1 (4-6.2)	4.7 (4-6.5)	NS
Absolute monocyte count, $\times 10^9/L$, median (IQR)	0.5 (0.3-0.7)	0.4 (0.3-0.4)	< .001
Hydroxyurea use, n (%)	39 (53)	—	—
NT-proBNP level, pg/dL, median (IQR)	133.5 (60.3-326)	49 (49-59)	< .001
Creatinine level, mg/dL, median (IQR)	0.75 (0.6-1)	0.8 (0.7-0.9)	NS
Total bilirubin level, mg/dL, median (IQR)	2.2 (1.2-2.9)	0.3 (0.2-0.4)	< .001
Direct bilirubin level, mg/dL, median (IQR)	0.09 (0.09-0.1)	0.09 (0.09-0.09)	NS
LDH level, U/L, median (IQR)	898 (658-1199)	488 (413-516)	< .001
PIGF level, pg/mL, median (IQR)	9.03 (6-11.4)	0.23 (0-6.8)	< .001

SCD indicates sickle cell disease; IQR, interquartile range; NS, not significant; PASP, pulmonary arterial systolic pressure; TRjet, tricuspid regurgitant jet velocity; NT-proBNP, N terminal pro-brain natriuretic peptide; LDH, lactate dehydrogenase; and PIGF, placenta growth factor.

and reticulocyte counts (6.9% vs 1.5%; $P < .001$) than did healthy control subjects. In addition, patients with SCD had significantly lower hemoglobin levels (8.9 g/dL vs 12.4 g/dL; $P < .001$) and higher total bilirubin (2.2 mg/dL vs 0.3 mg/dL; $P < .001$), lactate dehydrogenase (898 U/L vs 488 U/L; $P < .001$), and NT-proBNP (133.5 pg/mL vs 49 pg/mL; $P < .001$) levels than did healthy controls. Plasma levels of PIGF (9.03 pg/mL vs 0.23 pg/mL; $P < .001$) were significantly higher in patients with SCD than in healthy control subjects (Figure 1).

Association of PIGF with SCD-related clinical complications

We found no significant correlation between PIGF levels and the number of acute pain episodes in the past year in patients with SCD (Figure 2). However, we observed that plasma levels of PIGF were significantly higher in patients with PHT than in patients without PHT (Figure 3; Table 2). Median plasma levels of PIGF also appeared to be higher in patients with a history of thrombotic stroke than in patients without this complication, although the difference did not achieve statistical significance (11.0 pg/mL vs 8.8 pg/mL; $P = .08$). No differences were observed in PIGF levels when patients with a history of acute chest syndrome, leg ulcers, priapism, or retinopathy were compared with patients without these complications (data not shown). In addition, there was no significant difference in PIGF levels in patients on hydroxyurea compared with patients not on hydroxyurea. Although adherence to hydroxyurea was not evaluated in this study, patients identified as being on hydroxyurea had significantly higher mean corpuscular volume values (104 fL; interquartile range [IQR], 99-110 fL vs 88 fL; IQR, 79-96 fL; $P < .001$) and HbF levels (10.4%; IQR, 5.2%-17.5% vs 4.6%; IQR, 1.7%-6.1%; $P = .002$) compared with patients not on hydroxyurea. There were, however, no significant differences in white blood cell counts and hemoglobin levels in patients receiving hydroxyurea compared with patients not on hydroxyurea.

Correlation of PIGF and laboratory variables

Although PIGF has been reported to induce ET-1 release in vitro,²⁷ we found no correlation between PIGF and ET-1 ($r = -0.136$, $P = .3$). However, we observed significant correlations between PIGF levels and both lactate dehydrogenase level ($r = 0.36$, $P = .002$; Figure 4A) and reticulocyte count ($r = 0.32$, $P = .007$; Figure 4B) and a negative correlation between PIGF and hemoglobin levels ($r = -0.44$, $P < .001$; Figure 4C). In addition, we observed a correlation between PIGF and soluble VCAM levels ($r = 0.42$, $P < .001$; Figure 4D; Table 3).




Figure 1. PIGF levels in patients with SCD and healthy controls. Plasma levels of PIGF in patients with SCD were significantly higher than in healthy control subjects (9.03 pg/mL vs 0.23 pg/mL; $P < .001$). Data are shown as median with range.




Figure 2. PIGF is not associated with the frequency of acute pain episodes. Data are shown as median with range and stratified according to disease severity (mild vs severe).

Association of ET-1 with PHT

There was a trend toward a higher level of ET-1 in patients with PHT than in patients without this complication (0.07 pg/mL vs 0.057 pg/mL, $P = .07$). Although there was a weak correlation between ET-1 and NT-proBNP ($r = 0.102$, $P = .015$), we did not observe any association between ET-1 and hemoglobin level, lactate dehydrogenase level, or reticulocyte count.

Predictive accuracy of PIGF for PHT

When the PASP was adjusted for age, sex, and body mass index, the area under the ROC with the use of PIGF to diagnose PHT was 0.71 (Figure 5). We observed a correlation of PIGF with NT-proBNP ($r = 0.25$, $P = .03$), a laboratory marker of cardiac dysfunction and an established predictor of PHT in SCD.³⁴ The area under the ROC with the use of soluble VCAM and NT-proBNP to diagnose PHT was 0.77 for each biomarker (Figure 5). In addition, there was a significant correlation between soluble VCAM and NT-proBNP ($r = 0.46$, $P < .001$). Using the 75th percentile for each variable as the threshold level, we observed no substantial difference in the predictive accuracies of the 3 biomarkers for the diagnosis of PHT in our cohort (Table 4). The sensitivity of PIGF for the presence of PHT was 53%, compared with sensitivities of 60% for NT-proBNP and 50% for soluble VCAM; the specificity of PIGF was 83%, compared with 87% for NT-proBNP and 85% for soluble VCAM; and the likelihood ratio for PIGF was 3.18, compared with 4.63 for NT-proBNP and 3.31 for soluble VCAM.




Figure 3. PIGF levels in PHT in SCD. PIGF is elevated in PHT (11.5 pg/mL vs 8.5 pg/mL; $P = .009$). Data are shown as median with range.

Table 2. Measures of hemolysis and inflammation in PHT and stroke

Variable*	PHT (n = 20)	No PHT (n = 54)	P	Stroke (n = 11)	No stroke (n = 62)	P
Hb level, g/dL	7.6 (6.3-8.8)	9.4 (8.3-10.2)	< .001	7.5 (6.9-9.4)	9 (7.8-10.2)	.03
LDH level, U/L	1077 (879-1364)	808 (620-1079)	.03	1136 (947-1367)	840 (646-1123)	.05
sVCAM level, ng/mL	1123 (758-1346)	624 (486-947)	< .001	1135 (813-1563)	753 (515-1044)	.05
PlGF level, pg/mL	11.5 (8.8-15.4)	8.5 (5.7-10.8)	.009	11.0 (8.4-16.0)	8.8 (6.0-11.3)	.08

PHT indicates pulmonary hypertension; Hb, hemoglobin; LDH, lactate dehydrogenase; sVCAM, soluble vascular cell adhesion molecule-1; and PlGF, placenta growth factor.

*Data are presented as medians and interquartile ranges (25th-75th percentiles).

Logistic regression analysis

To assess whether PlGF is an independent predictor of PHT in SCD, we used logistic regression analyses to explore whether controlling for reticulocyte count, lactate dehydrogenase, and hemoglobin, separately, affected the observed association between PlGF and PHT. Because of the small number of participants with PHT in our cohort, it is not possible to adequately control for potential confounding variables, especially more than one at a time. Controlling for either reticulocyte count or lactate dehydrogenase had little effect on the magnitude or significance of the estimated odds ratio associated with a one-unit change in PlGF (Table 5). However, when we controlled for hemoglobin, the estimate of the odds ratio for PlGF was closer to one and was no longer statistically significant ($P = .43$).

Discussion

The elevated levels of PlGF in our study patients compared with healthy control subjects further confirms the inflammatory

nature of SCD. Despite the abundant evidence of inflammation in patients with SCD, its contribution to the pathophysiology of specific complications remains poorly defined. The correlation of PlGF with both lactate dehydrogenase levels and reticulocyte count, as well as its inverse correlation with hemoglobin levels, suggests that hemolysis plays a role in the up-regulation of PlGF (and inflammation) in patients with SCD. Because hemolysis results in increased erythropoietin production and erythroid hyperplasia, our finding is consistent with the previous report of increased PlGF expression in erythroid cells after the addition of erythropoietin.²⁰

Our finding that PlGF is elevated in patients with SCD with PHT and possibly in thrombotic stroke, but not with such vasoocclusive complications such as acute chest syndrome or acute pain episodes, is contrary to a previous report that suggested a correlation between PlGF levels and the frequency of acute pain episodes.²⁰ In our study, PHT and history of stroke do not appear to be independent (data not shown), although any inference of dependence is limited by the relatively small number of subjects with




Figure 4. Correlation between PlGF and markers of hemolysis and inflammation. (A) PlGF is correlated with lactate dehydrogenase levels ($r = 0.36$, $P = .002$). (B) PlGF is correlated with reticulocyte count ($r = 0.32$, $P = .007$). (C) PlGF is inversely correlated with hemoglobin levels ($r = -0.44$, $P = .001$). (D) PlGF is correlated with soluble VCAM ($r = 0.42$, $P < .001$). Data shown with regression line. Spearman correlation data are shown in Table 2.

Table 3. Association of PIGF and measures of hemolysis and inflammation

Variable	r	P
Hemoglobin level	-0.44	< .001
Reticulocyte count	0.32	.007
Lactate dehydrogenase level	0.36	.002
sVCAM level	0.42	< .001
Absolute neutrophil count	0.18	.13
Absolute monocyte count	0.18	.14

PIGF indicates placenta growth factor; and sVCAM, soluble vascular cell adhesion molecule-1.

these events in this study cohort. Every single one of the patients with a history of stroke also had PHT, and 55% of those patients with PHT had a history of stroke. There is an increasing body of evidence that hemolysis is associated with complications such as PHT, leg ulcers, priapism, and possibly stroke but not with acute pain episodes or acute chest syndrome. Hemolysis may predispose to certain SCD complications by decreasing the bioavailability of nitric oxide.³⁵ This reduced bioavailability is thought to impair downstream effects of nitric oxide, such as inhibition of platelet activation,³⁶ coagulation activation,²⁹ and transcriptional repression of cell adhesion molecules, VCAM-1, intracellular adhesion molecule-1, P-selectin, and E-selectin.^{29,37} The correlation of PIGF with soluble VCAM suggests that PIGF may also contribute to the endothelial activation observed in patients with SCD.

The finding that PIGF is independent of either lactate dehydrogenase level or reticulocyte count as a predictor of PHT, but not independent of hemoglobin level, suggests that PIGF may predispose to PHT by a mechanism independent of hemolysis. The dependence of PIGF on hemoglobin level in predicting for PHT may be related to the decreased oxygen-carrying capacity associated with anemia. In view of our relatively small sample size, more studies are needed to fully validate the role of PIGF in the pathogenesis of SCD-associated PHT.

Despite the observed correlations, our data do not provide a causal link between PIGF and either PHT or stroke in SCD. It is particularly interesting that no significant correlation was observed between PIGF and ET-1. Markedly elevated PIGF levels during pregnancy have been reported to be associated with a decline in the level of ET-1 in the first trimester of pregnancy, and these levels remain low throughout gestation.³⁸ Although there is a suggestion that ET-1 is associated with PHT in our study, elevated PIGF levels may produce a vasculopathy by an inflammatory mechanism not dependent on ET-1.

Although the observed elevation of PIGF levels in our study patients with PHT and stroke are relatively modest, being approximately 1.4- and 1.2-fold, respectively, compared with SCD patients without these complications, the presence of chronic, low-level elevation of this cytokine may be sufficient to produce vasculopathic changes in SCD patients, possibly in combination with increased levels of other angiogenic cytokines such as VEGF,




Figure 5. Potential predictive accuracy of biomarkers in PHT. ROC curves of PIGF, soluble VCAM, and NT-proBNP in patients with SCD.

erythropoietin, and angiopoietin.²¹ Elevated PIGF levels may also play a role in the formation of fragile, tortuous blood vessels in the brain (Moyamoya), a risk factor for stroke.³⁹ Furthermore, PIGF-induced remodeling of the pulmonary vasculature may play a role in the formation of the plexiform lesions noted in PHT.⁴⁰

Doppler echocardiography is usually used to screen for PHT. However, only approximately 80% to 90% of patients with increased right ventricular systolic pressures have a Doppler tricuspid regurgitant signal that is sufficient to predict pulmonary artery systolic pressure.⁴¹ The availability of biomarkers that predict the presence of PHT may be complementary to Doppler echocardiography in the diagnosis of this life-threatening complication in SCD. Furthermore, these biomarkers may provide prognostic information, as has been shown for NT-proBNP.³⁴ Our current data show that, although relatively specific, PIGF, NT-proBNP, and soluble VCAM have low predictive accuracies for the presence of PHT, with low sensitivities and positive predictive values. The predictive accuracies of currently available biomarkers may however be enhanced by using statistical modeling techniques that combine biomarkers as well as other types of measurements.

In view of the potential role of hemolysis in the up-regulation of PIGF, therapeutic strategies that reduce red cell destruction or increase the oxygen-carrying capacity of the blood, including chronic red blood cell transfusion or treatment with compounds such as the investigational Gardos channel blocker, senicapoc,⁴² may be beneficial in decreasing levels of PIGF. It is interesting that no differences in PIGF levels were observed in patients on hydroxyurea compared with those patients not on hydroxyurea. Despite the increased mean corpuscular volume and HbF values in

Table 4. Summary of ROC curves for PIGF, NT-proBNP, and sVCAM

Variable	AUC	95% CI	75th percentile	Sensitivity, %	Specificity, %	Likelihood ratio	Positive predictive value, %
PIGF, pg/mL	0.71	0.57-0.85	11.5	53	83	3.18	59
NT-proBNP, pg/mL	0.77	0.63-0.91	326	60	87	4.63	60
sVCAM, ng/mL	0.77	0.65-0.88	1199	50	85	3.31	50

ROC indicates receiver-operator characteristic curve; PIGF, placenta growth factor; NT-proBNP, N-terminal pro-brain natriuretic peptide; sVCAM, soluble vascular cell adhesion molecule-1; and AUC, area under the curve.

Table 5. Estimates of the unadjusted and adjusted odds ratio for PIGF and PHT

Markers of hemolysis	Odds ratio estimate for PIGF in PHT	95% CI	P
None	1.19	(1.04-1.35)	.01
Lactate dehydrogenase level	1.17	(1.02-1.34)	.03
Reticulocyte count	1.19	(1.04-1.37)	.01
Hemoglobin level	1.06	(0.92-1.23)	.4

PIGF indicates placenta growth factor; and PHT, pulmonary hypertension.

patients on hydroxyurea, there were no significant differences in hemoglobin levels between those patients on hydroxyurea and those patients not on the drug, potentially accounting for the similarities in PIGF levels between the 2 groups.

Because our study had a cross-sectional design, longer term prospective studies are required to validate the clinical significance of the modest elevation of PIGF levels observed in SCD-associated PHT and to further define the role of PIGF in SCD.

In conclusion, PIGF is higher in patients with SCD than in healthy control subjects and is correlated with the presence of PHT. The association of PIGF with measures of hemolysis suggests that treatments that improve survival of red blood cells may be beneficial in the treatment or prevention of SCD-associated vasculopathy. Prospective studies evaluating PIGF are required to conclusively define its contribution to the pathogenesis of SCD.

References

- Okpala I. The intriguing contribution of white blood cells to sickle cell disease - a red cell disorder. *Blood Rev.* 2004;18(1):65-73.
- Wun T. The role of inflammation and leukocytes in the pathogenesis of sickle cell disease; haemoglobinopathy. *Hematology.* 2001;5(5):403-412.
- Hebbel RP, Osarogigabon R, Kaul D. The endothelial biology of sickle cell disease: inflammation and a chronic vasculopathy. *Microcirculation.* 2004;11(2):129-151.
- Ataga KI, Orringer EP. Hypercoagulability in sickle cell disease: a curious paradox. *Am J Med.* 2003;115(9):721-728.
- Francis RB Jr. Platelets, coagulation, and fibrinolysis in sickle cell disease: their possible role in vascular occlusion. *Blood Coagul Fibrinolysis.* 1991;2(2):341-353.
- Key NS, Slungaard A, Dandele L, et al. Whole blood tissue factor procoagulant activity is elevated in patients with sickle cell disease. *Blood.* 1998;91(11):4216-4223.
- Shet AS, Aras O, Gupta K, et al. Sickle blood contains tissue factor-positive microparticles derived from endothelial cells and monocytes. *Blood.* 2003;102(7):2678-2683.
- Lee SP, Ataga KI, Zayed M, Manganello JM, et al. Phase I study of eptifibatid in patients with sickle cell anaemia. *Br J Haematol.* 2007;139(4):612-620.
- Lee SP, Ataga KI, Orringer EP, Phillips DR, Parise LV. Biologically active CD40 ligand is elevated in sickle cell anemia: potential role for platelet-mediated inflammation. *Arterioscler Thromb Vasc Biol.* 2006;26(7):1626-1631.
- Brittain JE, Parise LV. Cytokines and plasma factors in sickle cell disease. *Curr Opin Hematol.* 2007;14(5):438-443.
- Pathare A, Kindi SA, Daar S, Dennison D. Cytokines in sickle cell disease. *Hematology.* 2003;8(5):329-337.
- Platt OS, Brambilla DJ, Rosse WF, et al. Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N Engl J Med.* 1994;330(23):1639-1644.
- Ohene-Frempong K, Weiner SJ, Sleeper LA, et al. Cerebrovascular accidents in sickle cell disease: rates and risk factors. *Blood.* 1998;91(1):288-294.
- Frenette PS. Sickle cell vasoocclusion: heterotypic, multicellular aggregations driven by leukocyte adhesion. *Microcirculation.* 2004;11(2):167-177.
- Turhan A, Weiss LA, Mohandas N, Collier BS, Frenette PS. Primary role for adherent leukocytes in sickle cell vascular occlusion: a new paradigm. *Proc Natl Acad Sci U S A.* 2002;99(5):3047-3051.
- Platt OS. Sickle cell anemia as an inflammatory disease. *J Clin Invest.* 2000;106(3):337-338.
- Athanassiades A, Lala PK. Role of placenta growth factor (PIGF) in human extravillous trophoblast proliferation, migration and invasiveness. *Placenta.* 1998;19(7):465-473.
- Persico MG, Vincenti V, DiPalma T. Structure, expression and receptor-binding properties of placenta growth factor (PIGF). *Curr Top Microbiol Immunol.* 1999;237:31-40.
- Tordjman R, Delaire S, Plouet J, et al. Erythroblasts are a source of angiogenic factors. *Blood.* 2001;97(7):1968-1974.
- Perelman N, Selvaraj SK, Batra S, et al. Placenta growth factor activates monocytes and correlates with sickle cell disease severity. *Blood.* 2003;102(4):1506-1514.
- Duits AJ, Rodriguez T, Schnog JJ. Serum levels of angiogenic factors indicate a pro-angiogenic state in adults with sickle cell disease. *Br J Haematol.* 2006;134(1):116-119.
- Torry RJ, Tomanek RJ, Zheng W, Miller SJ, Labarrere CA, Torry DS. Hypoxia increases placenta growth factor expression in human myocardium and cultured neonatal rat cardiomyocytes. *J Heart Lung Transplant.* 2009;28(2):183-190.
- Terman B, Khandke L, Dougher-Vermazan M, et al. VEGF receptor subtypes KDR and FLT1 show different sensitivities to heparin and placenta growth factor. *Growth Factors.* 1994;11(3):187-195.
- Sawano A, Takahashi T, Yamaguchi S, Aonuma M, Shibuya M. Flt-1 but not KDR/Flk-1 tyrosine kinase is a receptor for placenta growth factor, which is related to vascular endothelial growth factor. *Cell Growth Differ.* 1996;7(2):213-221.
- Pipp F, Heil M, Issbrucker K, et al. VEGFR-1-selective VEGF homologue PIGF is arteriogenic: evidence for a monocyte-mediated mechanism. *Circ Res.* 2003;92(4):378-385.
- Patel N, Gonsalves CS, Yang M, Malik P, Kalra VK. Placenta growth factor induces 5-lipoxygenase-activating protein to increase leukotriene formation in sickle cell disease. *Blood.* 2009;113(5):1129-1138.
- Patel N, Gonsalves CS, Malik P, Kalra VK. Placenta growth factor augments endothelin-1 and endothelin-B receptor expression via hypoxia-inducible factor-1 alpha. *Blood.* 2008;112(3):856-865.
- Moldovan NI. Role of monocytes and macrophages in adult angiogenesis: a light at the tunnel's end. *J Hematother Stem Cell Res.* 2002;11(2):179-194.
- Ataga KI, Moore CG, Hillery CA, et al. Coagulation activation and inflammation in sickle cell disease-associated pulmonary hypertension. *Haematologica.* 2008;93(1):20-26.
- McQuillan BM, Picard MH, Leavitt M, Weyman AE. Clinical correlates and reference intervals for pulmonary artery systolic pressure among echocardiographically normal subjects. *Circulation.* 2001;104(23):2797-2802.
- Charache S, Terrin ML, Moore RD, et al. Effect of hydroxyurea on the frequency of painful crises in sickle cell anemia. Investigators of the Multicenter Study of Hydroxyurea in Sickle Cell Anemia. *N Engl J Med.* 1995;332(20):1317-1322.
- Platt OS, Thorington BD, Brambilla DJ, et al. Pain in sickle cell disease. Rates and risk factors. *N Engl J Med.* 1991;325(1):11-16.
- Vichinsky EP, Neumayr LD, Earles AN, et al.

Acknowledgments

We thank Sang Lee for his technical work on this project and Rosalie Dominik, DrPH, for help with statistical analyses. J.E.B. is an Office of Research on Women's Health BIRCWH scholar.

This work was supported by National Institutes of Health grants K12 HD01441 (J.E.B.), UL1RR025747, HL091265 (K.I.A.), and HL079915 (M.J.T. and K.I.A.) and by an award from the North Carolina State Sickle Cell Program.

Authorship

Contribution: J.E.B. and K.I.A. designed and performed the research, analyzed and interpreted data, and wrote the paper; B.H. performed the research; S.K.J. and D.S. collected data; L.D.C. and E.P.O. interpreted data; M.J.T. interpreted data and contributed to the manuscript; and A.H. collected data, performed the research, and analyzed the data.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Kenneth I. Ataga, Division of Hematology/Oncology, University of North Carolina at Chapel Hill, Rm 3149, Physicians Office Bldg, CB# 7305, Chapel Hill, NC 27599-7305; e-mail: kataga@med.unc.edu.

- Causes and outcomes of the acute chest syndrome in sickle cell disease. National Acute Chest Syndrome Study Group. *N Engl J Med*. 2000;342(25):1855-1865.
34. Machado RF, Anthi A, Steinberg MH, et al. N-terminal pro-brain natriuretic peptide levels and risk of death in sickle cell disease. *JAMA*. 2006;296(3):310-318.
 35. Kato GJ, Gladwin MT, Steinberg MH. Deconstructing sickle cell disease: reappraisal of the role of hemolysis in the development of clinical subphenotypes. *Blood Rev*. 2007;21(1):37-47.
 36. Villagra J, Shiva S, Hunter LA, Machado RF, Gladwin MT, Kato GJ. Platelet activation in patients with sickle disease, hemolysis-associated pulmonary hypertension, and nitric oxide scavenging by cell-free hemoglobin. *Blood*. 2007;110(6):2166-2172.
 37. Kato GJ, Martyr S, Blackwelder WC, et al. Levels of soluble endothelium-derived adhesion molecules in patients with sickle cell disease are associated with pulmonary hypertension, organ dysfunction, and mortality. *Br J Haematol*. 2005;130(6):943-953.
 38. Lygnos MC, Pappa KI, Papadaki HA, P et al. Changes in maternal plasma levels of VEGF, bFGF, TGF-beta1, ET-1 and sKL during uncomplicated pregnancy, hypertensive pregnancy and gestational diabetes. *In Vivo*. 2006;20(1):157-163.
 39. Scott RM, Smith ER. Moyamoya disease and moyamoya syndrome. *N Engl J Med*. 2009;360(12):1226-1237.
 40. Haque AK, Gokhale S, Rampy BA, Adegboyega P, Duarte A, Saldana MJ. Pulmonary hypertension in sickle cell hemoglobinopathy: a clinicopathologic study of 20 cases. *Hum Pathol*. 2002;33(10):1037-1043.
 41. Yock PG, Popp RL. Noninvasive estimation of right ventricular systolic pressure by Doppler ultrasound in patients with tricuspid regurgitation. *Circulation*. 1984;70(4):657-662.
 42. Ataga KI, Smith WR, De Castro LM, et al. Efficacy and safety of the Gardos channel blocker, senicapoc (ICA-17043), in patients with sickle cell anemia. *Blood*. 2008;111(8):3991-3997.
 43. Ataga KI, Jones SK, Hulkower B, Orringer EP, Brittain JE. Association of placenta growth factor with hemolysis and inflammation in sickle cell disease (SCD). *Blood (ASH Annual Meeting Abstracts)*. 2008;112(11):Abstract 2500.



blood[®]

2010 115: 2014-2020
doi:10.1182/blood-2009-04-217950 originally published
online December 29, 2009

Placenta growth factor in sickle cell disease: association with hemolysis and inflammation

Julia E. Brittain, Ben Hulkower, Susan K. Jones, Dell Strayhorn, Laura De Castro, Marilyn J. Telen, Eugene P. Orringer, Alan Hinderliter and Kenneth I. Ataga

Updated information and services can be found at:
<http://www.bloodjournal.org/content/115/10/2014.full.html>

Articles on similar topics can be found in the following Blood collections
[Clinical Trials and Observations](#) (4511 articles)
[Red Cells, Iron, and Erythropoiesis](#) (781 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>