

PSEUDO-ANALYTIC FINITE PARTITION APPROACH TO TEMPERATURE DISTRIBUTION PROBLEM IN HUMAN LIMBS

V.P. SAXENA

School of Mathematics and Allied Sciences
Jiwaji University
Gwalior

and

J.S. BINDRA

Department of Applied Sciences
Mathematics Section
G.N. Engineering College
Ludhiana

(Received March 10, 1987 and in revised form September 8, 1988)

ABSTRACT. The main object of this paper is to introduce the Pseudo Analytic Finite Partition method and to illustrate its use in solving physiological heat distribution problems pertaining to limbs and similar body organs. A two dimensional circular region resembling the cross section of a human or animal limb is considered. The biological properties are assumed to vary along the radial direction. The theoretical model incorporates the effect of blood mass flow and metabolic heat generation. The region is divided into annular sub-regions and Ritz variational finite element method is applied along the radial direction, while for the angular direction, Fourier Series has been used due to uniformity in each annular part.

KEY WORDS AND PHRASES. Blood mass flow rate, rate of metabolic heat generation, variational finite element method.

1980 AMS SUBJECT CLASSIFICATION CODES.

1. INTRODUCTION.

A human body maintains its body core temperature at a uniform temperature under the normal atmospheric conditions. In order to maintain this core temperature, parameters like rate of blood mass flow, rate of metabolic heat generation and, thermal conductivity vary in response to changes in atmospheric conditions. However, in extreme parts of a human body, the core temperature is not uniform at low atmospheric temperatures, where the core temperature of limbs varies extensively as we move away from the body core. This may be because the arterial blood has cooled down while travelling towards the extremities. The heat flow in in-vivo tissues is given by W. Perl [1] as given below.

$$\bar{\rho} \bar{c} \frac{\partial u}{\partial t} = \text{Div}(K \text{ grad } u) + m_b c_b (u_b - u) + s \quad (1.1)$$

where $\bar{\rho}$, \bar{c} , K and S are respectively the density, specific heat, thermal conductivity and, rate of metabolic heat generation in tissues. m_b and c_b are the mass blood flow rate and specific heat of the blood respectively. W. Perl [1] derived and used this equation to study simple problems of heat flow in tissue medium. Chao, Easley and Yang [2] and Chao and Yang [3] also studied temperature distributions in infinite tissue mediums. Cooper and Trezek [4] obtained a solution for a cylindrical symmetry with all the parameters as constant. Saxena [5,6], Saxena and Arya [7,8], and Saxena and Bindra [9,10] used this model to study temperature distributions in skin and subcutaneous tissues using analytical and numerical techniques. This study was performed for a one dimensional steady state case. Later Arya [11], Arya and Saxena [12] and Saxena and Bindra [13] investigated this problem for a two dimensional steady state case in skin and subcutaneous tissues. Here a cylindrical limb having circular cross section with layers of tissues with different properties is considered. The outer boundary is assumed to be exposed to the environment and heat loss takes place due to convection, radiation and, evaporation. The innermost solid cross section is assumed to be at a known variable temperature. This case may occur when one side of the limb contains major blood vessels and thus heated constantly by the blood coming out of main trunk.

2. MATHEMATICAL FORMULATION.

Equation (1.1) in polar steady state form can be written as:

$$\frac{\partial}{\partial r} (Kr \frac{\partial u}{\partial r}) + \frac{K}{r} \frac{\partial^2 u}{\partial r^2} + m_b c_b r (u_b - u) + rS = 0. \quad (2.1)$$

The region is divided into N layers with inner and outer radii equal to a_0 and a_N respectively. (see Fig.1).

The boundary and initial conditions imposed are

$$\left. -K \frac{\partial u}{\partial r} \right|_{r=a_N} = h(u - u_a) + LE \quad (2.2)$$

$$\text{and } u(a_0, \theta) = f(\theta) = u_0 \quad (2.3)$$

where $f(\theta)$ is known. Here h is the heat transfer coefficient and, L and E are respectively the latent heat and rate of sweat evaporation. S is assigned temperature dependent values given by

$$S = s (u_b - u) / u_b, \text{ where } s \text{ is constant.}$$

Thus equation (2.1) along with boundary conditions (2.2) and (2.3) is reduced to discretized polar variational form as given below:

$$I_1 = \frac{1}{2} \int_{a_{i-1}}^{a_i} \left[\bar{r} K_i \left(\frac{\partial v^{(i)}}{\partial r} \right)^2 - \frac{K_i}{\bar{r}} \left(\frac{\partial^2 v^{(i)}}{\partial \theta^2} + \bar{M}_i \bar{r} v^{(i)} \right)^2 \right] d\bar{r} + \delta_1, \quad i=1(1)N \quad (2.4)$$

$$\text{where } \delta_N = \left[\frac{h}{2} (v^{(N)} - v_a)^2 - LE v^{(N)} / u_b \right]_{r=a_N}$$

$$\delta_1 = 0, \quad i=1(1)N-1$$

$\bar{a}_i = a_i/a_o$, $a_i (i=1(1)N)$ is the external radius of i th annular region as shown in Fig.

(1), K_i , \bar{M}_i and $v^{(i)}$ are values of K , \bar{M} and, v in the i th region,

$$\bar{M} = a_o^2 (m_b c_b + s/u_b), \bar{r} = r/a_o, v = (u_b - u)/u_b, v_a = (u_b - u_a)/u_b,$$

The following linear shape functions have been taken for each region:

$$v^{(i)} = \frac{a_i v_{i-1} - a_{i-1} v_i}{a_i - a_{i-1}} + \frac{v_i - v_{i-1}}{a_i - a_{i-1}} \frac{r}{r_i} \tag{2.5}$$

Evaluating integrals (2.4) and assembling these we get

$$I = \sum_{i=1}^N I_i \tag{2.6}$$

Now I is extremized with respect to v_i and the following equations are obtained:

$$\sum_{j=0}^N (E_j^i v_j + F_j^i \frac{\partial^2 v_j}{\partial \theta^2}) = H_i \tag{2.7}$$

Where E_j^i , F_j^i and H_i are constants depending upon physical and physiological parameters.

Now Fourier Series is applied to eliminate the θ variable from the equation (2.7).

We take

$$v_o = A_{oo} + \sum_{n=1}^{\infty} (A_{no} \text{Cos}n\theta + B_{no} \text{Sinn}\theta) \tag{2.8}$$

and

$$v_i = A_{oi} + \sum_{n=1}^{\infty} (A_{ni} \text{Cos}n\theta + B_{ni} \text{Sinn}\theta) \tag{2.9}$$

Here the coefficients A_{oo} , A_{no} and B_{no} are known due to boundary condition (2.3). All the coefficients A_{oi} , A_{ni} and $B_{ni} (i=1,2,\dots,N)$ are unknown. Accordingly, the following system of linear equations is obtained:

$$\begin{aligned} \bar{E}^o \bar{A}_o &= \bar{G}^o \\ \bar{E}^1 \bar{A}_n &= \bar{G}^1 \\ \bar{E}^2 \bar{B}_n &= \bar{G}^2 \end{aligned} \tag{2.10}$$

where $\bar{E}^v = [E_{ij}^v] (i=1(1)N, j=1(1)N)$ are square matrices of order N .

$\bar{A}_0 = [A_{0i}]$, $\bar{A}_n = [A_{ni}]$, $\bar{B}_n = [B_{ni}]$ and, $\bar{G}^v = [G_1^v]$ where $i=1(1)N$, are vectors. E_{ij}^v , A_{0i} , A_{ni} , B_{ni} and G_1^v are constants and they depend on the parameters mentioned earlier. Here only a special case in which the annular cross-section of the limb has been divided into two layers i.e. $N=2$ is considered. These two layers have different biological properties. The outermost layer is supposed to be made of mainly dead tissues. Hence it does not have any metabolic heat generation or blood flow. These quantities have been prescribed in the inner annular part. The core of the limb is assumed to have unsymmetric temperature. Parabolic variation of temperature along the circular boundary has been taken, so

$$V_0 = F(\theta) = \frac{(\theta-\pi)(\theta-2\pi)}{2\pi^2} w_0 + \frac{2\theta(\theta-2\pi)}{(-2\pi^2)} w_1 + \frac{\theta(\theta-\pi)}{2\pi^2} w_0$$

where $w_0 = (u_b - u_\alpha)/u_b$, $w_1 = (u_b - u_\beta)/u_b$, u_α and u_β are temperatures at

$\theta=0$ and $\theta=\pi$ respectively. Using $F(\theta)$ in terms of Fourier series, the constants A_{00} , A_{n0} , and B_{n0} are determined and then substituted in the system of equations (2.10). These equations are solved to find the values of A_{0i} , A_{ni} , and B_{ni} which in turn are substituted in expressions (2.9) to obtain v_1 . Using (2.5) and (2.9), the temperature profiles are obtained for each sub-region.

3. NUMERICAL RESULTS

The following values of physical and physiological parameters and constants have been taken:

$$\begin{aligned} K_1 &= 0.06 \text{ Cal/cm-min. deg. C, } K_2 = 0.03 \text{ Cal/cm-min. deg. C.,} \\ a_0 &= 2.5 \text{ cm, } a_1 = 5 \text{ cm, } a_2 = 7.5 \text{ cm, } L = 579 \text{ Cal/gm,} \\ h &= 0.009 \text{ Cal/cm}^2\text{-min. deg. C, } u_b = 37^\circ\text{C,} \\ M_1 &= (m_b c_b)_1 = 0.003 \text{ Cal/cm}^3\text{-min. deg. C, } M_2 = (m_b c_b)_2 = 0 \\ (S)_1 &= s_1 = 0.0357 \text{ Cal/cm}^3\text{-min, } (S)_2 = s_2 = 0, \\ u_\alpha &= 30^\circ\text{C, } u_\beta = 34^\circ\text{C, } u_a = 15^\circ, E = 0, \end{aligned}$$

Graphs have been plotted between u and θ and the temperature profiles have been shown in fig. 2, 3 and, 4. Fig. 2 is a geometrical representation of the boundary conditions and figures 3 and 4 give temperature variation in direction around the two annular partitions. These two curves are significantly different from the curve in Fig. 2. There is a slow rise in the temperature at $\theta=0$. This rise becomes sharper later on and obviously the temperature takes on its maximum value at $\theta=\pi$. The linear variation of temperature with respect to r is easily seen by comparing the figures 3 and 4.

The number of computations involved in this method are less as compared to those in variational finite element method for two dimensional case.




Fig. 1. ANNULAR CROSS-SECTION OF A HUMAN LIMB DIVIDED INTO N LAYERS. SHADED PORTION IS SOLID CIRCULAR SECTION.




Fig.2. GRAPH SHOWING THE GEOMETRICAL REPRESENTATION OF THE BOUNDARY CONDITIONS.




Fig. 3. GRAPH BETWEEN U_1 AND θ , SHOWING THE TEMPERATURE VARIATIONS AROUND THE INNER ANNULAR PARTITION.




Fig. 4. GRAPH BETWEEN U_2 AND θ , SHOWING THE TEMPERATURE VARIATIONS AROUND THE OUTER ANNULAR PARTITION.

REFERENCES

1. PERL, W., Heat and Water Migration in Body Tissues and Determination of Tissue Blood Flow by Local Clearance Methods, J. Theo. Biol. 2 (1962), 201-235.
2. CHAO, K.N., ELSELY, J.G. and YANG, W.J., Heat and Water Migration in Regional Skins and Subcutaneous Tissues, Bio-Mech symp. ASME, 1975, 69-72.
3. CHAO, K.N. and YANG, W.J., Response of Skin and Tissue Temperature in Sauna and Steam Baths, Bio-Mech. symp., ASME, 69-71.
4. COOPER, T.E. and TREZEK, G.J., Analytical Determination of Cylindrical Source Temperature Fields and their Relation to Thermal Diffusivity of Brain Tissues, Thermal Problems in Bio-Technology, ASME, 1968, NY, 1-15.
5. SAXENA, V.P., Application of Similarity Transformation to Unsteady State Heat Migration Problems in Human SST, Proc. 6th Int. Heat. Trans. Conf. Vol. III, 1978, 65-68.
6. SAXENA, V.P., Temperature Distribution in Human Skin and Subdermal Tissues, J. Theo. Biol., 1983, 102, 277-286.
7. SAXENA, V.P. and ARYA, D., Exact Solution of the Temperature Distribution Problem in Epidermis and Dermis Regions of Human Body. Proc. VNM, Medical and Biological Engineering, Sweden, 1981, 364-366.
8. SAXENA, V.P. and ARYA, D., Variational Finite Element Approach to Heat Distribution Problems in Human Skin and Subdermal Tissues, Proc. 1st Int. Conf. Numerical Methods in Thermal Problems, Pineridge Press, U.K., 1979, 1067-1076.
9. SAXENA, V.P. and BINDRA, J.S., Steady State Temperature Distribution in Dermal Regions of Human Body with Variable Blood Flow, Perspiration and Selfcontrolled Metabolic Heat Generation, Ind. J. Pure app. Math, 15(1), 1984, 31-42.
10. SAXENA, V.P. and BINDRA, J.S., Quadratic Shape Functions in Variational Finite Element Approach to Heat Distribution in Cutaneous and Subcutaneous Tissues, Ind. J. Pure Appl. Math. 18(9), 1987, 846-855.
11. ARYA, D., Two Dimensional Steady State Temperature Distribution in Skin and Subcutaneous Tissues, Ind. J. Pure Appl. Math. 12(1), 1981, 1361-1371.
12. ARYA, D. and SAXENA, V.P., Temperature Variation in Skin and Subcutaneous Layers Under Different Environmental Conditions- A Two Dimensional Study, Ind. J. Pure. Appl. Mathe 17(1), 1986, 84-99.
13. SAXENA, V.P. and BINDRA, J.S., Two Dimensional Steady State Temperature Distribution in Non-Uniform Dermal Layers with Variable Physical and Physiological Conditions. (Under Publication).

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mssklai@cityu.edu.hk