

# FIXED POINT THEOREMS FOR GENERALIZED WEAKLY $S$ -CONTRACTIVE MAPPINGS IN PARTIAL METRIC SPACES

Lakshmi Narayan Mishra<sup>1,2</sup>, Shiv Kant Tiwari<sup>3</sup> and  
Vishnu Narayan Mishra<sup>4,†</sup>

**Abstract** In this paper, we establish some unique fixed point theorems for generalized weakly  $S$ -contractive with nondecreasing and weakly increasing mappings in complete partial metric space. Also, we give some examples for strengthens of our main results.

**Keywords** Fixed point, complete partial metric space, weakly  $S$ -contractive mapping, decreasing mapping, weakly increasing mappings.

**MSC(2010)** 47H10, 54H25, 54E50.

## 1. Introduction and Preliminaries

In 1906, Frechet [13] was first presented the concept of metric space. The notion of partial metric space was introduced by Matthew [17, 18], which is generalization of the metric space. Matthews [17] generalized the Banach contraction principle to the class of complete partial metric space as follows: a self mapping  $f$  on a complete partial metric space  $(X, p)$  has a unique fixed point, if there exist  $0 \leq k < 1$  such that

$$p(fx, fy) \leq kp(x, y), \quad \forall x, y \in X.$$

Several authors have been focused on partial metric spaces and its topological properties (see [4, 5, 7, 19, 23, 28] and references therein). Some interesting work related to generalization of the contraction mapping and metric space can be seen in [10–12, 20–22, 26, 27] and its references. Also, some authors have to generalized fixed point theorems from class of metric space to the class of partial metric space.

First, we recall some useful definitions and results, which is useful throughout the paper.

**Definition 1.1** ([17, 18]). A partial metric  $p$  on a nonempty set  $X$  is a function  $p : X \times X \rightarrow [0, \infty)$  such that:

---

<sup>†</sup>The corresponding author.

<sup>1</sup>Department of Mathematics, National Institute of Technology, Silchar - 788 010, District - Cachar (Assam), India

<sup>2</sup>L. 1627 Awadh Puri Colony Beniganj, Phase III, Opposite Industrial Training Institute (I.T.I.), Ayodhya Main Road, Faizabad - 224 001 (Uttar Pradesh), India  
Email address: lakshminarayanmishra04@gmail.com, l.n.mishra@yahoo.co.in

<sup>3</sup>Department of Applied Mathematics, University Institute of Technology, Rajiv Gandhi Technical University, Bhopal - 462 033 (M.P.) India  
Email address: shivkant.math@gmail.com

<sup>4</sup>Department of Applied Mathematics and Humanities, Sardar Vallabhbhai National Institute of Technology, Ichchhanath Mahadev Dumas Road, Surat - 395 007 (Gujarat), India  
Email address: vishnunarayanmishra@gmail.com, vishnu\_narayanmishra@yahoo.co.in

( $P_1$ )  $p(x, x) = p(x, y) = p(y, y)$  if and only if  $x = y$  (equality),

( $P_2$ )  $p(x, x) \leq p(x, y)$  (small self distance),

( $P_3$ )  $p(x, y) = p(y, x)$  (symmetry),

( $P_4$ )  $p(x, y) + p(z, z) \leq p(x, z) + p(z, y)$  (triangularity)

for all  $x, y, z \in X$ . Then the pair  $(X, p)$  is called a partial metric space. Throughout this paper,  $(X, p)$  represent a partial metric space equipped with a partial metric  $p$ , unless or otherwise stated.

**Example 1.1** ([5, 19]). Consider a mapping  $p : [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$  such that  $p(x, y) = \max\{x, y\}$  for all  $x, y \in [0, \infty)$ . Then  $p$  will be satisfy all the property of partial metric, and hence  $([0, \infty), p)$  is a partial metric space, but fails to the condition of  $p(x, x) = 0$  for all non zero  $x \in [0, \infty)$ . Therefore  $([0, \infty), p)$  is not a metric space.

**Remark 1.1** ([14]). (1) If  $p(x, y) = 0$ , then  $x = y$  but if  $x = y$ , then  $p(x, y)$  may not be zero. (2)  $x \neq y$ , then  $p(x, y) > 0$ .

Also, each partial metric  $p$  on  $X$  generates a  $T_0$  topology  $\tau_p$  on  $X$  with a base of the family of open p-balls  $\{B_p(x, \varepsilon), x \in X, \varepsilon > 0\}$ , where  $B_p(x, \varepsilon) = \{y \in X : p(x, y) < p(x, x) + \varepsilon\}$ , for all  $x \in X$  and  $\varepsilon > 0$  [7]. Let  $(X, p)$  be a partial metric space on  $X$ , then a function  $d^p : X \times X \rightarrow [0, \infty)$  defined as

$$d^p(x, y) = 2p(x, y) - p(x, x) - p(y, y)$$

is a (usual) metric on  $X$  [7]. Furthermore, it is possible to observe that the following

$$\begin{aligned} d^m(x, y) &= \max\{p(x, y) - p(x, x), p(x, y) - p(y, y)\} \\ &= p(x, y) - \min\{p(x, x), p(y, y)\}, \end{aligned}$$

also defines a metric on  $X$ . In fact  $d^p$  and  $d^m$  are equivalent [5].

**Definition 1.2** ([15]). In a partial metric space  $(X, p)$ ,

(1) a sequence  $\{x_n\}$  is said to be convergent to a point  $x \in X$  if and only if  $\lim_{n \rightarrow \infty} p(x_n, x) = p(x, x)$ .

(2) a sequence  $\{x_n\}$  is called Cauchy sequence if and only if  $\lim_{n, m \rightarrow \infty} p(x_n, x_m)$  is finite.

(3) if every Cauchy sequence  $\{x_n\}$  is converges to a point  $x \in X$  such that

$$\lim_{n, m \rightarrow \infty} p(x_n, x_m) = p(x, x),$$

then  $(X, p)$  is known as complete partial metric space.

Chatterjea [6] has been generalized the concept of contractive mappings into C-contractive mappings and Choudhury [9] introduced the weak C-contraction or weakly C-contractive mapping, which is more generalization of C-contractive mappings, as follows:

**Definition 1.3** ([9, 24]). Let  $(X, d)$  be a metric space, then a self mapping  $f$  on  $X$ , satisfying

$$d(fx, fy) \leq \frac{1}{2}[d(x, fy) + d(fx, y)] - \phi(d(x, fy), d(fx, y))$$

for all  $x, y \in X$  and  $\phi : [0, \infty)^2 \rightarrow [0, \infty)$  is a continuous function with  $\phi(x, y) = 0$  if and only if  $x = y = 0$ , is called weakly C-contractive.

In the parallel consequences of weakly C-contractive mapping [9, 24], Shukla and Tiwari [25] has been introduced the concept of S-contractive mapping and generalization its, known as weakly S-contractive mapping, as follows:

**Definition 1.4** ([25]). Let  $(X, d)$  be a metric space, then a self mapping  $f$  on  $X$  is said to be weakly S-contractive mapping or a weak S-contraction, if the following inequality holds:

$$d(fx, fy) \leq \frac{1}{3}[d(x, fy) + d(fx, y) + d(x, y)] - \phi(d(x, fy), d(fx, y), d(x, y))$$

for all  $x, y \in X$  and  $\phi : [0, \infty)^3 \rightarrow [0, \infty)$  is a continuous mapping with  $\phi(x, y, z) = 0$  if and only if  $x = y = z = 0$ .

**Definition 1.5.** A self mapping  $f$  on a partial metric space  $X$  is called nondecreasing, if for all  $x_1, x_2 \in X$ , such that

$$x_1 \leq x_2 \Rightarrow f(x_1) \leq f(x_2).$$

**Definition 1.6** ([3]). In a partial metric space  $(X, p)$ , two self mappings  $f$  and  $g$  are said to be weakly increasing mappings, if for all  $x \in X$  such that

$$g(x) \leq fg(x) \quad \text{and} \quad f(x) \leq gf(x).$$

**Lemma 1.1** ([1, 8]). In a partial metric space  $(X, p)$ , if a sequence  $\{x_n\}$  convergent to a point  $x \in X$ , then  $\lim_{n \rightarrow \infty} p(x_n, x) \leq p(x, z)$  for all  $z \in X$ . Also, if  $p(x, x) = 0$ , then

$$\lim_{n \rightarrow \infty} p(x_n, z) = p(x, z), \quad \forall z \in X.$$

**Lemma 1.2** ([7]). If  $\{x_{2n}\}$  is not a Cauchy sequence in  $(X, p)$ , and two sequences  $\{m(k)\}$  and  $\{n(k)\}$  of positive integers such that  $n(k) > m(k) > k$ , then the following four sequences

$$p(x_{2m(k)}, x_{2n(k)+1}), p(x_{2m(k)}, x_{2n(k)}), p(x_{2m(k)-1}, x_{2n(k)+1}), p(x_{2m(k)-1}, x_{2n(k)})$$

tend to  $\varepsilon > 0$ , when  $k \rightarrow \infty$ .

**Lemma 1.3** ([5, 19]). In a partial metric space  $(X, p)$ ,

(a) a sequence  $\{x_n\}$  is a Cauchy if and only if, it is a Cauchy  $(X, d^p)$ .

(b)  $X$  is complete if and only if it is complete in  $(X, d^p)$ . Also,  $\lim_{n \rightarrow \infty} d^p(x_n, x) = 0$  if and only if

$$\lim_{n, m \rightarrow \infty} p(x_n, x_m) = \lim_{n \rightarrow \infty} p(x_n, x) = p(x, x).$$

(c) if  $\{x_n\}$  is a Cauchy sequence in the metric space  $(X, d^p)$ , we have

$$\lim_{n, m \rightarrow \infty} d^p(x_n, x_m) = 0$$

and therefore by definition of  $d^p$ , we have

$$\lim_{n, m \rightarrow \infty} p(x_n, x_m) = 0.$$

The aim of this paper, is to prove some fixed point results in complete partial metric space for more generalization of weakly S-contractive mappings described in equations (2.1) and (2.16).

## 2. Main results

**Theorem 2.1.** *In a complete partial metric space  $(X, p)$ , a self continuous nondecreasing mapping  $f$  on  $X$ , satisfying the condition*

$$p(fx, fy) \leq M(x, y), \quad (2.1)$$

$$M(x, y) = \max \left\{ \phi(p(x, y)), \phi(p(fx, y)), \phi(p(x, fy)), \right. \\ \left. \phi\left(\frac{p(x, y) + p(fx, y) + p(x, fy)}{3}\right) \right\} \quad (2.2)$$

for all  $x, y \in X$  and  $\phi : [0, \infty) \rightarrow [0, \infty)$  is a continuous function with  $\phi(t) < t \forall t > 0$  and  $\phi(0) = 0$ , has a unique fixed point in  $X$ .

**Proof.** We choose an arbitrary point  $x_0 \in X$ , if  $x_0 = fx_0$ , then theorem follows trivially. Now, we discuss about  $x_0 < fx_0$ , then we can choose  $x_1 \in X$  such that  $fx_0 = x_1$ . Since  $f$  is nondecreasing function, then we have

$$x_0 < x_1 = fx_0.$$

Again, let  $x_2 = fx_1$ , then we get

$$x_0 < x_1 = fx_0 \leq fx_1 = x_2 \leq fx_2.$$

Similarly, proceeding this work, we can construct a sequence  $\{x_n\}$  in  $X$ , such that  $x_{n+1} = fx_n$  with

$$x_0 < x_1 \leq x_2 \leq x_3 \leq \dots \leq x_n \leq x_{n+1} \leq \dots$$

Suppose that  $p(x_{n_0}, x_{n_0+1}) = 0$  for some  $n_0 \geq 0$ , then by Remark 1.1,  $x_{n_0} = x_{n_0+1} = fx_{n_0}$ , that is,  $x_{n_0}$  is a fixed point of  $f$ . So we assume that  $p(x_n, x_{n+1}) \geq 0$  for all  $n \geq 0$ .

Next, we prove that  $\{p(x_n, x_{n+1})\}$  is non increasing sequence in  $X$ . Now, we putting  $x = x_{2n+1}, y = x_{2n}$  in (2.1) we get

$$p(fx_{2n}, fx_{2n+1}) = p(x_{2n+1}, x_{2n+2}) \leq M(x_{2n}, x_{2n+1}). \quad (2.3)$$

By (2.2), we have

$$M(x_{2n}, x_{2n+1}) = \max \left\{ \phi(p(x_{2n}, x_{2n+1})), \phi(p(fx_{2n}, x_{2n+1})), \phi(p(x_{2n}, fx_{2n+1})), \right. \\ \left. \phi\left(\frac{p(x_{2n}, x_{2n+1}) + p(fx_{2n}, x_{2n+1}) + p(x_{2n}, fx_{2n+1})}{3}\right) \right\} \\ = \max \left\{ \phi(p(x_{2n}, x_{2n+1})), \phi(p(x_{2n+1}, x_{2n+1})), \phi(p(x_{2n}, x_{2n+2})), \right. \\ \left. \phi\left(\frac{p(x_{2n}, x_{2n+1}) + p(x_{2n+1}, x_{2n+1}) + p(x_{2n}, x_{2n+2})}{3}\right) \right\}. \quad (2.4)$$

**Case-I:** If  $M(x_{2n}, x_{2n+1}) = \phi(p(x_{2n}, x_{2n+1}))$ , then by equation (2.3) and using the fact  $\phi(t) < t$  for all  $t > 0$ , we have

$$p(x_{2n+1}, x_{2n+2}) < p(x_{2n}, x_{2n+1}).$$

**Case-II:** If  $M(x_{2n}, x_{2n+1}) = \phi(p(x_{2n+1}, x_{2n+1}))$ , then by similar argument of case-I, we get

$$p(x_{2n+1}, x_{2n+2}) < p(x_{2n+1}, x_{2n+1}),$$

which is contradiction of  $(P_2)$ .

**Case-III:** If  $M(x_{2n}, x_{2n+1}) = \phi(p(x_{2n}, x_{2n+2}))$ , then by similar argument of case-I and applying the property of  $(P_4)$  and  $(P_2)$ , we have

$$p(x_{2n+1}, x_{2n+1}) < p(x_{2n}, x_{2n+1}).$$

**Case-IV:** If  $M(x_{2n}, x_{2n+1}) = \phi\left(\frac{p(x_{2n}, x_{2n+1}) + p(x_{2n+1}, x_{2n+1}) + p(x_{2n}, x_{2n+2})}{3}\right)$ , using similar argument of case-III, we have

$$p(x_{2n+1}, x_{2n+2}) < p(x_{2n}, x_{2n+1}).$$

Thus, in all possible cases, we say that  $\{p(x_{2n}, x_{2n+1})\}$  for all  $n \geq 0$ , is a monotonically decreasing sequence in  $X$ . Since, a monotonic decreasing bounded below sequence convergent to its greatest lower bound [16]. Thus,

$$\lim_{n \rightarrow \infty} p(x_n, x_{n+1}) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} p(x_n, x_n) = 0, \quad \forall n \geq 0. \quad (2.5)$$

Now, we have required to prove that the sequence  $\{x_n\}$  is a Cauchy sequence in the partial metric space  $(X, p)$  and so in  $(X, d^p)$  ( by Lemma 1.3(a). On the contrary, we suppose that the sequence  $\{x_n\}$  is not a Cauchy sequence in  $(X, p)$ , then sequences in Lemma 1.2 tend to  $\varepsilon > 0$ , when  $k \rightarrow \infty$ . Now, we putting  $x = x_{2n(k)-1}$  and  $y = x_{2m(k)}$  in equation (2.1), we get

$$\begin{aligned} & p(x_{2n(k)}, x_{2m(k)+1}) \\ &= p(fx_{2n(k)-1}, fx_{2m(k)}) \leq M(x_{2n(k)-1}, x_{2m(k)}) \\ &= \max \left\{ \phi(p(x_{2n(k)-1}, x_{2m(k)})), \phi(p(fx_{2n(k)-1}, x_{2m(k)})), \phi(p(x_{2n(k)-1}, fx_{2m(k)})), \right. \\ & \quad \left. \phi\left(\frac{p(x_{2n(k)-1}, x_{2m(k)}) + p(fx_{2n(k)-1}, x_{2m(k)}) + p(x_{2n(k)-1}, fx_{2m(k)})}{3}\right) \right\} \\ &= \max \left\{ \phi(p(x_{2n(k)-1}, x_{2m(k)})), \phi(p(x_{2n(k)}, x_{2m(k)})), \phi(p(x_{2n(k)-1}, x_{2m(k)+1})), \right. \\ & \quad \left. \phi\left(\frac{p(x_{2n(k)-1}, x_{2m(k)}) + p(x_{2n(k)}, x_{2m(k)}) + p(x_{2n(k)-1}, x_{2m(k)+1})}{3}\right) \right\}. \end{aligned}$$

Taking  $k \rightarrow \infty$  and using Lemma 1.2 in above inequality, we have

$$\varepsilon \leq \max \{ \phi(\varepsilon), \phi(\varepsilon), \phi(\varepsilon), \phi(\varepsilon) \}.$$

Using the fact  $\phi(t) < t$  for all  $t > 0$ , we get

$$\varepsilon \leq \phi(\varepsilon) < \varepsilon,$$

which is a contradiction with respect to  $\varepsilon > 0$ . Thus  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, p)$  and so, in  $(X, d^p)$  (by Lemma 1.3(a)). Since  $(X, p)$  is complete, so  $(X, d^p)$  is also complete (by Lemma 1.3(b)). Therefore, the Cauchy sequence  $\{x_n\}$  converges in the metric space  $(X, d^p)$ , say  $\lim_{n \rightarrow \infty} d^p(x_n, z) = 0$  then again applying Lemma 1.3(b), we have

$$p(z, z) = \lim_{n \rightarrow \infty} p(x_n, z) = \lim_{n, m \rightarrow \infty} p(x_n, x_m). \quad (2.6)$$

By Lemma 1.3(c) and definition of  $d^p$ , we get

$$\begin{aligned} \lim_{n,m \rightarrow \infty} d^p(x_n, x_m) &= 0 \quad \text{and} \\ d^p(x_n, x_m) &= 2p(x_n, x_m) - p(x_n, x_n) - p(x_m, x_m). \end{aligned}$$

Taking  $n, m \rightarrow \infty$  and using equation (2.5) in above inequality, we obtain

$$\lim_{n,m \rightarrow \infty} p(x_n, x_m) = 0. \quad (2.7)$$

From equations (2.6) and (2.7), we get

$$p(z, z) = \lim_{n \rightarrow \infty} p(x_n, z) = 0. \quad (2.8)$$

By  $(P_4)$ , we have

$$\begin{aligned} p(z, fz) &\leq p(z, x_n) + p(x_n, fz) - p(x_n, x_n). \\ p(z, fz) &\leq p(z, x_n) + p(x_n, fz) = p(z, x_n) + p(fx_{n-1}, fz). \end{aligned}$$

Taking  $n \rightarrow \infty$  in above inequality and using equation (2.8) and Lemma 1.1, we have

$$p(z, fz) \leq p(fz, fz). \quad (2.9)$$

From  $(P_2)$ , we have

$$p(fz, fz) \leq p(z, fz). \quad (2.10)$$

From (2.9) and (2.10), we get

$$p(z, fz) = p(fz, fz). \quad (2.11)$$

Next, we show that  $p(fz, fz) = 0$ . On contrary, we suppose that  $p(fz, fz) > 0$ , then by equation (2.1) and (2.8), we have

$$\begin{aligned} p(fz, fz) &\leq M(z, z) \\ &= \max \left\{ \phi(p(z, z)), \phi(p(fz, z)), \phi(p(z, fz)) \right. \\ &\quad \left. \phi\left(\frac{p(z, z) + p(fz, z) + p(z, fz)}{3}\right) \right\} \\ &< \max \left\{ 0, p(fz, fz), p(fz, fz), \frac{2}{3}p(fz, fz) \right\} \\ &= p(fz, fz), \end{aligned}$$

which is a contradiction, so

$$p(fz, fz) = 0.$$

Using equation (2.11) in above, we get

$$p(z, fz) = 0 \Rightarrow fz = z. \quad (2.12)$$

Hence  $z$  is a fixed point of  $f$ . Now, we claim that the fixed point of  $f$  is unique. Against of our claim, we assume that  $u, v \in X$  be two fixed points of  $f$  with  $u \neq v$ ,

such that  $fu = u$  and  $fv = v$ .

From equation (2.1), we have

$$\begin{aligned} p(u, v) &= p(fu, fv) \leq M(u, v) \\ &= \max \left\{ \phi(p(u, v)), \phi(p(fu, v)), \phi(p(u, fv)), \phi\left(\frac{p(u, v) + p(fu, v) + p(u, fv)}{3}\right) \right\} \\ &< p(u, v), \end{aligned} \quad (2.13)$$

which is a contradiction. Hence  $u = v$ , that is, fixed point of  $f$  is unique.  $\square$

**Example 2.1.** Consider a complete partial metric space  $(X = [0, 1], p)$ , such that  $p(x, y) = \max\{x, y\}$  for all  $x, y \in X$ . We define a self mapping  $f$  on  $X$  in such a way

$$f(x) = \begin{cases} x - 0.91x^3 & : \quad \forall x \leq 0.605227532, \\ \alpha & : \quad \forall \alpha \in [\frac{1}{2}, \infty], x > 0.605227532, \end{cases}$$

and a map  $\phi : [0, \infty) \rightarrow [0, \infty)$  such that

$$\phi(t) = \begin{cases} t - \frac{t^2}{2} - \frac{t^3}{3} & : \quad \forall t \in [0, 1], \\ \beta & : \quad \forall \beta \in [0, 1], t \geq 1. \end{cases}$$

Then equation (2.1) holds for all comparable  $x, y \in X$  (that is,  $x \leq y$  or  $x \geq y$ ), and satisfies all the requirements of Theorem 2.1. Therefore  $f$  has a unique fixed point 0 in  $X$ .

**Corollary 2.1.** In a complete partial metric space  $(X, p)$ , a self continuous non-decreasing mapping  $f$  on  $X$ , satisfying the condition

$$p(fx, fy) \leq \phi(M(x, y)), \quad (2.14)$$

$$M(x, y) = \max \left\{ p(x, y), p(fx, y), p(x, fy), \frac{p(x, y) + p(fx, y) + p(x, fy)}{3} \right\} \quad (2.15)$$

for all  $x, y \in X$  and all other are same as in theorem 2.1, then  $f$  has a unique fixed point in  $X$ .

**Corollary 2.2.** In Corollary 2.1, if we replaced second and third ordinates by  $p(x, fx)$  and  $p(y, fy)$  respectively, and fourth ordinate is replace by average of only  $p(x, fy)$  and  $p(fx, y)$  on  $M(x, y)$ , then it reduces to the similar results of [2, 4].

**Theorem 2.2.** In a complete partial metric space  $(X, p)$ , two self weakly increasing continuous mappings  $f$  and  $g$  on  $X$ , satisfying the condition

$$p(fx, gy) \leq M(x, y), \quad (2.16)$$

$$M(x, y) = \max \left\{ \phi(p(x, y)), \phi(p(fx, y)), \phi(p(x, gy)), \phi\left(\frac{p(x, y) + p(fx, y) + p(x, gy)}{3}\right) \right\} \quad (2.17)$$

for all  $x, y \in X$  and  $\phi : [0, \infty) \rightarrow [0, \infty)$  is a continuous function with  $\phi(t) < t$   $\forall t > 0$  and  $\phi(0) = 0$ , then  $f$  and  $g$  have a unique common fixed point in  $X$ .

**Proof.** We choose an arbitrary point  $x_0 \in X$ , such that  $x_0 = fx_0$  and  $x_0 = gx_0$ , then theorem follows trivially. Therefore, we discuss about  $x_0 \neq fx_0$  and  $x_0 \neq gx_0$ . Now, we choose  $x_1 \in X$ , such that  $fx_0 = x_1$  and  $x_2 \in X$ , such that  $gx_1 = x_2$ . Then, we can construct a sequence  $\{x_n\}$  in  $X$ , in such a way

$$x_{2n+1} = fx_{2n}, \quad x_{2n+2} = gx_{2n+1}.$$

Since  $f$  and  $g$  are weakly increasing mappings on  $X$ , then we have

$$\begin{aligned} x_1 &= fx_0 \leq gfx_0 = gx_1 = x_2, \\ x_2 &= gx_1 \leq fgx_1 = fx_2 = x_3, \\ x_3 &= fx_2 \leq gfx_2 = gx_3 = x_4 \end{aligned}$$

and proceeding this work, we get

$$x_0 \leq x_1 \leq x_2 \leq x_3 \leq \cdots \leq x_n \leq x_{n+1} \leq \cdots .$$

Thus  $\{x_n\}$  is a nondecreasing sequence in  $X$ . If we suppose that  $p(x_{n_0}, x_{n_0+1}) = 0$  for some  $n_0 \geq 0$ , then by Remark 1.1  $x_{n_0} = x_{n_0+1} = fx_{n_0}$ , that is,  $x_{n_0}$  is a fixed point of  $f$  and  $g$ . So we assume that  $p(x_n, x_{n+1}) \geq 0$  for all  $n \geq 0$ .

Firstly, we show that the sequence  $\{p(x_n, x_{n+1})\}$  is non-increasing in  $X$ . Now, we putting  $x = x_{2n}, y = x_{2n+1}$  in (2.16), we get

$$p(fx_{2n}, gx_{2n+1}) = p(x_{2n+1}, x_{2n+2}) \leq M(x_{2n}, x_{2n+1}). \quad (2.18)$$

By (2.16), we have

$$\begin{aligned} M(x_{2n}, x_{2n+1}) &= \max \left\{ \phi(p(x_{2n}, x_{2n+1})), \phi(p(fx_{2n}, x_{2n+1})), \phi(p(x_{2n}, gx_{2n+1})), \right. \\ &\quad \left. \phi\left(\frac{p(x_{2n}, x_{2n+1}) + p(fx_{2n}, x_{2n+1}) + p(x_{2n}, gx_{2n+1})}{3}\right) \right\} \\ &= \max \left\{ \phi(p(x_{2n}, x_{2n+1})), \phi(p(x_{2n+1}, x_{2n+1})), \phi(p(x_{2n}, x_{2n+2})), \right. \\ &\quad \left. \phi\left(\frac{p(x_{2n}, x_{2n+1}) + p(x_{2n+1}, x_{2n+1}) + p(x_{2n}, x_{2n+2})}{3}\right) \right\}. \end{aligned} \quad (2.19)$$

**Case-I:** If  $M(x_{2n}, x_{2n+1}) = \phi(p(x_{2n}, x_{2n+1}))$ , then by equation (2.18) and using the fact  $\phi(t) < t$  for all  $t > 0$ , we have

$$p(x_{2n+1}, x_{2n+2}) < p(x_{2n}, x_{2n+1}).$$

**Case-II:** If  $M(x_{2n}, x_{2n+1}) = \phi(p(x_{2n+1}, x_{2n+1}))$ , then by similar argument of case-I, we get

$$p(x_{2n+1}, x_{2n+2}) < p(x_{2n+1}, x_{2n+1}),$$

which is contradiction of  $(P_2)$ .

**Case-III:** If  $M(x_{2n}, x_{2n+1}) = \phi(p(x_{2n}, x_{2n+2}))$ , using the similar argument of case-I and applying the property of  $(P_4)$  and  $(P_2)$ , we have

$$p(x_{2n+1}, x_{2n+1}) < p(x_{2n}, x_{2n+1}).$$

**Case-IV:** If  $M(x_{2n}, x_{2n+1}) = \phi\left(\frac{p(x_{2n}, x_{2n+1}) + p(x_{2n+1}, x_{2n+1}) + p(x_{2n}, x_{2n+2})}{3}\right)$ , then by similar argument of case-III, we obtain

$$p(x_{2n+1}, x_{2n+2}) < p(x_{2n}, x_{2n+1}).$$

Thus, in all possible cases, we observe that  $\{p(x_{2n}, x_{2n+1})\}$  for all  $n \geq 0$ , is a monotonically decreasing sequence in  $X$ . Then, by the similar argument as in Theorem 2.1, we have

$$\lim_{n \rightarrow \infty} p(x_n, x_{n+1}) = 0 \text{ and } \lim_{n \rightarrow \infty} p(x_n, x_n) = 0, \quad \forall n \geq 0. \quad (2.20)$$

Also, we will show that  $\{x_n\}$  is a Cauchy sequence in the partial metric space  $(X, p)$ . By similar arguments as used in case of proving Theorem 2.1, we find that the sequence  $\{x_n\}$  is a Cauchy sequence. Now, we putting  $x = x_{2n(k)}$  and  $y = x_{2m(k)-1}$  in equation (2.16). We get

$$\begin{aligned} & p(x_{2n(k)+1}, x_{2m(k)}) \\ &= p(fx_{2n(k)}, gx_{2m(k)-1}) \leq M(x_{2n(k)}, x_{2m(k)-1}) \\ &= \max \left\{ \phi(p(x_{2n(k)}, x_{2m(k)-1})), \phi(p(fx_{2n(k)}, x_{2m(k)-1})), \phi(p(x_{2n(k)}, gx_{2m(k)-1})), \right. \\ & \quad \left. \phi\left(\frac{p(x_{2n(k)}, x_{2m(k)-1}) + p(fx_{2n(k)}, x_{2m(k)-1}) + p(x_{2n(k)}, gx_{2m(k)-1})}{3}\right) \right\} \\ &= \max \left\{ \phi(p(x_{2n(k)}, x_{2m(k)-1})), \phi(p(x_{2n(k)+1}, x_{2m(k)-1})), \phi(p(x_{2n(k)}, x_{2m(k)})), \right. \\ & \quad \left. \phi\left(\frac{p(x_{2n(k)}, x_{2m(k)-1}) + p(x_{2n(k)+1}, x_{2m(k)-1}) + p(x_{2n(k)}, x_{2m(k)})}{3}\right) \right\}. \end{aligned}$$

Taking  $k \rightarrow \infty$  and applying Lemma 1.2 in above inequality, we have

$$\varepsilon \leq \max \{ \phi(\varepsilon), \phi(\varepsilon), \phi(\varepsilon), \phi(\varepsilon) \}.$$

Using the fact  $\phi(t) < t$  for all  $t > 0$ , we get

$$\varepsilon \leq \phi(\varepsilon) < \varepsilon,$$

which is a contradiction with respect to  $\varepsilon > 0$ . Thus  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, p)$  and so, in  $(X, d^p)$  (by Lemma 1.3(a)). Further, the similar argument of Theorem 2.1, we have  $\lim_{n \rightarrow \infty} d^p(x_n, z) = 0$  then by Lemma 1.3(b), gives that

$$p(z, z) = \lim_{n \rightarrow \infty} p(x_n, z) = \lim_{n, m \rightarrow \infty} p(x_n, x_m). \quad (2.21)$$

From Lemma 1.3(c), we have

$$\lim_{n, m \rightarrow \infty} d^p(x_n, x_m) = 0.$$

So, by definition of  $d^p$ , we get

$$d^p(x_n, x_m) = 2p(x_n, x_m) - p(x_n, x_n) - p(x_m, x_m).$$

Taking  $n, m \rightarrow \infty$  and using equation (2.20) in above, we obtain

$$\lim_{n, m \rightarrow \infty} p(x_n, x_m) = 0. \quad (2.22)$$

From equations (2.21) and (2.22), we get

$$p(z, z) = \lim_{n \rightarrow \infty} p(x_n, z) = 0. \quad (2.23)$$

By  $(P_4)$ , we have

$$\begin{aligned} p(z, fz) &\leq p(z, x_{2n+1}) + p(x_{2n+1}, fz) - p(x_{2n+1}, x_{2n+1}) \\ &\leq p(z, x_{2n+1}) + p(x_{2n+1}, fz) = p(z, x_n) + p(fx_{2n}, fz). \end{aligned}$$

Taking  $n \rightarrow \infty$  and using equation (2.23), we obtain

$$p(z, fz) \leq p(fz, fz). \quad (2.24)$$

By  $(P_2)$ , we have

$$p(fz, fz) \leq p(z, fz). \quad (2.25)$$

From (2.24) and (2.25), we get

$$p(z, fz) = p(fz, fz). \quad (2.26)$$

Similarly,

$$p(z, gz) = p(gz, gz). \quad (2.27)$$

Using equations (2.26) and (2.27) and applying property of  $(P_4)$ , we have

$$\begin{aligned} p(z, gz) &\leq p(z, fz) + p(fz, gz) - p(fz, fz), \\ p(z, gz) &\leq p(fz, gz). \end{aligned} \quad (2.28)$$

Similarly,

$$p(z, fz) \leq p(fz, gz). \quad (2.29)$$

Now, we prove that  $p(fz, gz) = 0$ . On contrary, we suppose that  $p(fz, gz) > 0$ , then by equations (2.16), (2.23) and using above inequality, we get

$$\begin{aligned} &p(fz, gz) \\ &\leq M(z, z) \\ &= \max \left\{ \phi(p(z, z)), \phi(p(fz, z)), \phi(p(z, gz)), \phi\left(\frac{p(z, z) + p(fz, z) + p(z, gz)}{3}\right) \right\} \\ &< \max \left\{ p(z, z), p(fz, z), p(z, gz), \frac{p(z, z) + p(fz, z) + p(z, gz)}{3} \right\} \\ &= \max \left\{ p(fz, z), p(z, gz), \frac{p(fz, z) + p(z, gz)}{3} \right\} \\ &= \max \left\{ p(fz, gz), p(fz, gz), \frac{2p(fz, gz)}{3} \right\} = p(fz, gz), \\ &\Rightarrow p(fz, gz) < p(fz, gz), \end{aligned}$$

which is a contradiction to our assumption. Thus we get

$$p(fz, gz) = 0 \Rightarrow p(z, fz) = 0 \text{ and } p(z, gz) = 0.$$

Applying Remark 1.1, we have

$$fz = gz, fz = z \text{ and } gz = z. \quad (2.30)$$

Hence,  $z$  is a common fixed point of  $f$  and  $g$ . Next, we prove that the common fixed point of  $f$  and  $g$  is unique. Let us suppose that  $u, v \in X$  be two fixed points of  $f$  and  $g$  with  $u \neq v$ , such that  $fu = gu = u$  and  $fv = gv = v$ .

From equation (2.16), we have

$$\begin{aligned} & p(u, v) \\ & = p(fu, gv) \leq M(u, v) \\ & = \max \left\{ \phi(p(u, v)), \phi(p(fu, v)), \phi(p(u, gv)), \phi\left(\frac{p(u, v) + p(fu, v) + p(u, gv)}{3}\right) \right\} \\ & < p(u, v), \end{aligned} \tag{2.31}$$

which is a contradiction with respect to  $u \neq v$ . Therefore,  $u = v$ . Thus the proof is complete.  $\square$

**Corollary 2.3.** *In a complete partial metric space  $(X, p)$ , two self weakly increasing continuous mappings  $f$  and  $g$  on  $X$ , satisfying the condition*

$$p(fx, gy) \leq \phi(M(x, y)), \tag{2.32}$$

$$M(x, y) = \max \left\{ p(x, y), p(fx, y), p(x, gy), \frac{p(x, y) + p(fx, y) + p(x, gy)}{3} \right\} \tag{2.33}$$

for all  $x, y \in X$  and all other are same as in Theorem 2.2, then  $f$  and  $g$  have a unique common fixed point in  $X$ .

**Corollary 2.4.** *In Theorem 2.2, if we replaced second and third ordinates by  $p(x, fx)$  and  $p(y, fy)$  respectively, and fourth ordinate is replaced by average of only  $p(x, fy)$  and  $p(fx, y)$  on  $M(x, y)$ , where  $f$  and  $g$  are weakly isotone increasing mappings such that  $fx \leq gfx \leq fgfx$  and  $gx \leq fgx \leq fgfx$  instead of weakly increasing mappings, then it reduces to the main result of [23].*

## Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

## Acknowledgements

The authors take this opportunity to express their deep gratitude to Dr. Deepmala, SQC & OR Unit, Indian Statistical Institute, Kolkata, for her valuable suggestions for the improvement of the paper. The authors are extremely grateful to the reviewer for a careful reading of the manuscript and making valuable suggestions leading to a better presentation of the paper. The first author Lakshmi Narayan Mishra is thankful to the Ministry of Human Resource Development (MHRD) India for financial support to carry out the above research work. The authors are grateful to all the editorial board members and reviewers of esteemed journal i.e. Journal of Applied Analysis and Computation (JAAC). The second author Shiv Kant Tiwari is grateful to his elder brother Mr. Vaidurya Mani Tiwari and Ms. Jagrati Bilthare for their encouragement and support. The third author Vishnu Narayan Mishra

acknowledges that this project was supported by the Cumulative Professional Development Allowance (CPDA), SVNIT, Surat (Gujarat), India. All the authors carried out the proof of Lemmas, Theorems, Corollaries and Example. Each author contributed equally in the development of the manuscript. Vishnu Narayan Mishra conceived of the study and participated in its design and coordination. All the authors read and approved the final version of manuscript. The authors declare that there is no conflict of interests regarding the publication of this research article.

## References

- [1] T. Abdeljawad, *Fixed points for generalized weakly contractive mappings in partial metric spaces*, Math. Comput. Modelling, 54(2011), 2923–2927.
- [2] I. Altun and A. Erduran, *Fixed point theorems for monotone mappings on partial metric spaces*, Fixed Point Theory Appl., (2011) Article ID 508730 doi: 10.1155/2011/508730.
- [3] I. Altun, B. Damjanovic and D. Djoric, *Fixed point and common fixed point theorems on ordered cone metric spaces*, Appl. Math. Lett., 23(2009), 310–316.
- [4] I. Altun, F. Sola and H. Simsek, *Generalized contraction on partial metric spaces*, Topology Appl., 157(18)(2010), 2778–2785.
- [5] H. Aydi, S. Hadj-Amor and E. Karapinar, *Berinde-Type generalized contractions on partial metric spaces*, Abstr. Appl. Anal., (2013) Article ID 312479 doi: 10.1155/2013.
- [6] S.K. Chatterjee, *Fixed point theorems*, C.R. Acad. Bulgare Sci., 25(1975), 727–730.
- [7] C. Chen and C. Zhu, *Fixed point theorems for weakly C-contractive mappings in partial metric spaces*, Fixed Point Theory Appl., (2013) doi: 10.1186/1687-1812-2013-107.
- [8] K.P. Chi, E. Karapinar and T.D. Thanh, *A generalized contraction principle in partial metric spaces*, Math. Comput. Modelling, 55(2012), 71673–1681.
- [9] B.S. Choudhury, *Unique fixed point theorem for weak C-contractive mappings*, Kathmandu Univ. J. Sci. Eng. Tech., 5(1)(2009), 6–13.
- [10] Deepmala, *A Study of Fixed Point Theorems for Nonlinear Contractions and Its Applications* [Ph.D. thesis], Pt. Ravishankar Shukla University, Raipur, India, 2014.
- [11] Deepmala, *Existence theorems for solvability of a functional equation arising in dynamic programming*, International Journal of Mathematics and Mathematical Sciences, 2014(2014), Article ID 706585, 9 pages.
- [12] Deepmala and H.K. Pathak, *A study on some problems on existence of solutions for nonlinear functional-integral equations*, Acta Mathematica Scientia, Series B English Edition, 33(5)(2013), 1305–1313.
- [13] M. Frechet, *Sur quelques point du calcul fonctionnel*, Rendiconti del circolo Matematico di palermo, 22(1906), 1–74.
- [14] E. Karapinar and I.M. Erhan, *Fixed point theorems for operators on partial metric spaces*, Appl. Math. Lett., 24(2011), 1894–1899.

- [15] E. Karapinar and I.S. Yuce, *Fixed point theory for cyclic generalized weak  $\phi$ -contraction on partial metric spaces*, Abstr. Appl. Anal., (2012) Article ID 491542 doi: 10.1155/2012.
- [16] S.C. Malik and S. Arora, *Mathematical analysis, fourth edition*, New age Int. Publ., (2010), page-87.
- [17] S.G. Matthews, *Partial Metric Topology, in proceeding of 8<sup>th</sup> summer conference on General Topology and Applications*, at Queens College (1922) in: Annals of the New York Academy of Sciences, 728(1994), 183–197.
- [18] S.G. Matthews, *Partial Metric Topology*, Research Report 212, Dept. of Comput. Sci., University of Warwick, 1992.
- [19] L.N. Mishra, S.K. Tiwari, V.N. Mishra and I.A. Khan, *Unique fixed point theorems for generalized contractive mappings in partial metric spaces*, J. Function Spaces, (2015) Manuscript Id 960827, in press.
- [20] V.N. Mishra, M.L. Mittal, and U. Singh, *On best approximation in locally convex space*, Varahmihir Journal of Mathematical Sciences India, 6(1)(2006), 43–48.
- [21] V.N. Mishra, *Some Problems on Approximations of Functions in Banach spaces [Ph.D. thesis]*, Indian Institute of Technology, Uttarakhand, India, 2007.
- [22] V.N. Mishra and L.N. Mishra, *Trigonometric approximation in  $L_p(p \geq 1)$ -spaces*, International Journal of Contemporary Mathematical Sciences, 7(2012), 909–918.
- [23] H.K. Nashine, Z. Kadelburg and S. Radenovic, *Common fixed point theorems for weakly isotone increasing mappings in ordered partial metric spaces*, Math. Comput. Modelling, 57(2013), 2355–2365.
- [24] W. Shatanawi, *Fixed point theorems for nonlinear weakly  $C$ -contractive mappings in metric spaces*, Math. Comput. Modelling, 54(2011), 2816–2826.
- [25] D.P. Shukla and S.K. Tiwari, *Unique fixed point theorem for weakly  $S$ -contractive mappings*, Gen. Math. Notes, 4(1)(2011), 28–34.
- [26] D.P. Shukla, S.K. Tiwari and S.K. Shukla, *Fixed point theorems for a pair of compatible mappings in integral type equation*, Int. J. of Math. Sci. & Engg. Appls, 7(VI)(2013), 413–419.
- [27] D.P. Shukla, S.K. Tiwari and S.K. Shukla, *Unique common fixed point theorems for compatible mappings in complete metric space*, Gen. Math. Notes, 18(1)(2013), 13–23.
- [28] S. Wang, *Multidimensional fixed point theorems for isotone mappings in partially ordered metric spaces*, Fixed Point Theory Appl., 54(2014), 2014–137.