

# Three-dimensional evaluation of midfacial asymmetry in patients with nonsyndromic unilateral cleft lip and palate by cone-beam computed tomography

Youn-Kyung Choi<sup>a</sup>  
Soo-Byung Park<sup>a</sup>  
Yong-Il Kim<sup>b,c</sup>  
Woo-Sung Son<sup>a</sup>

<sup>a</sup>Department of Orthodontics, Pusan National University Dental Hospital, Yangsan, Korea

<sup>b</sup>Department of Orthodontics, School of Dentistry, Pusan National University, Busan, Korea

<sup>c</sup>Biomedical Research Institute, Pusan National University Hospital, Busan, Korea

**Objective:** To compare three-dimensionally the midfacial hard- and soft-tissue asymmetries between the affected and the unaffected sides and determine the relationship between the hard tissue and the overlying soft tissue in patients with nonsyndromic complete unilateral cleft lip and palate (UCLP) by cone-beam computed tomography (CBCT) analysis. **Methods:** The maxillofacial regions of 26 adults (18 men, 8 women) with nonsyndromic UCLP were scanned by CBCT and reconstructed by three-dimensional dental imaging. The frontal-view midfacial analysis was based on a 3 × 3 grid of vertical and horizontal lines and their intersecting points. Two additional points were used for assessing the dentoalveolar area. Linear and surface measurements from three reference planes (Basion-perpendicular, midsagittal reference, and Frankfurt horizontal planes) to the intersecting points were used to evaluate the anteroposterior, transverse, and vertical asymmetries as well as convexity or concavity. **Results:** Anteroposteriorly, the soft tissue in the nasolabial and dentoalveolar regions was significantly thicker and positioned more anteriorly on the affected side than on the unaffected side ( $p < 0.05$ ). The hard tissue in the dentoalveolar region was significantly retruded on the affected side compared with the unaffected side ( $p < 0.05$ ). The other midfacial regions showed no significant differences. **Conclusions:** With the exception of the nasolabial and dentoalveolar regions, no distinctive midfacial hard- and soft-tissue asymmetries exist between the affected and the unaffected sides in patients with nonsyndromic UCLP. [Korean J Orthod 2013;43(3):113-119]

**Key words:** Computed tomography, Cleft lip and palate, Asymmetry, Soft tissue

Received September 10, 2012; Revised November 13, 2012; Accepted November 14, 2012.

**Corresponding author:** Yong-Il Kim.

Assistant Professor, Department of Orthodontics, Pusan National University, Biomedical Research Institute, Pusan National University Hospital, Gudeok-ro 137, Seo-gu, Busan 626-731, Korea.

Tel +82-51-240-7432 e-mail kimyongil@pusan.ac.kr

\*This work was supported by a 2-Year Research Grant of Pusan National University. Choi YK and Park SB were equally contributed to this article.

The authors report no commercial, proprietary, or financial interest in the products or companies described in this article.

© 2013 The Korean Association of Orthodontists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**INTRODUCTION**

Cleft lip, palate, or both are the most frequently occurring congenital facial deformities, having an incidence rate of 0.65% among newborns, with ethnic and geographic variations.<sup>1,2</sup> The typical manifestations of cleft lip and/or palate include midfacial deficiency related to retruded maxilla, contralaterally deviated anterior nasal spine, distorted nasal septum, collapsed piriform rim, and subsided nasal base.<sup>3,4</sup> In the unilateral cleft, asymmetrical maxillary bone defects and distortions are common.<sup>3,5-11</sup> Even after primary corrective surgery to improve the appearance, patients with unilateral cleft lip and palate (UCLP) show a distinct facial asymmetry.<sup>12,13</sup>

Numerous studies of the asymmetry of the cranial base, nasolabial region, and mandible of patients with UCLP have been conducted.<sup>6,8-11,14-16</sup> In the nasolabial region, nasal septum distortion and deviation toward the affected side have been reported.<sup>6,8-11</sup> However, the

mandibular asymmetry is controversial.<sup>8,14-16</sup> Some authors reported significant differences in the lower face, especially in the mandible,<sup>14,15</sup> whereas others offered contradictory data.<sup>8,16</sup> Nevertheless, midfacial analysis to distinguish areas of obvious asymmetry has rarely been conducted.

This lacuna of research is attributable to the serious limitations of lateral cephalometric radiographs: specifically, their two-dimensional nature, superimposition of bony structures in the craniofacial and midfacial regions, and incomplete assessment of soft tissue.<sup>17</sup> Kim et al.<sup>18</sup> suggested grid-based cone-beam computed tomography (CBCT) analysis of hard- and soft-tissue changes, and Meyer-Marcotty et al.<sup>7</sup> recommended registration of the original and mirrored data by using three-dimensional surface scans to analyze facial asymmetry. CBCT is a good tool for quantitative analysis of hard and soft tissues in the midface.<sup>19,20</sup> Nonetheless, only a few reports on the use of CBCT for measuring superficial

**Table 1.** Reference points and planes

Reference point and plane	Description
Porion (Po)	Most superior point of external auditory meatus
Orbitale (Or)	Most inferior point of infraorbital margin
Basion (Ba)	Most posterior inferior point of occipital bone at anterior margin of foramen magnum
Nasion (Na)	Most anterior point of frontonasal suture on midsagittal plane
Frankfort horizontal (FH) plane	Plane was constructed by both sides of Po and non-affected side of Or
Midsagittal reference (MSR) plane	Plane was perpendicular to FH plane and passed through Na and Ba
Ba-perpendicular plane	Plane was perpendicular to FH and midsagittal planes and passed through Ba
Superior horizontal (SH) plane	Plane parallel to the FH plane and passing through the lowest point of the orbit
Inferior horizontal (IH) plane	Plane parallel to the FH plane and passing through the inferior border of the zygomaticomaxillary suture
Middle horizontal (MH) plane	Plane bisecting the SH and IH planes
Inner sagittal (IS) plane	Plane parallel to the midsagittal plane and passing through the outer rim of the piriform aperture
Lateral sagittal (LS) plane	plane parallel to the FH plane and passing through the lateral border of the orbit
Middle sagittal (MS) plane	Plane bisecting the IS and LS planes
P1 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between SH plane and IS plane
P2 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between SH plane and MS plane
P3 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between SH plane and LS plane
P4 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between MH plane and IS plane
P5 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between MH plane and MS plane
P6 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between MH plane and LS plane
P7 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between LH plane and IS plane
P8 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between LH plane and MS plane
P9 <sub>hard/soft</sub>	Intersecting point <sub>hard/soft</sub> between LH plane and LS plane
P10 <sub>hard/soft</sub>	Midpoint <sub>hard/soft</sub> of distance between alare and cheilion
P11 <sub>hard/soft</sub>	Midpoint <sub>hard/soft</sub> of distance between P10 <sub>hard/soft</sub> and midsagittal plane

midfacial asymmetry in UCLP exist.

The aims of this study were to compare three-dimensionally the midfacial hard- and soft-tissue asymmetries between the affected and the unaffected sides and determine the relationship between the hard tissue and the overlying soft tissue in patients with complete UCLP by CBCT analysis. The null hypothesis was that no significant midfacial hard- and soft-tissue differences would exist between the affected and the unaffected sides in complete UCLP.

## MATERIALS AND METHODS

### Subjects

Twenty-six adults (18 men, 8 women; mean age, 22.79  $\pm$  6.38 years; range, 19 - 43 years) who had undergone complete UCLP correction were selected from the Department of Orthodontics, Pusan National University Dental Hospital (Busan, Korea). The inclusion criteria were (1) availability of birth and treatment records, (2) surgical correction of the cleft in infancy (primary lip correction within 1 year; primary palate correction before 2 years), and (3) no orthognathic surgery. The exclusion criteria were (1) diagnosis of a syndrome, (2) either growth or mental retardation, and (3) maxillary distraction osteogenesis in adolescence. This study was reviewed and approved by the Ethics Committee of Pusan National University Hospital (E-2011040).


### Data acquisition

Before the orthodontic treatment, CBCT scans (PaX-Zenith 3D; Vatech Co., Gyeonggi-Do, Korea) were recorded, with the subject in an upright position for maximum intercuspation. The Frankfurt horizontal (FH) plane was parallel to the floor. The maxillofacial regions were scanned by using a field of view of 20  $\times$  19 cm, tube voltage of 90 kVp, tube current of 4.0 mA, and scan time of 24 s. The scans were reconstructed by using three-dimensional dental imaging software (Ez3D2009; E-WOO Technology Co., Ltd., Gyeonggi-Do, Korea). Then, the craniofacial structures in the three-dimensional images were re-orientated according to reference planes: the Basion-perpendicular, midsagittal reference (MSR), and FH planes (Table 1).

One operator performed all the measurements twice with an interval of 2 weeks. Under the same conditions, the linear and surface measurements of the hard and soft tissues were obtained (window width: 4,000; window level: 1,000).

### Linear measurements

The analytical method involved the use of a 3  $\times$  3 grid of vertical and horizontal planes to divide the midfacial region unilaterally. The vertical planes were the inner




**Figure 1.** Planes parallel to the reference planes and their intersecting points. SH, Superior horizontal; MH, middle horizontal; IH, inferior horizontal; LS, lateral sagittal; MS, middle sagittal; IS, inner sagittal. See Table 1 for the definitions.

sagittal (IS), middle sagittal (MS), and lateral sagittal (LS) planes and the horizontal planes were the superior horizontal (SH), middle horizontal (MH), and inferior horizontal (IH) planes (Table 1, Figure 1). Their intersecting points were P1<sub>hard/soft</sub> to P9<sub>hard/soft</sub>. Two additional points were used unilaterally to assess the dentoalveolar area: P10<sub>hard/soft</sub>, midpoint of the distance between the alare and the cheilion, and P11<sub>hard/soft</sub>, midpoint of the distance between P10<sub>hard/soft</sub> and the MSR plane (Table 1, Figure 1).

We obtained 22, five, and two linear measurements for assessing anteroposterior, transverse, and vertical hard- and soft-tissue asymmetries, respectively, in the midfacial and dentoalveolar areas on each side. To evaluate the anteroposterior asymmetry, the distances from the Ba-perpendicular plane to the corresponding hard- and soft-tissue points (P1<sub>hard/soft</sub> to P11<sub>hard/soft</sub>) on the affected and unaffected sides were measured (Figure 2A). To evaluate the transverse asymmetry, the distances from the MSR plane to the IS and LS planes, along with the most lateral soft-tissue point on each of the vertical lines, were measured (Figure 2B). Finally, the vertical asymmetry was calculated by measuring the distances from the FH plane to the SH and IH planes. The SH plane on the unaffected side, referred to as the FH plane, was used as the reference line (Figure 2C).

### Surface measurements

To evaluate the midfacial asymmetry in terms of convexity or concavity between the affected and the unaffected sides, the surface areas of the maxilla and



**Figure 2.** Linear and surface measurements. A, Distance between the Ba-perpendicular plane and the corresponding hard- and soft-tissue points. B, Distance between the midsagittal reference (MSR) plane and the inner sagittal (IS) and lateral sagittal (LS) planes as well as the most lateral soft-tissue point on each of the vertical lines. C, Distance between the Frankfurt horizontal (FH) plane and the superior horizontal (SH) and inferior horizontal (IH) planes. D, Surface measurements between IS plane and line from inner measurement point to lateral measurement point. See Table 1 for the abbreviations.

zygomatic bone on each of the horizontal planes were measured. The superior area was between the anterior surface of the maxilla and the line from P1<sub>hard</sub> to P3<sub>hard</sub>, the middle area was between the anterior surface of the maxilla and the line from P4<sub>hard</sub> to P6<sub>hard</sub>, and the inferior area was between the anterior surface of maxilla and the line from P7<sub>hard</sub> to P9<sub>hard</sub>. Negative values indicated concavity and positive values indicated convexity (Figure 2D).

**Statistical analysis**

To assess the intraobserver variability and reproducibility, the measurements were checked twice with an interval of 14 days. A paired t-test was used for all the comparisons. The differences were considered significant at  $p < 0.05$ . PASW Statistics version 18.0 for Windows (IBM Co., Armonk, NY, USA) was used for the statistical analyses.

**RESULTS**

The intraobserver agreement of the measurements was excellent (intraclass correlation coefficient range: 0.966 - 0.998).

**Anteroposterior asymmetry**

In the soft-tissue analysis, the distances from the Ba-perpendicular plane to P7<sub>soft</sub> on the unaffected side ( $p = 0.010$ ) and P11<sub>soft</sub> on the affected side ( $p = 0.006$ ) were significantly decreased (Table 2). In the hard-tissue analysis, the distances from the Ba-perpendicular plane to P10<sub>hard</sub> ( $p = 0.034$ ) and P11<sub>hard</sub> ( $p = 0.006$ ) on the affected side showed a significant reduction. None of the other midfacial measurements showed significant differences ( $p > 0.05$ ).

**Table 2.** Results of the CBCT analysis of the sagittal hard- and soft-tissue differences

Difference (mm)	Affected side	Non-affected side	p-value
<b>Soft tissue difference</b>			
P1 <sub>hard/soft</sub>	86.74 ± 3.83	86.70 ± 3.83	0.898
P2 <sub>hard/soft</sub>	82.88 ± 3.73	82.28 ± 3.38	0.145
P3 <sub>hard/soft</sub>	77.93 ± 3.97	76.50 ± 4.28	0.089
P4 <sub>hard/soft</sub>	86.84 ± 4.65	86.57 ± 4.42	0.244
P5 <sub>hard/soft</sub>	83.47 ± 3.75	82.70 ± 3.76	0.129
P6 <sub>hard/soft</sub>	78.85 ± 3.64	77.95 ± 3.67	0.206
P7 <sub>hard/soft</sub>	85.34 ± 4.04	84.07 ± 5.56	0.010*
P8 <sub>hard/soft</sub>	83.84 ± 3.96	83.18 ± 4.06	0.097
P9 <sub>hard/soft</sub>	78.99 ± 3.68	77.92 ± 3.72	0.194
P10 <sub>hard/soft</sub>	88.18 ± 5.15	87.88 ± 5.55	0.415
P11 <sub>hard/soft</sub>	90.72 ± 5.76	91.77 ± 5.58	0.006*
<b>Hard tissue difference</b>			
P1 <sub>hard/soft</sub>	79.39 ± 4.74	79.18 ± 4.24	0.696
P2 <sub>hard/soft</sub>	75.77 ± 4.49	75.00 ± 3.73	0.184
P3 <sub>hard/soft</sub>	69.36 ± 4.69	67.25 ± 3.73	0.050
P4 <sub>hard/soft</sub>	76.32 ± 4.49	77.09 ± 4.52	0.194
P5 <sub>hard/soft</sub>	72.18 ± 4.76	71.63 ± 5.01	0.492
P6 <sub>hard/soft</sub>	66.71 ± 4.56	66.25 ± 4.06	0.187
P7 <sub>hard/soft</sub>	73.27 ± 4.02	73.59 ± 4.36	0.497
P8 <sub>hard/soft</sub>	67.42 ± 4.64	66.82 ± 3.75	0.638
P9 <sub>hard/soft</sub>	62.80 ± 4.25	61.26 ± 3.72	0.168
P10 <sub>hard/soft</sub>	72.83 ± 6.24	74.23 ± 5.62	0.034*
P11 <sub>hard/soft</sub>	76.83 ± 5.38	78.05 ± 5.23	0.006*

Values are presented as mean ± standard deviation. \*Significant difference by the paired t-test ( $p < 0.05$ ). See Table 1 for the definitions of landmarks.

**Table 3.** Results of the CBCT analysis of the soft-tissue thickness at the intersecting points

Measurement (mm)	Affected side	Non-affected side	p-value
P1 <sub>thickness</sub>	7.54 ± 1.82	7.54 ± 1.58	0.646
P2 <sub>thickness</sub>	7.11 ± 1.67	7.17 ± 1.46	0.842
P3 <sub>thickness</sub>	8.98 ± 2.45	9.69 ± 3.43	0.349
P4 <sub>thickness</sub>	10.75 ± 1.92	9.73 ± 2.75	0.021*
P5 <sub>thickness</sub>	11.23 ± 2.10	11.12 ± 2.18	0.228
P6 <sub>thickness</sub>	11.84 ± 3.67	12.77 ± 2.87	0.187
P7 <sub>thickness</sub>	12.14 ± 3.65	10.82 ± 3.47	0.000*
P8 <sub>thickness</sub>	16.64 ± 2.57	16.35 ± 2.27	0.323
P9 <sub>thickness</sub>	16.07 ± 2.61	16.57 ± 3.21	0.471
P10 <sub>thickness</sub>	15.58 ± 2.77	14.23 ± 2.93	0.003*
P11 <sub>thickness</sub>	13.89 ± 2.58	13.50 ± 2.13	0.303

Values are presented as mean ± standard deviation.

\*Significant difference by the paired t-test ( $p < 0.05$ ).

See Table 1 for the definitions of landmarks.

### Transverse and vertical asymmetries

In terms of the transverse asymmetry, only the distance from the MSR plane to the IS plane was significantly different ( $p < 0.05$ ): it was greater on the affected side than on the unaffected side. None of the soft-tissue measurements showed significant differences ( $p > 0.05$ ). Further, neither of the measurements for vertical asymmetry were significant ( $p > 0.05$ ).

### Soft tissue thickness

In terms of the soft-tissue thickness, P4<sub>thickness</sub> ( $p = 0.021$ ), P7<sub>thickness</sub> ( $p = 0.000$ ), and P10<sub>thickness</sub> ( $p = 0.003$ ) showed significant differences on the affected side (Table 3). The middle and lateral regions of the midface showed no significant differences between the sides ( $p > 0.05$ ).

### Surface measurements

No significant differences in the surface measurements were noted ( $p > 0.05$ ; Table 4).

## DISCUSSION

Although much research on asymmetry in patients with UCLP has been conducted, conclusions remain elusive.<sup>6,7,17,21,22</sup> Simply, the midfacial hard- and soft-tissue asymmetries in adults have not been sufficiently evaluated. In the present study, CBCT, with the use of established reference lines and grid planes,<sup>18,22</sup> enabled effective simultaneous analysis of hard- and soft-tissue asymmetries of the midface in adults with nonsyndromic UCLP. As these patients show midfacial defects and

**Table 4.** Results of the CBCT analysis of the surface measurements

Measurement (mm <sup>2</sup> )	Affected side	Non-affected side	p-value
S-area	56.88 ± 28.16	50.11 ± 26.83	0.516
M-area	13.79 ± 34.96	10.05 ± 32.60	0.225
I-area	11.25 ± 49.47	-0.61 ± 64.62	0.872

Values are presented as mean ± standard deviation. Negative and positive values represent concavity and convexity, respectively.

\*Significant difference by the paired t-test ( $p < 0.05$ ).

S-area, Superior area; M-area, middle area; I-area, inferior area.

See Table 1 for the definitions of landmarks.

variations, a cranial structure manifesting a relatively small variation was used as the basis for the reference planes. The FH and MSR planes, given the previous research showing the porion to be an adequately stable structure for use as a reference point,<sup>17,23</sup> were considered acceptable for evaluating midfacial asymmetry caused by a relatively small variation of the cleft.<sup>17</sup>

Tissue defects and scar tissue resulting from early surgery in patients with UCLP can impede normal growth and induce anteroposterior and transverse asymmetries. Nasal asymmetry is a long-recognized stigma of these patients.<sup>9,11,21,24,25</sup> Even so, the midfacial area has rarely been the focus of study.

The results showed that no significant differences existed between the affected and the unaffected sides or between the hard and the soft tissues at most sites ( $p > 0.05$ ). Similarly, previous research has shown significant differences between the affected and the unaffected sides only in the nasolabial and dentoalveolar areas.<sup>8,17</sup> Suri et al.<sup>17</sup> observed no significant differences in bilateral measurements of the craniofacial structures far from the cleft, indicating the absence of major transverse and sagittal asymmetries of the deeper midfacial structures in their study sample. The maxillary sinuses on both sides of the face were similar to the reference in size and lateromedial and anteroposterior locations. Further, Suzuki et al.<sup>24</sup> reported that the size of the sinus associated with UCLP is not dependent on the cleft side. However, others have reported contradictory results, showing asymmetry in the midfacial region, especially of the orbital, maxillary, and nasal regions, by using three-dimensional skull models. Lateral dislocation of the maxilla on the affected side has been noted, and mean retrograde movement has been shown in 52% of the patients.<sup>22,26,27</sup>

With regard to the hard-tissue measurements, only those in the dentoalveolar area were significantly different ( $p < 0.05$ ): the tissue on the affected side was

retruded compared with that on the unaffected side. This seems reasonable, because the dentoalveolar area and piriform aperture directly lie in the contraction area of the cleft.<sup>6,7,17,26</sup> P11<sub>hard</sub> seems to have been affected by asymmetrical posterior displacement of the maxilla itself, due to the absence of the lateral incisor on the affected side. This depression of the dentoalveolar area on the cleft side is similar to the sagittal depression of the bony alar base in the findings of Mølsted and Dahl<sup>9</sup> and Kolbenstvedt et al.<sup>25</sup>

Among the soft-tissue measurements, only in the dentoalveolar (P11<sub>soft</sub>) and nasal (P7<sub>soft</sub>) areas were significant differences noted ( $p < 0.05$ ); however, their positions varied. Specifically, the dentoalveolar area was located relatively posterior on the affected side, because P11<sub>soft</sub> was closer to the cleft. P11<sub>soft</sub> was the site of scar-tissue formation, owing to the flap facilitation and suturing that had been performed in the previous cheiloplasty.<sup>28,29</sup> Conversely, the lower portion of the midface adjacent to the nose was located relatively anteriorly on the affected side. This might have been caused by the soft-tissue thickness, which was greater near the nose (P4<sub>soft</sub> and P7<sub>soft</sub>) than elsewhere. Further, the difference in the soft-tissue thickness may have been caused by the gap between the cleft segment and the postsurgical changes in muscle position and function. Nevertheless, these results are consistent with previous findings, showing greater soft-tissue thickness on the affected side.<sup>3,9,30</sup> In their research on untreated unilateral cleft nose deformities by using a combined photography/CBCT measurement method, Li et al.<sup>3</sup> insisted that bony deformities would determine the soft-tissue contours and the soft tissue, in turn, could camouflage the underlying bony deformities in various patterns and to different extents, making the external configuration less deformed than its bony basis.

The surface measurements of bone concavity or convexity did not show any significant difference between the affected and the unaffected sides in any area ( $p > 0.05$ ). These results confirm that the affected side does not present any particular bony depression. However, with respect to the impact of facial asymmetry on visual perception, Meyer-Marcotty et al.<sup>7</sup> insisted that the greater the facial asymmetry near the midline of the face, the more negative is the evaluation of the face in direct face-to-face interactions among patients with UCLP. The greatest asymmetry in adult patients is in the nasolabial and dentoalveolar areas, which can affect the perception of midfacial asymmetry. Better midfacial symmetry is achievable, therefore, by performing rhinoplasty after orthognathic surgery.

## CONCLUSION

The nasolabial and dentoalveolar regions show significant differences in linear measurements and soft-tissue thickness. No other distinctive hard- and soft-tissue asymmetries exist between the affected and the unaffected sides of the midface in patients with nonsyndromic UCLP.

## REFERENCES

1. Mossey PA, Little J, Munger RG, Dixon MJ, Shaw WC. Cleft lip and palate. *Lancet* 2009;374:1773-85.
2. Baik HS, Keem JH, Kim DJ. The prevalence of cleft lip and/or cleft palate in Korean male adult. *Korean J Orthod* 2001;31:63-9.
3. Li J, Shi B, Liu K, Zheng Q. A preliminary study on the hard-soft tissue relationships among unoperated secondary unilateral cleft nose deformities. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2012;113:300-7.
4. Atherton JD. Morphology of facial bones in skulls with unoperated unilateral cleft palate. *Cleft Palate J* 1967;4:18-30.
5. Joos U. Skeletal growth after muscular reconstruction for cleft lip, alveolus and palate. *Br J Oral Maxillofac Surg* 1995;33:139-44.
6. Stauber I, Vairaktaris E, Holst A, Schuster M, Hirschfelder U, Neukam FW, et al. Three-dimensional analysis of facial symmetry in cleft lip and palate patients using optical surface data. *J Orofac Orthop* 2008;69:268-82.
7. Meyer-Marcotty P, Alpers GW, Gerdes AB, Stellzig-Eisenhauer A. Impact of facial asymmetry in visual perception: a 3-dimensional data analysis. *Am J Orthod Dentofacial Orthop* 2010;137:168.e1-8.
8. Son WS, Kim MK. Facial asymmetry of unilateral cleft lip and palate patients. *Korean J Orthod* 1995; 25:13-8.
9. Mølsted K, Dahl E. Asymmetry of the maxilla in children with complete unilateral cleft lip and palate. *Cleft Palate J* 1990;27:184-90.
10. Mølsted K, Kjaer I, Dahl E. Cranial base in newborns with complete cleft lip and palate: radiographic study. *Cleft Palate Craniofac J* 1995;32:199-205.
11. Ras F, Habets LL, van Ginkel FC, Prahl-Andersen B. Three-dimensional evaluation of facial asymmetry in cleft lip and palate. *Cleft Palate Craniofac J* 1994; 31:116-21.
12. Broder HL, Smith FB, Strauss RP. Effects of visible and invisible orofacial defects on self-perception and adjustment across developmental eras and gender. *Cleft Palate Craniofac J* 1994;31:429-36.
13. Lee LW, Chen SH, Yu CC, Lo LJ, Lee SR, Chen YR. Stigma, body image, and quality of life in women

- seeking orthognathic surgery. *Plast Reconstr Surg* 2007;120:225-31.
14. Laspos CP, Kyrkanides S, Tallents RH, Moss ME, Subtelny JD. Mandibular asymmetry in noncleft and unilateral cleft lip and palate individuals. *Cleft Palate Craniofac J* 1997;34:410-6.
  15. Smahel Z, Brejcha M. Differences in craniofacial morphology between complete and incomplete unilateral cleft lip and palate in adults. *Cleft Palate J* 1983;20:113-27.
  16. Kurt G, Bayram M, Uysal T, Ozer M. Mandibular asymmetry in cleft lip and palate patients. *Eur J Orthod* 2010;32:19-23.
  17. Suri S, Utreja A, Khandelwal N, Mago SK. Craniofacial computerized tomography analysis of the midface of patients with repaired complete unilateral cleft lip and palate. *Am J Orthod Dentofacial Orthop* 2008;134:418-29.
  18. Kim YI, Kim JR, Park SB. Three-dimensional analysis of midfacial soft tissue changes according to maxillary superior movement after horizontal osteotomy of the maxilla. *J Craniofac Surg* 2010;21:1587-90.
  19. de Moraes ME, Hollender LG, Chen CS, Moraes LC, Balducci I. Evaluating craniofacial asymmetry with digital cephalometric images and cone-beam computed tomography. *Am J Orthod Dentofacial Orthop* 2011;139:e523-31.
  20. Hwang HS, Hwang CH, Lee KH, Kang BC. Maxillofacial 3-dimensional image analysis for the diagnosis of facial asymmetry. *Am J Orthod Dentofacial Orthop* 2006;130:779-85.
  21. Kyrkanides S, Klambani M, Subtelny JD. Cranial base and facial skeleton asymmetries in individuals with unilateral cleft lip and palate. *Cleft Palate Craniofac J* 2000;37:556-61.
  22. Zemann W, Santler G, Kärcher H. Analysis of mid-face asymmetry in patients with cleft lip, alveolus and palate at the age of 3 months using 3D-COSMOS measuring system. *J Craniomaxillofac Surg* 2002;30:148-52.
  23. Kim YH, Sato K, Mitani H, Shimizu Y, Kikuchi M. Asymmetry of the sphenoid bone and its suitability as a reference for analyzing craniofacial asymmetry. *Am J Orthod Dentofacial Orthop* 2003;124:656-62.
  24. Suzuki H, Yamaguchi T, Furukawa M. Rhinologic computed tomographic evaluation in patients with cleft lip and palate. *Arch Otolaryngol Head Neck Surg* 1999;125:1000-4.
  25. Kolbenstvedt A, Aaløkken TM, Arctander K, Johannessen S. CT appearances of unilateral cleft palate 20 years after bone graft surgery. *Acta Radiol* 2002;43:567-70.
  26. Breitsprecher L, Fanghänel J, Metelmann HR, Mlynski G, Würfel F, Freise K, et al. The influence of the muscles of facial expression on the development of the midface and the nose in cleft lip and palate patients. A reflection of functional anatomy, facial esthetics and physiology of the nose. *Ann Anat* 1999;181:19-25.
  27. Markus AF, Precious DS. Effect of primary surgery for cleft lip and palate on mid-facial growth. *Br J Oral Maxillofac Surg* 1997;35:6-10.
  28. Markus AF, Delaire J, Smith WP. Facial balance in cleft lip and palate. II. Cleft lip and palate and secondary deformities. *Br J Oral Maxillofac Surg* 1992;30:296-304.
  29. Sade Hoefert C, Bacher M, Herberts T, Krimmel M, Reinert S, Göz G. 3D soft tissue changes in facial morphology in patients with cleft lip and palate and class III mal occlusion under therapy with rapid maxillary expansion and delaire facemask. *J Orofac Orthop* 2010;71:136-51.
  30. Carstens MH. Functional matrix cleft repair: principles and techniques. *Clin Plast Surg* 2004;31:159-89.