

A MEASURE OF GRAPH VULNERABILITY: SCATTERING NUMBER

ALPAY KIRLANGIÇ

Received 2 April 2001 and in revised form 20 August 2001

The *scattering number* of a graph G , denoted $sc(G)$, is defined by $sc(G) = \max\{c(G-S) - |S| : S \subseteq V(G) \text{ and } c(G-S) \neq 1\}$ where $c(G-S)$ denotes the number of components in $G-S$. It is one measure of graph vulnerability. In this paper, general results on the *scattering number* of a graph are considered. Firstly, some bounds on the *scattering number* are given. Further, *scattering number* of a binomial tree is calculated. Also several results are given about *binomial trees* and *graph operations*.

2000 Mathematics Subject Classification: 05C40, 05C85.

1. Introduction. In a communication network, vulnerability measures the resistance of the network to disruption of operation after the failure of certain stations or communication links. To measure vulnerability we have some parameters that are *toughness*, *binding number*, *vertex integrity*, and *scattering number* [5]. In this paper, we discuss the *scattering number* of a graph.

The *scattering number* of a graph G , denoted $sc(G)$, was introduced in [4]. Formally the *scattering number* is defined by

$$sc(G) = \max\{c(G-S) - |S| : S \subseteq V(G), c(G-S) \neq 1\}, \quad (1.1)$$

where $c(G-S)$ denotes the number of components in $G-S$. A cutset S of a graph G fulfilling $sc(G) = c(G-S) - |S|$ is said to be a *scattering set*. The problem “given a graph G , decide whether the scattering number is larger than zero” is NP-complete.

The scattering number of a graph is closely related to the toughness of a graph and to the existence of Hamilton cycles and paths. The *toughness* of a graph G , denoted $t(G)$, was defined by Chvátal [1]: for the complete graph K_n we have $t(K_n) = \infty$; if G is not complete, then $t(G) = \min\{|S|/c(G-S) : S \subseteq V(G), c(G-S) > 1\}$. A graph G is said to be *t-tough* if $t(G) \geq t$, that is, $|S| \geq tc(G-S)$ for any cutset S . It follows from the definitions that $t(G) \geq 1$ if and only if $sc(G) \leq 0$ for any graph G [3]. Moreover, Jung [4] calls the scattering number the “additive dual” of the toughness.

A Hamilton cycle in a graph G is a cycle containing every vertex of G . Similarly, a Hamilton path in a graph G is a path that contains every vertex of G . If a graph G has a Hamilton cycle, then $sc(G) \leq 0$; and if a graph G has a Hamilton path, then $sc(G) \leq 1$ [3].

The following theorem is given by Deogun et al. [3].

THEOREM 1.1. (a) *The scattering number of a graph G $sc(G) \geq sc(G-V') - |V'|$ holds for every subset $V' \subseteq V(G)$ in any graph G ;*

- (b) $\text{sc}(G) \geq \text{sc}(G - v) - 1$ for every vertex $v \in V(G)$ in any graph G ;
- (c) let G be a connected graph. Then there is a vertex $v \in V(G)$ such that $\text{sc}(G) \leq \text{sc}(G - v) - 1$;
- (d) for every connected graph G $\text{sc}(G) = \max_{v \in V(G)} \text{sc}(G - v) - 1$.

The *path cover number* of a graph G is the smallest number of disjoint paths covering the vertex set of G and is denoted by $\pi(G)$. For the next theorem a short proof is given in [3] and this theorem was also proven by Lehel without using order-theoretic tools [6].

THEOREM 1.2 (see [3]). *If G is cocomparability graph, then $\pi(G) = \max(1, \text{sc}(G))$.*

Now we give some definitions.

DEFINITION 1.3. The *connectivity* $\kappa = \kappa(G)$ of a graph G is the minimum number of vertices whose removal results in a disconnected or trivial graph.

DEFINITION 1.4. A subset X of V is called a covering of G if every edge of G has at least one end in X . A covering X is a *minimum covering* if G has no covering X' with $|X'| < |X|$. The *covering number*, $\alpha(G)$, is the number of vertices in a minimum covering of G .

DEFINITION 1.5. A subset X of V is called an *independent set* of G if no two vertices of X are adjacent in G . An independent set X is maximum if G has no independent set X' with $|X'| > |X|$. The *independence number* of G , $\beta(G)$, is the number of vertices in a maximum independent set of G .

In [Section 2](#), some bounds on the scattering number are given. [Section 3](#) gives several results about the scattering number and graph operations.

2. Bounds. Firstly, we give two theorems showing the relation between the toughness and the *scattering number*.

THEOREM 2.1. *If $t(G) \leq 0$, then $\text{sc}(G) \leq (\alpha(G)/t(G))(1 - t(G))$.*

PROOF. For any cutset S , we have

$$t(G) = \min \left\{ \frac{|S|}{c(G-S)} \right\} = \min \left\{ 1 - \frac{c(G-S) - |S|}{c(G-S)} \right\} = 1 - \max \left\{ \frac{c(G-S) - |S|}{c(G-S)} \right\}, \quad (2.1)$$

and so

$$\max \left\{ \frac{c(G-S) - |S|}{c(G-S)} \right\} = 1 - t(G). \quad (2.2)$$

Hence,

$$\frac{c(G-S) - |S|}{c(G-S)} \leq 1 - t(G) \Rightarrow c(G-S) - |S| \leq (1 - t(G))c(G-S). \quad (2.3)$$

On the other hand,

$$t(G) = \min \left\{ \frac{|S|}{c(G-S)} \right\} \Rightarrow \frac{|S|}{c(G-S)} \geq t(G), \quad c(G-S) \leq \frac{|S|}{t(G)} \quad (2.4)$$

for every cutset S . Since S is a cutset, we have $|S| \leq \alpha(G)$ and $c(G-S) \leq \alpha(G)/t(G)$. If $t(G) \leq 0$, then

$$(1-t(G))c(G-S) \leq \frac{\alpha(G)}{t(G)}(1-t(G)). \quad (2.5)$$

By (2.3) and (2.5),

$$c(G-S) - |S| \leq \frac{\alpha(G)}{t(G)}(1-t(G)), \quad t(G) \leq 0, \quad (2.6)$$

and so

$$sc(G) = \max_S \{c(G-S) - |S|\} \leq \frac{\alpha(G)}{t(G)}(1-t(G)). \quad (2.7)$$

The proof is completed. \square

THEOREM 2.2. *If $t(G) > 0$, then $sc(G) \leq \alpha(G)/t(G) - \kappa(G)$.*

PROOF. Consider any cutset S . Then $\kappa(G) \leq |S| \leq \alpha(G)$, obviously. Since $t(G) = \min\{|S|/c(G-S)\}$, we have $c(G-S) \leq |S|/t(G)$ and so

$$c(G-S) - |S| \leq \frac{|S|}{t(G)} - |S| \leq \frac{\alpha(G)}{t(G)} - \kappa(G). \quad (2.8)$$

Hence

$$sc(G) = \max_S \{c(G-S) - |S|\} \leq \frac{\alpha(G)}{t(G)} - \kappa(G). \quad (2.9)$$

The proof is completed. \square

Next, we give two theorems containing the relation between some graph parameters and the scattering number.

THEOREM 2.3. *If a graph G does not contain graph $2K_2$ as an induced subgraph, then*

$$sc(G) = \begin{cases} \beta(G) - \alpha(G), & \text{if } \kappa(G) = \alpha(G), \\ \beta(G) - \alpha(G) + 1, & \text{if } \alpha(G) = \kappa(G) + 1. \end{cases} \quad (2.10)$$

PROOF. If a graph G does not contain graph $2K_2$ as an induced subgraph, then we have $\alpha(G) = \kappa(G)$ or $\alpha(G) = \kappa(G) + 1$. That is, $|S|$ must be $\alpha(G)$ or $\alpha(G) - 1$.

$$\begin{aligned} \text{If } |S| = \alpha(G), & \quad \text{then } c(G-S) = \beta(G), \\ \text{If } |S| = \alpha(G) - 1, & \quad \text{then } c(G-S) = \beta(G). \end{aligned} \quad (2.11)$$

By (2.11) the proof is completed. \square




FIGURE 3.1

THEOREM 2.4. For any graph G , $\text{sc}(G) \leq \beta(G) - \kappa(G)$.

PROOF. For every $S \subset V(G)$, we have $c(G - S) \leq \beta(G)$. If S is a cutset, then $|S| \geq \kappa(G)$ and $c(G - S) - |S| \leq \beta(G) - \kappa(G)$. So

$$\max_S \{c(G - S) - |S|\} \leq \beta(G) - \kappa(G). \quad (2.12)$$

The proof is completed. \square

3. Binomial trees and scattering number. In this section, we consider the *binomial tree* B_n (Figure 3.1) (see [2]). The binomial tree B_n is an ordered tree defined recursively. The binomial tree B_0 consists of a single vertex. The binomial tree B_n consists of two binomial trees B_{n-1} that are linked together: the root of one is the leftmost child of the root of the other.

Now we give the scattering number of a binomial tree.

THEOREM 3.1. Let $n \geq 3$ be a positive integer. Then $\text{sc}(B_n) = 2^{n-2}$.

PROOF. In Figure 3.1, we call the vertex u top vertex of B_n . Let r be the number of removing vertices of B_n . If we remove top vertex u of B_n , then $B_{n-1}, B_{n-2}, \dots, B_1, B_0$ are components. Hence the number of components is n . Now if we remove top vertex of B_{n-1} , then we obtain the components $B_{n-2}, B_{n-3}, \dots, B_1, B_0$. Then we have $2(n-1)$ components. If we continue to remove the top vertex of each component, then we have two cases.

CASE 1. If $r = 2^i$ where $0 \leq i \leq n-1$, then the number of remaining components is exactly $(n-i)2^i$ where $0 \leq i \leq n-1$. Hence

$$\text{sc}(B_n) = \max_{0 \leq i \leq n-1} \{(n-i)2^i - 2^i\}. \quad (3.1)$$

CASE 2. If $2^{i-1} < r < 2^i$ where $2 \leq i \leq n-1$, then the number of remaining components is exactly $(n-(i-1))2^{i-1} + (r-2^{i-1})(n-(i+1))$. Hence

$$\text{sc}(B_n) = \max_{2 \leq i \leq n-1} \{(n-(i-1))2^{i-1} + (r-2^{i-1})(n-(i+1)) - r\}. \quad (3.2)$$

Now we can show that

$$\max_{2 \leq i \leq n-1} \{(n-(i-1))2^{i-1} + (r-2^{i-1})(n-(i+1)) - r\} \leq \max_{0 \leq i \leq n-1} \{(n-i)2^i - 2^i\}. \quad (3.3)$$

Then

$$\text{sc}(B_n) = \max_{0 \leq i < n} \{(n-i)2^i - 2^i\}. \quad (3.4)$$

The function $(n-i)2^i - 2^i$ takes its maximum value at $i = \lceil n - 1/\ln 2 - 1 \rceil$. It is obvious that $\lceil n - 1/\ln 2 - 1 \rceil = n - 2$ for every $n \geq 3$. Hence if we substitute this value in the function $(n-i)2^i - 2^i$, then the proof is completed. \square

DEFINITION 3.2. The tensor product of two graphs $G = (V(G), E(G))$ and $H = (V(H), E(H))$, denoted by $G \otimes H$, has the vertex set $V(G) \times V(H)$, the Cartesian product of $V(G)$ and $V(H)$, and an edge between vertices (x, y) and (u, v) , if and only if $\{x, u\} \in E(G)$ and $\{y, v\} \in E(H)$.

THEOREM 3.3. Let $m \geq 4$ and $n \geq 4$ be positive integers. Then

$$\text{sc}(B_m \otimes B_n) = \max \{2^{n-1}, 2^{m-1}\}. \quad (3.5)$$

PROOF. The graph $B_m \otimes B_n$ has the graphs B_m and B_n as subgraphs. We consider these graphs, respectively. Let r be the number of removing vertices from $B_m \otimes B_n$. Then we have two cases, depending on B_m or B_n .

CASE 1. Let u_1, u_2, \dots, u_{2^m} be the vertices of B_m and let v be the top vertex of B_n . If we remove the vertices $u_i v$ ($i = 1, 2, \dots, 2^m$), then the remaining components are $B_m \otimes B_{n-1}, B_m \otimes B_{n-2}, \dots, B_m \otimes B_1$, and $2^m B_0$. Now let the top vertex of B_{n-1} be v' . If we remove the vertices $u_i v'$ ($i = 1, 2, \dots, 2^m$), then we obtain the components $B_m \otimes B_{n-2}, \dots, B_m \otimes B_1$ and $2^m B_0$. If we continue to remove the vertices as mentioned above, then we obtain the following cases for r .

(a) If $r = 2^m 2^i$ where $0 \leq i \leq n-2$, then the number of components is $(n-i)2^i + 2^m 2^i$ and

$$\text{sc}(B_m \otimes B_n) = \max_{0 \leq i \leq n-2} \{2^i(n-i)\}. \quad (3.6)$$

(b) If $r = k2^m$ where $2^{i-1} < k < 2^i$ and $2 \leq i \leq n-2$, then the number of components is $(n-(i-1))2^{i-1} + (k-2^{i-1})(n-(i+1)) + 2^m k$ and

$$\text{sc}(B_m \otimes B_n) = \max_{2^{i-1} < k < 2^i, 2 \leq i \leq n-2} \{2^i + k(n-i-1)\}. \quad (3.7)$$

(c) If $r = k2^m$ and $2^{n-2} + 1 \leq k \leq 2^{n-1}$, then the number of components is $k2^m + 2(2^{n-1} - k)$ and

$$\text{sc}(B_m \otimes B_n) = \max_{2^{n-2} + 1 \leq k \leq 2^{n-1}} \{2(2^{n-1} - k)\}. \quad (3.8)$$

But we can show that

$$\begin{aligned} \max_{2^{i-1} < k < 2^i, 2 \leq i \leq n-2} \{2^i + k(n-i-1)\} &\leq \max_{0 \leq i \leq n-2} \{2^i(n-i)\}, \\ \max_{2^{n-2} + 1 \leq k \leq 2^{n-1}} \{2(2^{n-1} - k)\} &\leq \max_{0 \leq i \leq n-2} \{2^i(n-i)\}. \end{aligned} \quad (3.9)$$

Consequently, $\text{sc}(B_m \otimes B_n) = \max_{0 \leq i \leq n-2} \{2^i(n-i)\}$. The function $2^i(n-i)$ takes its maximum value at $i = \lceil n-1/\ln 2 \rceil$. It is obvious that $\lceil n-1/\ln 2 \rceil = n-1$ for every $n \geq 4$ and so

$$\text{sc}(B_m \otimes B_n) = 2^{n-1}. \quad (3.10)$$

CASE 2. Let v_1, v_2, \dots, v_{2^n} be the vertices of B_n and let u be the top vertex of B_m . If we remove the vertices uv_i ($i = 1, 2, \dots, 2^n$), then the remaining components are $B_{m-1} \otimes B_n, B_{m-2} \otimes B_n, \dots, B_1 \otimes B_n$, and $2^n B_0$. Now let the top vertex of B_{m-1} be u' . If we remove the vertices $u'v_i$ ($i = 1, 2, \dots, 2^n$), then we obtain the components $B_{m-2} \otimes B_n, \dots, B_1 \otimes B_n$ and $2^n B_0$. If we continue to remove the vertices as mentioned above, then we obtain the following cases for r .

(a) If $r = 2^n 2^i$ where $0 \leq i \leq m-2$, then the number of components is $(m-i)2^i + 2^n 2^i$ and

$$\text{sc}(B_m \otimes B_n) = \max_{0 \leq i \leq m-2} \{2^i(m-i)\}. \quad (3.11)$$

(b) If $r = k2^n$ where $2^{i-1} < k < 2^i$ and $2 \leq i \leq m-2$, then the number of components is $(m-(i-1))2^{i-1} + (k-2^{i-1})(m-(i+1)) + 2^i k$ and

$$\text{sc}(B_m \otimes B_n) = \max_{2^{i-1} < k < 2^i, 2 \leq i \leq m-2} \{2^i + k(m-i-1)\}. \quad (3.12)$$

(c) If $r = k2^n$ and $2^{m-2} + 1 \leq k \leq 2^{m-1}$, then the number of components is $k2^n + 2(2^{m-1} - k)$ and

$$\text{sc}(B_m \otimes B_n) = \max_{2^{m-2} + 1 \leq k \leq 2^{m-1}} \{2(2^{m-1} - k)\}. \quad (3.13)$$

But we can show that

$$\begin{aligned} \max_{2^{i-1} < k < 2^i, 2 \leq i \leq m-2} \{2^i + k(m-i-1)\} &\leq \max_{0 \leq i \leq m-2} \{2^i(m-i)\}, \\ \max_{2^{m-2} + 1 \leq k \leq 2^{m-1}} \{2(2^{m-1} - k)\} &\leq \max_{0 \leq i \leq m-2} \{2^i(m-i)\}. \end{aligned} \quad (3.14)$$

Consequently, $\text{sc}(B_m \otimes B_n) = \max_{0 \leq i \leq m-2} \{2^i(m-i)\}$. The function $2^i(m-i)$ takes its maximum value at $i = \lceil m-1/\ln 2 \rceil$. It is obvious that $\lceil m-1/\ln 2 \rceil = m-1$ for every $m \geq 4$ and so

$$\text{sc}(B_m \otimes B_n) = 2^{m-1}. \quad (3.15)$$

By (3.10) and (3.15) we have $\text{sc}(B_m \otimes B_n) = \max\{2^{m-1}, 2^{n-1}\}$.

The proof is completed. \square

DEFINITION 3.4. Let G_1 and G_2 be two graphs. The *union* $G = G_1 \cup G_2$ has $V(G) = V(G_1) \cup V(G_2)$ and $E(G) = E(G_1) \cup E(G_2)$. The *join* is denoted $V(G_1) + V(G_2)$ and consists of $V(G_1) \cup V(G_2)$ and all edges joining $V(G_1)$ with $V(G_2)$. For three or more disjoint graphs G_1, G_2, \dots, G_n , the *sequential join* $G_1 + G_2 + \dots + G_n$ is $(G_1 + G_2) \cup (G_2 + G_3) \cup \dots \cup (G_{n-1} + G_n)$.

THEOREM 3.5. *If n is an even number, then $\text{sc}(B_0 + B_1 + \cdots + B_n) = 4/3 - 2^n/3$.*

PROOF. To prove this theorem we have two cases.

CASE 1. If we remove the vertices of graphs B_1, B_3, \dots, B_{n-1} , then the remaining components are B_0, B_2, \dots, B_n and the number of removing vertices is $\sum_{i=1}^{n/2} |V(B_{2i-1})| = \sum_{i=1}^{n/2} 2^{2i-1}$. Moreover, we must delete $2^{2(i-1)}$ more vertices from each B_{2i} where $0 < i \leq n/2$ (except B_0). Hence $2 * 2^{2(i-1)}$ components are obtained from each B_{2i} where $0 < i \leq n/2$ (except B_0). Then the number of removing vertices is exactly

$$|S| = \sum_{i=1}^{n/2} 2^{2i-1} + \sum_{i=1}^{n/2} 2^{2(i-1)} \quad (3.16)$$

and the number of components is exactly

$$c((B_0 + B_1 + \cdots + B_n) - S) = 2 \sum_{i=1}^{n/2} 2^{2(i-1)} + 1. \quad (3.17)$$

So

$$\text{sc}(B_0 + B_1 + \cdots + B_n) = 1 - \frac{1}{4} \sum_{i=1}^{n/2} 2^{2i}. \quad (3.18)$$

CASE 2. If we remove the vertices of graphs B_0, B_2, \dots, B_n , then the remaining components are B_1, B_3, \dots, B_{n-1} and the number of removing vertices is $\sum_{i=0}^{n/2} |V(B_{2i})| = \sum_{i=0}^{n/2} 2^{2i}$. Moreover, we must delete 2^{2i-3} more vertices from each B_{2i-1} where $1 < i \leq n/2$ (except B_1). Hence $2 * 2^{2i-3}$ components are obtained from each B_{2i-1} where $1 < i \leq n/2$ (except B_1). Then the number of removing vertices is exactly

$$|S| = \sum_{i=1}^{n/2} 2^{2(i-1)} + \sum_{i=0}^{n/2} 2^{2i} \quad (3.19)$$

and the number of components is exactly

$$c((B_0 + B_1 + \cdots + B_n) - S) = 2 \sum_{i=2}^{n/2} 2^{2i-3} + 1. \quad (3.20)$$

So

$$\text{sc}(B_0 + B_1 + \cdots + B_n) = -5 - \sum_{i=2}^{n/2} 2^{2i}. \quad (3.21)$$

By (3.18) and (3.21), we have

$$\text{sc}(B_0 + B_1 + \cdots + B_n) = \max_i \left\{ 1 - \frac{1}{4} \sum_{i=1}^{n/2} 2^{2i}, -5 - \sum_{i=2}^{n/2} 2^{2i} \right\}. \quad (3.22)$$

Since $\sum a^t = a^t / (a - 1) + c(t)$ where $a \neq 1$, we have

$$\text{sc}(B_0 + B_1 + \cdots + B_n) = \max_i \left\{ \frac{4}{3} - \frac{2^n}{3}, \frac{1}{3} - \frac{4}{3} 2^n \right\} = \frac{4}{3} - \frac{2^n}{3}. \quad (3.23)$$

The proof is completed. \square

THEOREM 3.6. *If n is an odd number, then $\text{sc}(B_0 + B_1 + \cdots + B_n) = 2/3 - 2^n/3$.*

PROOF. The proof follows directly from [Theorem 3.5](#). □

REFERENCES

- [1] V. Chvátal, *Tough graphs and Hamiltonian circuits*, Discrete Math. **5** (1973), 215–228.
- [2] T. H. Cormen, C. E. Leiserson, and R. L. Rivest, *Introduction to Algorithms*, 4th ed., MIT Press, Massachusetts, 1990.
- [3] J. S. Deogun, D. Kratsch, and G. Steiner, *1-Tough cocomparability graphs are Hamiltonian*, Discrete Math. **170** (1997), 99–106.
- [4] H. A. Jung, *On a class of posets and the corresponding comparability graphs*, J. Combinatorial Theory Ser. B **24** (1978), 125–133.
- [5] D. Kratsch, T. Kloks, and H. Müller, *Measuring the vulnerability for classes of intersection graphs*, Discrete Appl. Math. **77** (1997), 259–270.
- [6] J. Lehel, *The path partition of cocomparability graphs*, unpublished, 1991.

ALPAY KIRLANGIÇ: DEPARTMENT OF MATHEMATICS, EGE UNIVERSITY, 35100 BORNOVA-İZMİR, TURKEY

E-mail address: alpkir@sci.ege.edu.tr

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br