

# Quantum affine wreath algebras

$$\begin{array}{c} \nearrow \\ \searrow \\ \circ \end{array} = \begin{array}{c} \nearrow \\ \searrow \\ \circ \end{array}, \quad \begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} - \begin{array}{c} \nearrow \\ \searrow \\ \searrow \\ \nearrow \end{array} = z \begin{array}{c} \uparrow \\ \bullet \\ \uparrow \\ \bullet \\ \uparrow \\ \bullet \\ \uparrow \end{array}$$

Alistair Savage  
University of Ottawa

Slides available online: [alistairsavage.ca/talks](http://alistairsavage.ca/talks)

Preprint: [arXiv:1902.00143](https://arxiv.org/abs/1902.00143) (joint with Daniele Rosso)

# Outline

**Goal:** Unify and generalize existing algebras by defining families of Hecke-like algebras depending on Frobenius algebras.

## Overview:

- 1 Strict monoidal categories and string diagrams
- 2 Warm up: symmetric groups, degenerate affine Hecke algebras
- 3 Frobenius algebras
- 4 Affine wreath product algebras
- 5 Quantum affine wreath product algebras

# Strict monoidal categories

A **strict monoidal category** is a category  $\mathcal{C}$  equipped with

- a bifunctor (the **tensor product**)  $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ , and
- a **unit object**  $\mathbb{1}$ ,

such that

- $(A \otimes B) \otimes C = A \otimes (B \otimes C)$  for all objects  $A, B, C$ ,
- $\mathbb{1} \otimes A = A = A \otimes \mathbb{1}$  for all objects  $A$ .

## Remark: Non-strict monoidal categories

In a (not necessarily strict) **monoidal category**, the equalities above are replaced by isomorphism, and we impose some **coherence conditions**.

Every monoidal category is monoidally equivalent to a strict one.

## $\mathbb{k}$ -linear monoidal categories

Fix a commutative ground ring  $\mathbb{k}$ .

A **strict  $\mathbb{k}$ -linear monoidal category** is a strict monoidal category such that

- each morphism space is a  $\mathbb{k}$ -module,
- composition of morphisms is  $\mathbb{k}$ -bilinear,
- tensor product of morphisms is  $\mathbb{k}$ -bilinear.

### The interchange law

The axioms of a strict monoidal category imply the **interchange law**: For  $A_1 \xrightarrow{f} A_2$  and  $B_1 \xrightarrow{g} B_2$ , the following diagram commutes:

$$\begin{array}{ccc} A_1 \otimes B_1 & \xrightarrow{1 \otimes g} & A_1 \otimes B_2 \\ f \otimes 1 \downarrow & \searrow f \otimes g & \downarrow f \otimes 1 \\ A_2 \otimes B_1 & \xrightarrow{1 \otimes g} & A_2 \otimes B_2 \end{array}$$

# String diagrams

Fix a strict monoidal category  $\mathcal{C}$ .

We will denote a morphism  $f: A \rightarrow B$  by:




The **identity map**  $1_A: A \rightarrow A$  is a string with no label:




We sometimes omit the object labels when they are clear or unimportant.

# String diagrams


Composition is **vertical stacking** and tensor product is **horizontal juxtaposition**:



The **interchange law** then becomes:



A morphism  $f: A_1 \otimes A_2 \rightarrow B_1 \otimes B_2$  can be depicted:



# Presentations of strict monoidal categories

One can give **presentations** of some strict  $\mathbb{k}$ -linear monoidal categories, just as for monoids, groups, algebras, etc.

**Objects:** If the objects are generated by some collection  $A_i, i \in I$ , then we have all possible tensor products of these objects:

$$\mathbb{1}, \quad A_i, \quad A_i \otimes A_j \otimes A_k \otimes A_\ell, \quad \text{etc.}$$

**Morphisms:** If the morphisms are generated by some collection  $f_j, j \in J$ , then we have all possible compositions and tensor products of these morphisms (whenever these make sense):

$$1_{A_i}, \quad f_j \otimes (f_i f_k) \otimes (f_\ell), \quad \text{etc.}$$

We then often impose some **relations** on these morphism spaces.

**String diagrams:** We can build complex diagrams out of our simple generating diagrams.

# The symmetric group category

Define a strict  $\mathbb{k}$ -linear monoidal category  $\mathcal{S}ym$  with one generating object  $\uparrow$  and denote

$$1_{\uparrow} = \uparrow$$

We have one generating morphism

$$\begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} : \uparrow \otimes \uparrow \rightarrow \uparrow \otimes \uparrow.$$

We impose the relations:

$$\begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} = \begin{array}{c} \uparrow \\ \uparrow \end{array}, \quad \begin{array}{c} \nearrow \\ \searrow \\ \searrow \\ \nearrow \end{array} = \begin{array}{c} \searrow \\ \nearrow \\ \searrow \\ \nearrow \end{array}.$$

Then

$$\text{End}_{\mathcal{S}ym}(\uparrow^{\otimes n}) = \mathbb{k}S_n$$

is the group algebra of the **symmetric group** on  $n$  letters.



# The symmetric group category

This monoidal presentation of  $S_n$  is very efficient! We only needed


- one generating morphism, and
- two relations,

to get **all** the symmetric groups.

Note that the “distant braid relation”

$$s_i s_j = s_j s_i, \quad |i - j| > 1$$

for simple transpositions follows for free from the interchange law:



# Degenerate affine Hecke algebras

The degenerate affine Hecke algebra  $H_n$  of type  $A$  is

$$\mathbb{k}[x_1, \dots, x_n] \otimes \mathbb{k}S_n$$

as a  $\mathbb{k}$ -module.

The factors  $\mathbb{k}[x_1, \dots, x_n]$  and  $\mathbb{k}S_n$  are subalgebras, and

$$\begin{aligned} s_i x_j &= x_j s_i, & j &\neq i, i+1, \\ s_i x_i &= x_{i+1} s_i - 1. \end{aligned}$$

# String diagrams

Obtain  $\mathcal{H}$  from  $Sym$  by adjoining one additional morphism (a **dot**)

$$\uparrow^{\circ} : \uparrow \rightarrow \uparrow$$

and one additional relation:

$$\begin{array}{c} \swarrow \circ \searrow \\ \times \\ \swarrow \searrow \circ \end{array} - \begin{array}{c} \swarrow \searrow \\ \times \\ \swarrow \circ \searrow \end{array} = \uparrow \uparrow .$$

Then

$$\text{End}_{\mathcal{H}}(\uparrow^{\otimes n}) = H_n$$

# Frobenius algebras

## Definition (symmetric Frobenius algebra)

An associative algebra  $A$  together with a linear **trace map**

$$\mathrm{tr}: A \rightarrow \mathbb{k}, \quad \mathrm{tr}(ab) = \mathrm{tr}(ba),$$

such that  $\ker \mathrm{tr}$  contains no nonzero left ideals.

## Example ( $\mathbb{k}$ )

$\mathbb{k}$  with  $\mathrm{tr} = \mathrm{id}_{\mathbb{k}}$ .

## Example ( $\mathbb{k}[x]/(x^k)$ )

$\mathbb{k}[x]/(x^k)$  with  $\mathrm{tr}(x^\ell) = \delta_{\ell, k-1}$ .

## Example (Matrix algebra)

Matrix algebras with the usual trace.

# Frobenius algebras: Examples

## Example (Group algebra)

If  $G$  is a finite group, then the **group algebra**  $\mathbb{k}G$  is a Frobenius algebra with

$$\mathrm{tr}(g) = \delta_{g,1_G}, \quad g \in G.$$

## Example (Hopf algebras)

Every f.d. Hopf algebra is a Frobenius algebra.

**From now on:**  $A$  is a symmetric Frobenius algebra with trace  $\mathrm{tr}$ .

**Remark:** Can actually work more generally, with graded Frobenius superalgebras (not necessarily symmetric).

# Wreath product algebras

The symmetric group  $S_n$  acts on  $A^{\otimes n}$  by **permutations**:

$$\pi(a_n \otimes \cdots \otimes a_1) = a_{\pi^{-1}(n)} \otimes \cdots \otimes a_{\pi^{-1}(1)},$$

## Wreath product algebra

The **wreath product algebra** is

$$\mathrm{Wr}_n(A) = A^{\otimes n} \otimes \mathbb{k}S_n$$

as  $\mathbb{k}$ -modules. Multiplication is determined by

$$(\mathbf{a} \otimes \pi)(\mathbf{b} \otimes \sigma) = \mathbf{a}\pi(\mathbf{b}) \otimes \pi\sigma.$$

# Wreath product algebras: Examples

Example ( $A = \mathbb{k}$ )

$$\mathrm{Wr}_n(\mathbb{k}) \cong \mathbb{k}S_n$$

Example ( $A = \mathrm{Cl}$ )

$\mathrm{Wr}_n(\mathrm{Cl})$  is the **Sergeev algebra**, which plays an important role in the projective representation theory of the symmetric group.

Example ( $A = \mathbb{k}G$ ,  $G = \mathbb{Z}/2\mathbb{Z}$ )

$\mathrm{Wr}_n(\mathbb{k}G)$  is the group algebra of the **hyperoctahedral group**, the Weyl group of type  $B$ .

Example ( $A = \mathbb{k}G$ ,  $G = \mathbb{Z}/r\mathbb{Z}$ )

$\mathrm{Wr}_n(\mathbb{k}G)$  is the group algebra of the **complex reflection group**  $G(r, 1, n)$ .

# The wreath product category

Define  $\mathcal{W}r(A)$  by adjoining to  $\mathcal{S}ym$  morphisms (tokens)

$$\uparrow \bullet a : \uparrow \rightarrow \uparrow, \quad a \in A,$$

subject to the relations

$$\uparrow \bullet \alpha a + \beta b = \alpha \uparrow \bullet a + \beta \uparrow \bullet b, \quad \begin{array}{c} \uparrow \\ \bullet a \\ \bullet b \end{array} = \uparrow \bullet ab, \quad \alpha, \beta \in \mathbb{k}, \quad a, b \in A,$$

(so  $A \mapsto \text{End}_{\mathcal{W}r(A)}(\uparrow)$ ,  $a \mapsto \uparrow \bullet a$  is an algebra homomorphism) and

$$\begin{array}{c} \nearrow \searrow \\ \bullet a \end{array} = \begin{array}{c} \nearrow \searrow \\ \bullet a \end{array}, \quad a \in A.$$

Then

$$\text{End}_{\mathcal{W}r(A)}(\uparrow^{\otimes n}) = \text{Wr}_n(A).$$



# Teleporters

Fix a basis  $B$  of  $A$ . The **dual basis** is

$$B^\vee = \{b^\vee \mid b \in B\} \quad \text{defined by} \quad \text{tr}(b^\vee c) = \delta_{b,c}, \quad b, c \in B.$$

**Exercise 1:**  $\sum_{b \in B} b \otimes b^\vee \in A \otimes A$  is independent of the basis  $B$ .

**Exercise 2:** For all  $a \in A$ , we have

$$\sum_{b \in B} ab \otimes b^\vee = \sum_{b \in B} b \otimes b^\vee a, \quad \sum_{b \in B} ba \otimes b^\vee = \sum_{b \in B} b \otimes ab^\vee.$$

Define the **teleporter**

$$\begin{array}{c} \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \end{array} := \sum_{b \in B} \begin{array}{c} \uparrow \quad \uparrow \\ \bullet \quad \bullet \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \end{array} b \quad b^\vee .$$

Then tokens “teleport” across teleporters:

$$\begin{array}{c} \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \end{array} a = \begin{array}{c} a \quad \uparrow \\ \bullet \quad \bullet \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \end{array}, \quad \begin{array}{c} \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \end{array} a = \begin{array}{c} \uparrow \quad \uparrow \\ \bullet \quad \bullet \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \end{array} a .$$

# Affine wreath product algebras

Define  $\mathcal{W}r^{\text{aff}}(A)$  by adjoining to  $\mathcal{W}r(A)$  the morphism (**dot**)

$$\uparrow_{\circ} : \uparrow \rightarrow \uparrow$$

and relations


$$\begin{array}{c} \nearrow \\ \circ \\ \searrow \end{array} - \begin{array}{c} \searrow \\ \circ \\ \nearrow \end{array} = \begin{array}{c} \uparrow \\ \bullet \\ \uparrow \end{array} \begin{array}{c} \uparrow \\ \bullet \\ \uparrow \end{array}, \quad \begin{array}{c} \uparrow \\ \bullet \\ \circ \\ \bullet \\ \uparrow \end{array} = \begin{array}{c} \uparrow \\ \circ \\ \bullet \\ \uparrow \end{array} a, \quad a \in A.$$

We define the **affine wreath product algebra** to be

$$\mathcal{W}r_n^{\text{aff}}(A) := \text{End}_{\mathcal{W}r^{\text{aff}}(A)}(\uparrow^{\otimes n}).$$

# Affine wreath product algebras

## Example ( $A = \mathbb{k}$ )

 and  $\text{Wr}_n^{\text{aff}}(\mathbb{k})$  is the **degenerate affine Hecke algebra**.

## Example ( $A = \text{Cl}$ , Clifford algebra)

$\text{Wr}_n^{\text{aff}}(\text{Cl})$  is the **affine Sergeev algebra**, aka the **degenerate affine Hecke–Clifford algebra**.

## Example ( $A = \mathbb{k}G$ )

$\text{Wr}_n^{\text{aff}}(\mathbb{k}G)$  is the **wreath Hecke algebra** (Wan–Wang).

## Example (Affine zigzag algebras)

When  $A$  is a certain zigzag algebra,  $\text{Wr}_n^{\text{aff}}(A)$  is related to imaginary strata for quiver Hecke algebras (Kleshchev–Muth).

# Hecke algebras

Fix  $z \in \mathbb{k}$ . Let  $\mathcal{H}(z)$  be the strict  $\mathbb{k}$ -linear monoidal category with one generating object  $\uparrow$ , generating morphisms

$$\begin{array}{c} \nearrow \searrow \\ \searrow \nearrow \end{array}, \begin{array}{c} \nearrow \nearrow \\ \searrow \searrow \end{array} : \uparrow \otimes \uparrow \rightarrow \uparrow \otimes \uparrow$$

and relations

$$\begin{array}{c} \nearrow \searrow \\ \searrow \nearrow \end{array} = \begin{array}{c} \uparrow \\ \uparrow \end{array}, \begin{array}{c} \nearrow \nearrow \\ \searrow \searrow \end{array} = \begin{array}{c} \uparrow \\ \uparrow \end{array}, \begin{array}{c} \nearrow \nearrow \\ \searrow \searrow \end{array} \begin{array}{c} \nearrow \searrow \\ \searrow \nearrow \end{array} = \begin{array}{c} \nearrow \searrow \\ \searrow \nearrow \end{array} \begin{array}{c} \nearrow \nearrow \\ \searrow \searrow \end{array}, \\ \begin{array}{c} \nearrow \searrow \\ \searrow \nearrow \end{array} - \begin{array}{c} \nearrow \nearrow \\ \searrow \searrow \end{array} = z \begin{array}{c} \uparrow \\ \uparrow \end{array} \quad (\text{skein relation}). \end{array}$$

Then

$$H_n(z) := \text{End}_{\mathcal{H}(z)}(\uparrow^{\otimes n})$$

is the **Iwahori–Hecke algebra** of type  $A_{n-1}$  (often  $z = q - q^{-1}$ ).

# Affine Hecke algebras

Define  $\mathcal{H}^{\text{aff}}(z)$  by adjoining to  $\mathcal{H}(z)$  the **invertible** morphism

$$\uparrow_{\circ} : \uparrow \rightarrow \uparrow$$

and relations

$$\begin{array}{c} \nearrow \\ \times \\ \searrow \\ \circ \end{array} = \begin{array}{c} \circ \\ \nearrow \\ \times \\ \searrow \end{array}, \quad \begin{array}{c} \nearrow \\ \times \\ \searrow \\ \circ \end{array} = \begin{array}{c} \nearrow \\ \circ \\ \times \\ \searrow \end{array}.$$

Then

$$H_n^{\text{aff}}(z) := \text{End}_{\mathcal{H}^{\text{aff}}}(\uparrow^{\otimes n})$$

is the **affine Hecke algebra** of type  $A_{n-1}$  (often  $z = q - q^{-1}$ ).

# Frobenius Hecke algebras

Define  $\mathcal{H}(A, z)$  by adjoining to  $\mathcal{H}(z)$  morphisms

$$\uparrow \bullet a : \uparrow \rightarrow \uparrow, \quad a \in A,$$

subject to the relations ( $\alpha, \beta \in \mathbb{k}, a, b \in A$ )

$$\uparrow \bullet \alpha a + \beta b = \alpha \uparrow \bullet a + \beta \uparrow \bullet b, \quad \begin{array}{c} \uparrow \bullet a \\ \uparrow \bullet b \end{array} = \uparrow \bullet ab,$$

$$\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array} - \begin{array}{c} \nwarrow \nearrow \\ \nearrow \searrow \end{array} = z \begin{array}{c} \uparrow \bullet \\ \uparrow \bullet \end{array},$$

$$\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array} \bullet a = \begin{array}{c} \nearrow \bullet \\ \nwarrow \searrow \end{array}, \quad \begin{array}{c} \nwarrow \nearrow \\ \nearrow \searrow \end{array} \bullet a = \begin{array}{c} \nwarrow \bullet \\ \nearrow \searrow \end{array}, \quad \begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array} \bullet a = \begin{array}{c} \nearrow \bullet \\ \nwarrow \nearrow \end{array}, \quad \begin{array}{c} \nwarrow \nearrow \\ \nearrow \searrow \end{array} \bullet a = \begin{array}{c} \nwarrow \nearrow \\ \nearrow \bullet \end{array}.$$

We call

$$H_n(A, z) := \text{End}_{\mathcal{H}(A, z)}(\uparrow^{\otimes n})$$

a Frobenius Hecke algebra.

# Frobenius Hecke algebras

Example ( $A = \mathbb{k}$ )

$H_n(\mathbb{k}, z)$  is an **Iwahori–Hecke algebra**.

Example ( $A = \mathbb{k}G$ ,  $G$  a cyclic group)

$H_n(\mathbb{k}G, z)$  is a **Yokonuma–Hecke algebra**.

Other choices of  $A$  yield **new** algebras.

# Quantum affine wreath algebras

Define  $\mathcal{W}r^{\text{aff}}(A, z)$  by adjoining to  $\mathcal{H}(A, z)$  the **invertible** morphism

$$\uparrow_{\circ} : \uparrow \rightarrow \uparrow$$

and relations

$$\begin{array}{c} \nearrow \\ \searrow \\ \circ \end{array} = \begin{array}{c} \circ \\ \nearrow \\ \searrow \end{array}, \quad \begin{array}{c} \nearrow \\ \searrow \\ \circ \end{array} = \begin{array}{c} \nearrow \\ \circ \\ \searrow \end{array}, \quad \begin{array}{c} \uparrow \\ \bullet a \\ \circ \end{array} = \begin{array}{c} \uparrow \\ \circ \\ \bullet a \end{array} .$$

We call

$$\mathcal{W}r_n^{\text{aff}}(A, z) := \text{End}_{\mathcal{W}r^{\text{aff}}(A, z)}(\uparrow^{\otimes n})$$

a **quantum affine wreath algebra**.

One could also call it a **affine Frobenius Hecke algebra**.



# Quantum affine wreath algebras

Example ( $A = \mathbb{k}$ )

$\mathrm{Wr}_n^{\mathrm{aff}}(\mathbb{k}, z)$  is an affine Hecke algebra.

Example ( $A = \mathbb{k}G$ ,  $G$  a cyclic group)

$\mathrm{Wr}_n^{\mathrm{aff}}(\mathbb{k}G, z)$  is an affine Yokonuma–Hecke algebra.

Other choices of  $A$  yield new algebras.

Example ( $A =$  zigzag algebra)

$\mathrm{Wr}_n^{\mathrm{aff}}(A, z)$  is a quantum analogue of affine zigzag algebras.

# Structure theory

One can prove many general structure theory results in a uniform way:

- Demazure operators (aka divided difference operators)
- Basis theorem:

$$\begin{aligned}\mathrm{Wr}_n^{\mathrm{aff}}(A) &\cong A^{\otimes n} \otimes \mathbb{k}[x_1, \dots, x_n] \otimes \mathbb{k}S_n \\ \mathrm{Wr}_n^{\mathrm{aff}}(A, z) &\cong A^{\otimes n} \otimes \mathbb{k}[x_1^{\pm 1}, \dots, x_n^{\pm 1}] \otimes H_n(z)\end{aligned}$$

as  $\mathbb{k}$ -modules.

- Center:

$$\begin{aligned}Z(\mathrm{Wr}_n^{\mathrm{aff}}(A)) &= (Z(A)^{\otimes n} \otimes \mathbb{k}[x_1, \dots, x_n])^{S_n} \\ Z(\mathrm{Wr}_n^{\mathrm{aff}}(A, z)) &= (Z(A)^{\otimes n} \otimes \mathbb{k}[x_1^{\pm 1}, \dots, x_n^{\pm 1}])^{S_n}\end{aligned}$$

- Jucys–Murphy elements
- Mackey theorem
- Cyclotomic quotients (basis theorem, Mackey theorem, etc.)

# Heisenberg categorification

Fix  $k \in \mathbb{Z}$ . The **Heisenberg category** (Khovanov, Mackaay–S., Brundan) is defined by adjoining to the degenerate affine Hecke category  $\mathcal{H}$  an object

$\downarrow$

and morphisms and relations so that

- $\uparrow$  is right dual to  $\downarrow$ ,
- we have an isomorphism

$$\begin{aligned}\uparrow \otimes \downarrow &\cong \downarrow \otimes \uparrow \oplus \mathbb{1}^{\oplus k} && \text{(when } k \geq 0\text{),} \\ \uparrow \otimes \downarrow \oplus \mathbb{1}^{\oplus(-k)} &\cong \downarrow \otimes \uparrow && \text{(when } k \leq 0\text{)}\end{aligned}$$

(the **inversion relation**).

Acts on modules for degenerate cyclotomic Hecke algebras, categorifies the Heisenberg algebra.

# Heisenberg categorification

We can now repeat this with our (quantum) affine wreath categories!

We get:

- quantum Heisenberg category (Licata–S., Brundan–S.–Webster)
- Frobenius Heisenberg category (Rosso–S., S.)
- quantum Frobenius Heisenberg category (Brundan–S.–Webster, work in progress)

These act on modules for the corresponding cyclotomic quotients.

Can also define an odd quantum Frobenius Heisenberg category...