

Article

New 3H-Indole Synthesis by Fischer's Method. Part I.

Sami Sajjadifar ^{1,2,*}, Hooshang Vahedi ¹, Abdolhossien Massoudi ¹ and Omid Louie ¹

¹ Department of Chemistry, Payame Noor University (PNU), P O Box 91735-433, Mashhad, Iran; E-Mails: hooshangvahedi@yahoo.co.uk (H.V.); Massoudi_h@pnu.ac.ir (A.M.); o_louie2001@yahoo.com (O.L.)

² Department of Chemistry, Payame Noor University (PNU), Ilam, Iran

* Author to whom correspondence should be addressed; E-Mail: ss_sajjadifar@yahoo.com; Tel.: +98 8412228316.

Received: 6 February 2010; in revised form: 12 March 2010 / Accepted: 19 March 2010 /

Published: 8 April 2010

Abstract: Methyl indolenines (**4a-c**) and (**5a-c**) were prepared in high yield by a Fischer indole synthesis reaction of *o,m*-tolylhydrazine hydrochlorides (**1a-b**) with isopropyl methyl ketone (**2**) and 2-methylcyclohexanone (**3**) in acetic acid at room temperature. *o,p*-Nitrophenylhydrazines (**1c-d**) were reacted with 2-methylcyclohexanone (**3**) in acetic acid at reflux to give nitroindolenines (**5d-e**), while the attempted reactions of *o,p*-nitrohydrazines with isopropyl methyl ketone (**2**) in acetic acid were not successful. Compounds (**1c-d**) were reacted with isopropyl methyl ketone (**2**) in acetic acid/HCl to give 2,3,3-trimethyl-5-nitro-indolenine (**4e**) and 2,3,3-trimethyl-7-nitroindolenine (**4d**).

Keywords: 3H-indole; indolenine; Fischer's synthesis method; acetic acid


Introduction

The Fischer indole synthesis is a chemical reaction that produces an aromatic heterocyclic indole from a (substituted) phenylhydrazine and an aldehyde or ketone under acidic conditions. The reaction was discovered in 1883 by Hermann Emil Fischer [1]. Today antimigraine drugs of the tryptan class are often synthesized by this method. The choice of acid catalyst is very important. Bronsted acids such as HCl, H₂SO₄, polyphosphoric acid and *p*-toluenesulfonic acid have been used successfully,

while. Lewis acids such as boron trifluoride, zinc chloride, iron chloride, and aluminium chloride are also useful catalysts [2–4].

The reaction of a (substituted) phenylhydrazine with an aldehyde or ketone initially forms a phenylhydrazone which isomerizes to the respective enamine (or 'ene-hydrazine'). After protonation, a cyclic [3,3]-sigmatropic rearrangement occurs producing an imine. The resulting imine forms a cyclic aminoacetal (or aminal), which under acid catalysis that eliminates NH_3 , resulting in the energetically favorable aromatic indole. The accepted mechanism for Fischer's synthesis was suggested by Robinson [2,3]. According to Robinson mechanism, this reaction has three steps: (a) tautomeric conversion of hydrazone to enehydrazine, (b) carbon-carbon bond formation, (c) cyclization with ammonia, elimination and finally indole synthesis. Some researchers [5] believed that the carbon-carbon bond formation step is the most important and rate-determining step. This step can control the amount of indoles. The presence of different ketones for indole synthesis has been investigated. Isotopic labeling studies show that the aryl nitrogen (N1) of the starting phenylhydrazine is incorporated into the resulting indole [6–7].

Scheme 1. The Fischer reaction for indolenine synthesis.



	R ₁	R ₂	R ₃
a)	H	H	Me
b)	H	Me	H
c)	NO ₂	H	H
d)	H	H	NO ₂

	R ₁	R ₂	R ₃	R ₄
a)	H	H	Me	H
b)	H	Me	H	H
c)	H	H	H	Me
d)	NO ₂	H	H	H
e)	H	H	NO ₂	H

Results and Discussion

As mentioned in the Introduction, the use of different ketones for indole synthesis has been investigated, but so far, the effects of substituent groups on the phenyl hydrazine benzene ring on the Fischer indole synthesis have not been reported. Therefore, the presence of methyl and nitro groups in different phenyl hydrazines and the use of isopropyl methyl ketone (**2**) and 2-methylcyclohexanone (**3**) resulted in a new synthesis of indolenines in which the yield significantly high, mainly a single product is obtained and it can be carried out under simple conditions.

$^1\text{H-NMR}$ spectrum of indolenine (**4a**) shows a singlet at $\delta = 1.16$ ppm corresponding to the two methyl groups at C-3, a singlet methyl group for C-2 at $\delta = 2.24$ ppm, and a methyl group singlet resonance for Ar-CH₃ at $\delta = 2.1$ ppm. In the UV spectrum of (**4a**), two absorption bands of 217 and 262 nm were seen. After acid addition, position shifts in the band maxima was noticed which are the result of presence of the non-bonding pair electrons in the H⁺ reaction. Moreover, the presence of the donating electron methyl groups in suitable positions supported this bathochromic effect.

m-Tolylhydrazine hydrochloride (**1b**) reacted at room temperature with isopropyl methyl ketone (**2**) at the presence of glacial acetic acid to afford two isomers - 2,3,3,4-tetramethylindolenine (**4c**) and 2,3,3,6-tetramethylindolenine (**4b**) - in 88% yield. Our efforts to separate the two isomers through TLC and column chromatography in different solvents were fruitless.

The $^1\text{H-NMR}$ spectrum of the two isomers displays two singlets at $\delta = 1.25$ and $\delta = 1.35$ ppm corresponding to the C-3 methyl hydrogens, two methyl group of C-2 at $\delta = 2.33$ and $\delta = 2.37$ ppm, a Ar-CH₃ methyl group singlet at $\delta = 2.19$ and in the aromatic region, a doublet at $\delta = 6.84$ ppm and a singlet at $\delta = 7.1$. In the IR spectrum, a C=N stretching vibration is seen at $1,700\text{ cm}^{-1}$. In the UV spectrum of this mixture, two absorption bands at 218 and 259 nm were seen and a bathochromic effect was noticed after acid addition.

The $^1\text{H-NMR}$ spectrum of indolenine (**5a**) showed a methyl group singlet at $\delta = 1.15$ ppm which provides strong support for the cyclization and creation of indolenine (**5a**). In the IR spectrum, a C=N stretching vibration was observed at $1,700\text{ cm}^{-1}$, which indicates the absence of the N-H bond. In the UV spectrum of (**5a**) two absorption bands at 212 and 263 nm were observed, which after acid addition shifted position. This indicates the presence of non-bonding pair electrons in the reaction with H⁺.


m-Tolylhydrazine hydrochloride (**1b**) was also reacted at room temperature with 2-methylcyclohexanone (**3**) in the presence of glacial acetic acid. After extraction, the products were passed through a short silica gel column. Two isomers, 4a,5- and 4a,7-dimethyl-1,2,3,4-tetrahydro-4aH-carbazole (**5b-c**), were obtained. The efforts to separate these products were fruitless. The $^1\text{H-NMR}$ spectrum of the mixture of isomers indicated two singlet for the C-4a methyl groups at $\delta = 1.14$ and $\delta = 1.24$, which proves the cyclization and formation of the indolenines (**5b-c**). The singlet signals of the methyl groups in the two isomers overlap at $\delta = 2.21$. In the IR spectrum the C=N stretching vibration was observed at $1,695\text{ cm}^{-1}$. In the UV spectrum the maximum absorptions were noticed at 214 and 259 nm which after acid addition, a shift in the position of the band maxima was also seen, which is indicative of the presence of non-bonding pair electrons in the reaction with H⁺ (bathochromic effect).

p-Nitrophenylhydrazine hydrochloride (**1d**) and isopropyl methyl ketone (**2**) were reacted in the presence of acetic acid and the mixture was refluxed for 1.5 h. The product, 2,3,3-trimethyl-5-nitroindolenine (**4d**), was obtained in very low yield (10%). During the long reaction under reflux conditions, the instability and sensitivity of the products led to their decomposition, oxidation and polymerization. The reaction was also carried out in a binary mixture of acetic acid and hydrochloric acid for 4 h. After extraction, the product (**4e**) was obtained in 30 % yield. The $^1\text{H-NMR}$ spectrum of indolenine (**4e**) showed a singlet signal for the two C-3 methyl groups at $\delta = 1.35$ and a methyl group singlet for C-2 at $\delta = 1.87$. In the aromatic region, three hydrogens were observed.

Conclusions


It is shown that by switching from –OMe to –NO₂, either the time of the reaction will be longer or reaction conditions will be harsher, so when there is an electron-withdrawing group on the benzene ring in the phenylhydrazine, the efficiency is lower and the reactions are much more difficult. This research demonstrates that Robinson's proposed mechanism is correct. The effects of the studied groups in the indolenine formation reaction are summarized in Tables 1-2.

Table 1. Reactions with isopropyl methyl ketone (2).



R	Condition	Time	Acid	Y %
p-OMe	RT	20 min	AcOH	85
p,o-OMe	RT	2 h	AcOH	>90
o,m,p-Cl	reflux	0.5 h	AcOH	>90
o,p-NO ₂	reflux	90 h	AcOH	9
		4 h	AcOH, HCl	31

Table 2. Reactions with 2-methylcyclohexanone (3).



R	Condition	Time	Acid	Y %
p-OMe	RT	20 min	AcOH	>90
p,o-OMe	RT	1.5 h	AcOH	>90
o,m,p-Cl	reflux	0.5 h	AcOH	>90
o,p-NO ₂	reflux	24 h	AcOH	53

Experimental

General

Melting points were recorded on a Philip Harris C4954718 apparatus and are uncorrected. ¹H- and ¹³C-NMR spectra were recorded on a Bruker Avance AQS 75 MHz spectrometer. Chemical shifts δ are in parts per million (ppm) measured in CDCl₃ as solvent and relative to TMS as the internal standard. Infrared spectra were recorded on a Thermo Nicolet-Nexus 670 FTIR instrument and elemental analyses were carried out on an Exeter analytical model CE440 C, H and N elemental analyzer.

2,3,3,5-Tetramethylindolenine (4a)

p-Tolylhydrazine hydrochloride (**1a**, 0.25 g, 1.62 mmol) and isopropyl methyl ketone (**3**, 0.14 g, 1.62 mmol) were added to glacial acetic acid (2 g, 0.03 mol). The mixture was refluxed for 2.25 h with stirring. TLC indicated the end of the reaction and formation of the product. The mixture was cooled and neutralized with 1 M NaOH then diluted with water (100 mL) and extracted with CDCl₃ (3 × 100 mL). Following the drying of the organic layer over Na₂SO₄, the solvent was removed by means of evaporation and the residue was passed through a short silica gel column for further purification. A red, viscous oil of 2,3,3,5-tetramethylindolenine (**4a**) was obtained. TLC: R_F 0.22 on silica gel eluting with PhCH₃-EtOAc (9:1); UV: (EtOH) λ_{max} (nm) 217 (H, 2.80) 262 (H, 1.557), (EtOH, HCl) 207 (H, 1.706), 236 (H, 1.487), 289 (H, 1.310); IR (cm⁻¹): 2940, 1680, 1570, 1450, 1363, 1190, 810; ¹H-NMR δ (ppm): 1.16 (s, 6H, 2×CH₃), 2.1 (s, 3H, Ar-CH₃), 2.24 (s, 3H, N=C-CH₃), 6.7-6.9 (b, 1H, Ar-H), 6.87 (s, 1H, Ar-H), 7.1-7.2 (b, 1H, Ar-H); ¹³C-NMR δ (ppm): 19.9, 24.6, 25.1, 37.8, 121.9, 127.6, 129.3, 136.6, 148.8, 152.7, 164.6; C₁₂H₁₅N, Mol. Wt: 173.2542; Elem. Anal.: C, 83.19; H, 8.73; N, 8.08.

Reaction of (1b) with ketone (2) to produce indolenines (4b) and (4c)

m-Tolylphenylhydrazine hydrochloride (**1b**, 0.3 g, 1.89 mmol) and isopropyl methyl ketone (**2**, 0.17 g, 1.98 mmol) were added to glacial acetic acid (3 g, 0.05 mol) and stirred for 2 h. TLC indicated the end of reaction and formation of the product. The mixture was then cooled, neutralized with 1 M NaOH, diluted with water (100 mL) and finally extracted with CDCl₃ (3 × 100 mL). Following the drying of the organic layer over Na₂SO₄, the solvent was removed by evaporation and the residue passed through a short column of silica gel for further purification. Attempts to separate the two isomers were not successful, but one of the isomers [2,3,3,6-tetramethylindolenine (**4b**)] was isolated and separated by preparative TLC eluting with CH₂Cl₂-CH₃COCH₃ (20:1) and the data of another isomer inferred by comparing with the data of the two isomer mixture.

2,3,3,6-Tetramethylindolenine (**4b**): TLC: R_F 0.3 on silica gel eluting with CH₂Cl₂-CH₃COCH₃ (20:1); UV: (EtOH) λ_{max} (nm) 219 (H, 2.292) 261 (H, 0.684), (EtOH, HCl) 210 (H, 1.513) 227 (H, 1.088) 280 (H, 0.625); IR (cm⁻¹): 2900, 1700, 1600, 1565, 1450, 1425, 1340, 1250, 1130, 800; ¹H-NMR: 1.25 (s, 6H, 2×CH₃), 2.19 (s, 3H, Ar-CH₃), 2.33 (s, 3H, N=C-CH₃), 6.7-6.9 (b, 2H, Ar-H), 7.1 (s, 1H, Ar-H); ¹³C-NMR: 19.9, 24.3, 25.1, 37.5, 123.6, 127.3, 127.4, 136.9, 149.8, 151.7, 164.6; C₁₂H₁₅N, Mol. Wt: 173.2542; Elem. Anal.: C, 83.19; H, 8.73; N, 8.08.

2,3,3,4-Tetramethylindolenine (**4c**): TLC: R_F 0.1-0.35 on silicagel eluting with CH₂Cl₂-CH₃COCH₃ (20:1); UV: (EtOH) λ_{max} (nm) 219 (H, 2.802) 259 (H, 1.007), (EtOH, HCl) 209 (H, 1.658) 235 (H, 1.234) 281 (H, 0.971); IR (cm⁻¹) 2900, 1680-1700, 1570, 1420, 1350, 1220-1250, 800; ¹H-NMR: 1.35 (s, 6H, 2×CH₃), 2.19 (s, 3H, Ar-CH₃), 2.37 (s, 3H, N=C-CH₃), 6.7-7.2 (b, 3H, Ar-H); ¹³C-NMR: 18.4, 19.9, 25.4, 31.0, 119.0, 127.2, 127.3, 135.3, 142.9, 151.7, 164.6; C₁₂H₁₅N, Mol. Wt: 173.2542; Elem. Anal.: C, 83.19; H, 8.73; N, 8.08.

4a,6-Dimethyl-1,3,3,4-tetrahydro-4aH-carbazole (5a)

p-Tolylydrazine hydrochloride (**1a**, 0.3 g, 1.89 mmol) and 2-methylcyclohexanone (**3**, 0.21 g, 1.89 mmol) were added to glacial acetic acid (3 g, 0.05 mol). The mixture was refluxed for 1.5 h with stirring until TLC indicated the end of the reaction and formation of product. The mixture was cooled and neutralized with 1 M NaOH, then diluted with water (100 mL) and extracted with CDCl₃ (3 × 100 mL). After drying the organic layer over Na₂SO₄, the solvent was removed by evaporation and the residue was passed through a short silica gel column for further purification. A red, viscous oil of 4a,6-dimethyl-1,2,3,4-tetrahydro-4aH-carbazole (**5a**, 0.32 g, 85%) was obtained. TLC: R_F 0.2 on silica gel eluting with PhCH₃-EtOAc (9:1); UV: (EtOH) λ_{max} (nm) 212 (H, 1.624) 263 (H, 0.709), (EtOH, HCl) 204 (H, 1.080) 237 (H, 0.635), 290(H, 1.603); IR (cm⁻¹): 2900, 1700, 1570, 1440, 1360, 1170, 810, 740. ¹H-NMR: 1.15 (s, 6H, 2×CH₃), 2.2 (s, 6H, Ar-CH₃ and N=C-CH₃), 1.1-2.8 (m, 8H, aliphatic ring hydrogens), 6.7-6.9 (b, 2H, Ar-H), 7.05-7.25 (b, 1H, Ar-H); ¹³C-NMR: 21.9, 22.9, 24.6, 25.4, 31.2, 34.1, 54.1, 122.0, 127.6, 131.3, 136.7, 143.4, 150.8, 187.3; C₁₄H₁₇N, Mol. Wt: 199.2915; Elem. Anal.: C, 84.37; H, 8.60; N, 7.03;

Reaction of (1b) with ketone (3) to produce a mixture of indolenines (5b) and (5c)

m-Tolylydrazine hydrochloride (**1b**, 0.3 g, 1.89 mmol) and 2-methylcyclohexanone (**3**, 0.21 g, 1.89 mmol) were added to glacial acetic acid (3 g, 0.05 mol). The mixture was refluxed for 2 h with stirring until TLC indicated the end of the reaction and formation of the product. The mixture was cooled and neutralized with 1 M NaOH then diluted with water (100 mL) and extracted with CDCl₃ (3 × 100 mL). Following drying the organic layer over Na₂SO₄, the solvent was removed by evaporation and the residue passed through short silica gel column for further purification. A red viscous oil that it is a mixture of the two isomers (**5b**) and (**5c**) is obtained. TLC: R_F 0.1-0.3 on silica gel eluting with CH₂Cl₂: CH₃COCH₃ (20: 1); UV (EtOH) λ_{max} (nm) 214 (H, 2.398) 259 (H, 0.157), (EtOH, HCl) 206 (H, 1.813) 235(H, 0.378) 281 (H, 0.150); IR (cm⁻¹): 2840-2910, 1695, 1570, 1430, 1365, 1215, 1125, 965, 800, 780, 743; ¹H-NMR: 1.14 (s, 3H), 1.1-2.8 (m, aliphatic ring), 1.24 (s, 3H), 2.21 (s, 3H), 6.4- 7.2 (m, Ar-H); ¹³C-NMR: 18.4, 21.9, 22.9, 23.2, 24.3, 25.4, 31.2, 34.1, 47.3, 53.8, 119.1, 123.7, 127.2, 127.4, 129.4, 133.6, 136.9, 137.3, 140.5, 153.7, 187.3; C₁₄H₁₇N, Mol. Wt: 199.2915; Elem. Anal.: C, 84.37; H, 8.60; N, 7.03.

4a-Methyl-6-nitro-1,2,3,4-tetrahydro-4aH-carbazole (5e)

p-Nitrophenylhydrazine (**1d**, 0.29 g, 1.89 mmol) and 2-methylcyclohexanone (**3**, 0.21 g, 1.89 mmol) were added to glacial acetic acid (3 g, 0.05 mol). The mixture was refluxed for 24 h with stirring until TLC indicated the end of the reaction and formation of product. The mixture cooled and neutralized with NaOH 1 M then diluted with water (100 mL) and extracted with CDCl₃ (3 × 100 mL). After drying the organic layer over Na₂SO₄, the solvent was removed by evaporation. A yellow oil of the product (**5e**) (0.24 g, 53%) was obtained and we used a short column of silica gel for further purification. TLC: R_F 0.44 on silicagel eluting with PhCH₃-CH₃COCH₃ (10:2); UV: (EtOH) λ_{max} (nm) 221 (H, 2.389) 304 (H, 2.413), (EtOH, HCl): 219 (H, 2.338) 302 (H, 2.370); IR: (cm⁻¹) 2840-2920, 1880, 1708, 1585, 1555, 1500, 1440, 1320, 1230, 1096, 1046, 970, 830, 730; ¹H-NMR: 1.32 (s, 3H),

1.1-2.8 (m, 8H, aliphatic ring), 7.3-7.5 (b, 1H, Ar-H), 7.85 (s, 1H, Ar-H), 7.9-8.1 (b, 1H, Ar-H); ^{13}C -NMR: 21.9, 22.9, 25.4, 31.2, 34.1, 52.8, 119.6, 123.0, 124.7, 144.4, 146.7, 159.9, 187.3; $\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_2$; Mol. Wt. 230.2625; Elem. Anal.: C, 67.81; H, 6.13; N, 12.17; O, 13.90.

4a-Methyl-8-nitro-1,2,3,4-tetrahydro-4aH-cabazole (5d)

A mixture of *o*-nitrophenylhydrazine (**1c**, 0.29 g, 1.89 mmol), 2-methylcyclohexanone (**3**, 0.21 g, 1.89 mmol) and glacial acetic acid (3 g, 0.05 mol) was refluxed for 24 h with stirring until TLC indicated the end of the reaction. The mixture was cooled and neutralized with NaOH 1 M then diluted with water (100 mL) and extracted with CDCl_3 (3 \times 100mL). After drying the organic layer over Na_2SO_4 , the solvent was removed by evaporation. A red oil of (**5d**) (0.23 g, 51%) was obtained and a short column of silica gel was used for further purification. TLC: R_F 0.2 on silica gel eluting with PhCH_3 - CH_3COCH_3 (9:1); UV: (EtOH) λ_{max} (nm) 233 (H, 1.92), 251 (H, 2.03), 440 (H, 0.66) (EtOH, HCl), 233 (H, 1.87), 251 (H, 1.94), 440 (H, 0.64); IR (cm^{-1}): 2820-2940, 1690, 1600, 1560, 1490, 1400, 1320, 1260, 1020, 740; ^1H -NMR: 1.3 (s, 3H), 1.1-2.8 (m, 8H, aliphatic ring) 6.5-7.8 (b, 3H, Ar-H); ^{13}C -NMR: 21.9, 22.9, 25.4, 31.2, 34.1, 52.8, 119.6, 128.0, 135.6, 141.7, 143.9, 144.4, 187.3; $\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_2$; Mol. Wt. 230.2625; Elem. Anal.: C, 67.81; H, 6.13; N, 12.17; O, 13.90.

2,3,3-Trimethyl-5-Nitroindolenine (4d)

A mixture of *o*-nitrophenylhydrazine (**1c**, 0.29 g, 1.89 mmol), 2-methylcyclohexanone (**3**, 0.21 g, 1.89 mmol), and glacial acetic acid (3 g, 0.05 mol) was refluxed for 24 h with stirring until TLC indicated the end of the reaction. The mixture was cooled and neutralized with 1 M NaOH, then diluted with water (100 mL) and extracted with CDCl_3 (3 \times 100 mL). After drying the organic layer over Na_2SO_4 , the solvent was removed by evaporation. A red oil of (**5d**) (0.23 g, 51%) was obtained and we used a short column of silica gel for additional purification. TLC: R_F 0.2 on silica gel eluting with PhCH_3 - CH_3COCH_3 (9:1); UV: (EtOH) λ_{max} (nm) 233 (H, 1.92), 251 (H, 2.03), 440 (H, 0.66): (EtOH, HCl), 233 (H, 1.87), 251 (H, 1.94), 440 (H, 0.64); IR (cm^{-1}): 2820-2940, 1690, 1600, 1560, 1490, 1400, 1320, 1260, 1020, 740. ^1H -NMR: 1.3 (s, 3H), 1.1-2.8 (m, 8H, aliphatic ring), 6.4-6.6 (b, 3H, Ar-H); ^{13}C -NMR δ : 19.9, 25.1, 36.5, 119.6, 122.7, 122.9, 146.6, 153.7, 157.9, 164.6; $\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_2$; Mol. Wt: 204.2252; Elem. Anal.: C, 64.69; H, 5.92; N, 13.72; O, 15.67.

Acknowledgements

The authors are grateful to the Payame Noor University (PNU) of Mashhad for financial support of this work.

References and Notes

1. Fischer, E.; Jourdan, F. Indole Synthesis. *Chem. Ber.* **1883**, *16*, 2241.
2. Robinson, B. The Fischer indole synthesis. *Chem. Ber.* **1963**, *63*, 373–401.
3. Robinson, B. Studies on the Fischer indole synthesis. *Chem. Rev.* **1969**, *69*, 227.
4. Van Orden, R.B.; Lindwell, H.G. Indole. *Chem. Rev.* **1942**, *30*, 69–96.

5. Robert E. Lyle and Leonidas Skarlos on the direction of cyclization of unsymmetrical ketone phenyl hydrazones in the Fischer indole synthesis. *Chem. Comm.* **1966**, 644.
6. Allen, C.F.H.; Wilson, C.V. The use of N¹⁵ as a tracer element in chemical reactions. The mechanism of the Fischer indole synthesis. *J. Am. Chem. Soc.* **1943**, 65, 611–612.
7. Schiess, P.; Grieder, A. Synthese und reaktivitaet von n-phenyl-N, N'-dimethyl-enhydrazinen. *Tetrahedron Lett.* **1969**, 10, 2097–2100.
8. Yamamoto, H. 1-Acylindoles. III. Novel synthesis of 9-acyltetrahydrocrbazole and 5-acyl-p-carboline derivatives. *J. Org. Chem.* **1967**, 32, 3693–3695.
9. Xu, D.Q.; Yang, W.L.; Luo, S.P.; Wang, B.T.; Wu, J.; Xu, Z.Y. Fischer indole synthesis in brønsted acidic ionic liquids: A green, mild, and regiospecific reaction system. *Eur. J. Org. Chem.* **2007**, 6, 1007–1012.
10. Illy, H.; Funderburk, L. Fischer indole synthesis. Direction of cyclization of isopropylmethyl ketone phenylhydrazone. *J. Org. Chem.* **1968**, 33, 4283–4285.
11. Grandberg, I.I.; Sorokin, V.I. The mode of cyclisation of the arylhydrazones and o-phenyl ethers of the oximes of asymmetric ketones under the conditions of the Fischer reaction. *Russ. Chem. Rev.* **1974**, 43, 115–128.
12. Rashidi, A.; Afghan, A.; Baradarani, M.M.; Joule, J.A. The Synthesis of 4-(3,3-Dimethyl-3H-pyrrolo[2,3-f]quinolin-2-yl)pyrazoles and 4-(3,3-Dimethyl-3H-pyrrolo[3,2-h]quinolin-2-yl)pyrazoles. *J. Heterocycl. Chem.* **2009**, 46, 428.
13. Humphrey, G.R.; Kueth, J.T. Practical methodologies for the synthesis of indoles. *Chem. Rev.* **2006**, 106, 2875–2911.
14. Miller, F.M.; Schinske, W.N. Direction of cyclization in the Fischer indole synthesis. Mechanistic considerations. *J. Org. Chem.* **1978**, 43, 3384–3388.
15. Venkatesh, C.; Singh, P.P.; Ila, H.; Junjappa, H. Highly diastereoselective [3+2] cyclopenta[b]annulation of indoles with 2-arylcyclopropyl ketones and diesters. *Eur. J. Org. Chem.* **2006**, 2006, 5378–5386.

Sample Availability: Not available.

© 2010 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).