


AAD Annual Meeting 2017:
Varicella-Zoster Virus: Diagnosis

Whitney A. High, MD, JD, MEng
 whitney.high@ucdenver.edu
 Associate Professor, Dermatology & Pathology
 Director, Dermatopathology Laboratory
 University of Colorado School of Medicine



March 4, 2017
 Orlando, FL

Varicella Zoster
 Diagnosis



www.giantmicrobe.com

Varicella Zoster

- Primary infection = “chickenpox”
 - crops of vesicular lesions
 - centripetal predominance
- Reactivation from ganglia = “shingles”
 - painful, unilateral vesicular eruption
 - most often dermatomal
- Most often both conditions are often diagnosed on clinical grounds, and this is ok

“Chickenpox”
 truncal
 predominance
 (centripetal)



“Shingles”
 extending down
 arm



VZV
 Ancillary Diagnostic Techniques

Biopsy – cytopathic effect under H&E examination

Tzanck – sensitive in experienced hands

DFA – rapid, allows for diff. of VZV from HSV

Culture – fastidious technique, ≥ 5 days for results





Antibodies – IgM rises first, IgG is sign of immunity

Slit-lamp – performed by specialists in VZO

PCR – 4x more sensitive, increasing in popularity


Typical Biopsy Findings

H&E stain alone
(cytopathic effect indicative of *Herpes* family infection)



Varicella Zoster Virus (SG1-1, SG1-SG4, NCP-1 & IE-62)



Add to Favorites



Analyte Specific Reagent: Analytical and performance characteristics are not determined

Species: Mouse cocktail Reactivity: N/A Dilution: N/A
 Visualization: N/A Isotype: Mixed Control: N/A


Labeling designation:				Instructions for use:	
ASR	IVD	IVD	RUO	IFU	MSDS

What is an “analyte specific reagent” (ASR)?

- FDA defines ASR as “antibodies... intended for use in a diagnostic application... in biological specimens.”
- An ASR is the active ingredient of an in-house test, and ***must be labeled as such*** in a report
- It requires a validation and disclaimer

Tzanck Prep Bedside Cytology



Dr. Arnauld Tzanck, 1886-1954


- Locate a fresh lesion and clean the area gently
- Unroof blister - scrape roof/base with #15 blade
- Transfer material to glass slide
- “Fix” with couple sprays of cytofix (preferred)
 - or with 95% EtOH, or gentle heat, or air dry
- Stain with Wright/Giemsa for about 40-60 sec
- Rinse gently and carefully dry slide by blotting
- Add drop of immersion oil and cover slip

Methods of Fixation




When nothing else is around?

- My “Kit” – #15 blade, glass slide, alcohol pads (kind used for injections), and shoes






Tzanck stains



Tzanck Findings

- 80% sensitivity
- 90% specificity




DFA Results

- Lower cost (relative to culture)
- Rapid turn around (90 minutes)



Allows for discrimination of:
HSV1, HSV2 and VZV



Antibody Tests - IgM

- IgM rises in acute infection (and in zoster)
- IgM testing is **LESS** sensitive than PCR
- Commercial IgM test can give **false negative**
- A positive IgM ELISA result does not exclude re-infection or reactivation (zoster)
- A positive IgM result from a person with characteristic rash is usually interpreted as primary varicella (but it doesn't have to be)

CDC Admonishment

- "Patients with zoster may mount an IgM response and are expected to mount a memory IgG response."
- Positive IgM by ELISA can be indicative of:
 - primary VZV infection
 - or reactivation (zoster)
- It is **difficult to detect an increase in IgG** for laboratory diagnosis of zoster as patients may have a high baseline antibody titers

Antibody Tests - IgG


- A positive IgG ELISA indicates antibodies from past wild-type infection or vaccination
- No commercial assay can detect immunity in all vaccinated persons (lacks sensitivity)
- IgG – positive result indicates immunity
- Paired acute and convalescent sera w/ 4x rise in IgG has excellent specificity for varicella, but is not as sensitive as PCR



More from the CDC...

“In people with compromised immune systems, it may be difficult to distinguish between varicella and disseminated herpes zoster by serological testing or even by physical examination.

In these instances, a **history of VZV exposure** or of a **rash that began in a dermatomal pattern**, with results of VZV antibody testing at/before the time of onset may help guide the diagnosis.”



“... discontinued culture for these viruses...”

- PCR on swabs from lesions
- Same day results



More about PCR...

- PCR genotyping – distinguishes wild-type VZV from vaccine type (Oka)-strain
- May be performed on blood, cerebrospinal fluid, or biopsy specimens
- Fast, cheap, and has the **highest sensitivity** of **ALL** diagnostic techniques

Methods of Skin Sampling

- Polyester Swab Method
 - a sterile needle is used to un-roof the vesicle
 - vigorously swab the base of the lesion, without causing bleeding
 - collect epithelial cells from the base of the lesion (highest concentration of virus)
 - avoid cotton and wooden swabs as these absorb extraction buffer and inhibit PCR


Methods of Skin Sampling

- Glass Slide Method
 - rake the edge of the slide over the lesion, abrading the lesion
 - ensure that skin cells are gathered
 - swab the abraded lesion and collect the material also
 - dried maculopapular lesion material is stable for several weeks at ambient temperature

Methods of Skin Sampling

- Collecting Crusts (Scabs)
 - crusts are also excellent for PCR detection
 - crusts can be lifted off with glass slide
 - transferred directly into break-resistant, snap-cap or screw-top tubes
 - shipped per “shipping instructions” to CDC

CDC Video available at:
<http://www.cdc.gov/shingles/lab-testing/collecting-specimens.html>



PCR does not have to be done only on skin...

J Clin Virol. 2002 Feb;24(1-2):7-12.

Fulminant disseminated Varicella Zoster virus infection without skin involvement.
 Grant RA¹, Weisman SS, Sherman GS, Quinn VA, Patel M, Teiber B.

Author information

ABSTRACT

BACKGROUND: Varicella Zoster virus (VZV) infection is potentially very serious in bone marrow transplant recipients, and may manifest as a disseminated visceral infection. This condition is generally accompanied by a vesicular rash.

OBJECTIVES: We review here a case of fulminant fatal disseminated VZV infection, not accompanied by skin involvement, and the laboratory approaches currently available to diagnose this disease.

STUDY DESIGN: Post mortem tissue samples were subjected to histopathological examination, and tested for herpesviruses by electron microscopy and PCR.

RESULTS: Intracellular inclusions were noted by histological examination in the lungs, liver, kidneys and bone marrow. Particles with a herpesvirus morphology were visualized in liver tissue. VZV DNA was detected in liver and bone marrow by PCR followed by sequencing of the amplicons. Viremia was documented by retrospective testing of the serum by PCR.

CONCLUSIONS: A disseminated VZV infection which proved rapidly fatal was demonstrated in a case without skin manifestations. This rare presentation of VZV infection is potentially underdiagnosed. Testing for VZV viremia by PCR can at the very least suggest the diagnosis although whether plasma-associated viremia is truly pathogenomic of visceral disseminated infection remains to be established.

PMID: 11784223 (PubMed - indexed for MEDLINE)

Thank you.

