

Management of Radicular Pain

Mel Cusi

MBBS, FACSP, FFSEM (UK)

Sport & Exercise Medicine Physician



ORTHOSPORTS

ORTHOPAEDIC & SPORTS MEDICINE SERVICE

Dr Mel Cusi
Sport & Exercise Medicine Physician


Management of Radicular Pain

- A. Background
- B. Epidemiology
- C. Diagnosis
- D. Treatment



A. Background

- Names and concepts
 - Radicular pain
 - Radiculopathy
- Structures that can produce radicular Sx
 - Sinu-vertebral nerve
 - Nerve root
- Mechanisms of pain
 - Direct toxic effect of disc material
 - Chemical substances



B. Epidemiology

- Occurs in 3-5% of the population
 - More frequent in males in their 40's
 - More frequent in females in their 50's
- In sporting population
 - More frequent in sports that combine spinal flexion/extension with rotation
 - Fast bowlers, gymnasts, dancers, RU backrowers, golfers, weightlifters, baseball pitchers



C. Diagnosis

- Radicular pain is only a descriptive symptom
- Diagnosis is made on the usual basis of
 - History
 - Clinical examination
 - Appropriate investigations (when required)




History

- Acute LBP radiating to buttock / lower limb
- Worse with flexion, sneezing, coughing.
Sitting worse than standing
- Some pointers
 - Referred pain from L1-3 does not reach the knee
 - Unusual Symptoms (weight loss, fever, chills) point to something else
 - Beware of *cauda equina*: surgical emergency




Neurological Examination

- Sensation
 - Subjective
 - Objective (light touch, pinprick)
- Dermatomal distribution is a poor indicator of the level of pathology (Albert et al. 2010)



Neurological Examination

- Power: Identify what levels are deficient



Segmental motor innervation of the lower limb



ORTHOSPORTS

ORTHOPAEDIC & SPORTS MEDICINE SERVICE

Dr Mel Cusi
Sport & Exercise Medicine Physician


Neurological Examination

- Reflexes
 - Patellar tendon (knee jerk)
 - L2/3/4
 - Ankle tap (ankle jerk)
 - S1



Diferential diagnosis

- Disc protrusion / extrusion (nerve root compression)
- Hip pathology (L1-2 dermatome distribution)
- Trochanteric bursitis (buttock / thigh pain)
- SIJ incompetence (pseudosciatica)
- Facet joint pain (usually no leg pain)




Investigations

- To confirm or exclude a specific diagnosis
- X-ray: to exclude organic bone pathology
- CT scan: good to assess bone and disc hernias
- MRI: best modality to assess soft tissues
- Caution: treat the patient, not the picture

35% of normal 35 y.o. females show some form of “pathology” on MRI



Investigations



D. Treatment

- Initial
 - Non-operative
 - Surgical
- Aspects of Rehabilitation and secondary prevention (of recurrences)



Non operative Treatment of disc herniation

- Regular analgesia (not p.r.n)
- Anti-inflammatories
 - NSAIDS
 - Early peri-radicular steroid injection?
- Maintain physical activity (ADLs at least)
- Extension exercises (if extrusion or sequestration extension can make pain worse)
- For how long???



Non operative Treatment of disc herniation

- As long as there is improvement, continue non-operative treatment for up to three months.
- If progress is not adequate patient will tell you



Surgical Options (for disc herniation)

- Surgical decompression is a QOL decision
- Long term results similar to non-operative measures
- There is a 10% recurrence rate
- Discectomy +/- laminotomy +/- rhizolysis of the nerve root
- Microdiscectomy does not mean a smaller incision, but the use of vision augmentation



Aspects of Rehabilitation

- Make patient self-sufficient, independent of
 - Medication
 - Health practitioners
 - (doctors, physios, osteos, chiros, massage Rx, etc)
- Prevention of recurrences
 - General fitness
 - Lumbopelvic stability training
 - Manual handling and postural training





ORTHOSPORT
ORTHOPAEDIC & SPORTS MEDICINE

Mel Cusi
Physician