



Occupational  
Cancer  
Research  
Centre

# Epidemiologic Studies of Pentachlorophenol Users

**Paul A. Demers, PhD**

Director

Occupational Cancer Research Centre

Toronto, Canada

# Studies of PCP Users



- Pentachlorophenol and its sodium salt were used extensively in the lumber and wood preservation industry as fungicides
- Epidemiologic evidence on users come from:
  - Cohort studies of workers in the wood industries
  - Case-control conducted in regions where use was relatively high
  - The studies were conducted in the Nordic countries, the NW US, Western Canada, New Zealand, and Australia

# BC Sawmill Study: Background



- Penta- and tetrachlorophenol were used as fungicides in the softwood lumber industry 1950-1990
  - Estimated that 100,000 workers were exposed in British Columbia (BC), Canada
- Widespread use in lumber mills was discontinued because of concerns regarding dioxin contamination in the late 1980's

# BC Sawmill Study [Demers et al, 2006]

- 14 large softwood sawmills with good records
  - 11 had used pentachlorophenol or tetrachlorophenol
- Enumeration of workers using mill records
  - All workers employed for 1+ years, 1950-1995
  - Info on all jobs abstracted from mill records
- Mortality: 1950-1995
  - 27,464 workers, 4% lost to follow-up
- Cancer Incidence: 1969-1995
  - 25,685 workers, <1% lost to follow-up

# Use of Chlorophenols in the Lumber Industry





~95% of Exposure Dermal



Towards a cancer-free workplace

# Exposure Assessment



- Historical information collected to establish time periods when technology and work practices were constant
- Panels of 9-20 workers with 5+ years experience in each time period established for each mill
- Reliable estimates (ICC=0.91) [Teschke et al, 1989]
- Validation study of most recent time period using urinary levels ( $r=0.72-0.76$ ) [Hertzman et al, 1988]
- Exposure metrics created for each fungicide

# BC Sawmill Workers Cancer Incidence: 1969-1995



<u>Disease Category</u>	<u>Cancers</u>	<u>SIR (95% CI)</u>
All Cancers	2,571	1.02 (0.98-1.06)
Liver	21	0.79 (0.49-1.21)
Sino-nasal	6	1.05 (0.39-2.29)
Lung	519	1.05 (0.96-1.15)
Kidney	79	1.10 (0.88-1.38)
Connective tissue	13	0.84 (0.49-1.44)
Non-Hodgkin's lymphoma	92	0.99 (0.81-1.21)
Hodgkin's disease	18	0.94 (0.56-1.49)
Multiple myeloma	25	0.80 (0.52-1.18)




# Pentachlorophenol and Cancer in BC Sawmill Workers: Dose-Response Analyses



Cancer Site*	1-2 exposure yrs RR (95% CI)	2-5 exposure yrs RR (95% CI)	5+ exposure yrs RR (95% CI)
Lung (n=519)	1.11 (0.86-1.45)	1.07 (0.84-1.36)	1.12 (0.87-1.44)
Liver (n=21)	4.09 (0.89-18.8)	8.47 (2.21-32.5)	1.41 (0.21-9.22)
Kidney (n=79)	1.03 (0.49-2.18)	1.79 (0.99-3.24)	1.66 (0.85-3.23)
STS (n=23)	0.64 (0.18-2.20)	0.18 (0.04-0.85)	0 observed
NHL (n=92)	1.33 (0.70-2.52)	1.88 (1.08-3.28)	1.71 (0.91-3.24)
Myeloma (n=25)	2.09 (0.57-7.61)	1.30 (0.34-4.98)	4.18 (1.36-12.9)

\* STS: Soft Tissue Sarcoma, NHL: Non-Hodgkin's Lymphoma

# Pentachlorophenol and Cancer in BC Sawmill Workers: Latency Analyses



#Analyses performed using Poisson Regression, adjusted for age, calendar period & race

\*Trend:  $P < 0.10$

\*\*Trend:  $p < 0.05$

# Selected Results from Other Cohort Studies



- Robinson et al 1987, mortality among 2,283 men employed 1+ years at four plywood mills in the NW US 1945-1955
  - Sub-cohort, 1+ years of PCP *or formaldehyde* exposure: lymphosarcoma (SMR=2.50, 3 cancers) and Hodgkin's lymphoma (SMR=3.33, 2 cancers)
- Partanen et al 1993, nested case-control study of men in a Finnish wood industry cohort
  - NHL (n=8)/HL (n=4)/leukemia (n=12) and  $\geq 1$  month chlorophenol exposure, OR=0.91 (0.18-4.51)

# Case-Control Studies of PCP, CP, and Non-Hodgkin's Lymphoma



Study	Population	Exposure*	Results OR (95% CI)
Hardell et al 1994, Sweden	Men age 25-85 105 cases, 355 controls	1+ week continuous or 1 month total exposure to PCP	8.8 (3.4-24)
Hardell et al 1999, Sweden	Men age 25+ 404 cases, 741 controls	Any PCP exposure (>1 year before diagnosis)	1.2 (0.7-1.8)
Garabedian et al, 1999, USA	Men 32-60 995 cases, 1783 controls	High confidence CP - High intensity	1.1 (0.9-1.4) 1.1 (0.8-1.5)
Woods et al 1987, USA	Men age 20-79 576 cases, 694 controls	Any job with possible CP exposure - Wood preservers	0.9 (0.8-1.2) > 1.5
Pearce et al 1986, New Zealand	Men age <70 83 cases, 168 cancer controls, 228 population controls	Ever exposed to CP - Work as fencing contractor or farmer	1.3 (0.6-2.7) 2.0 (1.3-3.0)

\* PCP: Pentachlorophenol, CP: Any Chlorophenol

# Case-Control Studies of PCP, CP, and Soft Tissue Sarcoma



Study	Population	Exposure	Results
Hardell et al 1995, Sweden	Meta-analysis, 4 studies men age 25-80 434 cases, 948 controls	1+ week continuous or 1 month total PCP	2.8 (1.5-5.4)
Hoppin et al, 1998, USA	Men age 30-60 295 cases, 1908 controls	High confidence CP - High intensity	1.7 (1.1-2.7) 2.1 (1.2-3.7)
Woods et al 1987, USA	Men age 20-79 128 cases, 694 controls	Any possible CP Jobs with possible exposure	1.0 (0.7-1.5) ORs from 0.8-4.8
Smith et al 1984, New Zealand	Men age 20-80 82 cases, 92 controls	1+ day CP exposure >5 days, 10 yr before diagnosis Jobs with possible exposure	1.3 (0.5-3.6) 1.6 (0.5-5.2) ORs from 0.7-1.9

\* PCP: Pentachlorophenol, CP: Any Chlorophenol



# Results from Case-Control Studies for Other Cancer Sites

- Case-control studies, primarily in the Nordic countries and the USA, have looked at the risk of other cancers, such as multiple myeloma, colon, nasal, nasopharyngeal, and respiratory cancers, with chlorophenol exposure
- Power often limited
- Results often inconsistent with cohort results
- Too little evidence to evaluate based on PCP users

# Challenges for Epidemiologic Studies

- Relatively few studies of Pentachlorophenol users specifically assessed pentachlorophenol exposure
- Many studies lack power
- Need for more studies with strong exposure assessment – difficult for dermal exposure
- Confounding exposures?
- Co-exposures or contaminants?

# Cancers\* associated with Potential Occupational Confounders & Co-Exposures



Carcinogen	Sufficient evidence	Limited evidence
Dioxins	All cancers combined	Lung, NHL, STS
Wood dust	Sino-nasal, nasopharyngeal	
Formaldehyde	Nasopharyngeal, leukemia	Sino-nasal
Arsenic	Lung, skin, bladder	Kidney, liver, prostate
Chromium	Lung	Sino-nasal

\* From IARC Monographs 100c & 100f

# Conclusions for PCP Users



- An association with non-Hodgkin's lymphoma in the major cohort with some support from case-control
- Relatively consistent results for soft tissue sarcoma from Swedish case-control studies, mixed results elsewhere
- Some associations observed with other sites, such as multiple myeloma and kidney cancer, but too little data to evaluate based on PCP users

# References



- Demers PA, Davies HW, Friesen MC, Hertzman C, Ostry A, Hershler R, Teschke K (2006). Cancer and occupational exposure to pentachlorophenol and tetrachlorophenol. *Cancer Causes Control* 17:749-58.
- Garabedian MJ, Hoppin JA, Tolbert PE, Herrick RF, Brann EA (1999). Occupational exposure and non-Hodgkin's lymphoma. *J Occup Environ Med* 41:267-272.
- Hardell L, Eriksson M, Degerman A (1994). Exposure to phenoxyacetic acids, chlorophenols, or organic solvents in relation to histopathology, stage, and anatomical localization of non-Hodgkin's lymphoma. *Cancer Research* 54:2386-2389.
- Hardell L, Eriksson M, Degerman A (1995). Meta-analysis of four Swedish case-control studies on exposure to pesticides as risk-factor for soft-tissue sarcoma including the relation to tumour localization and histopathological type. *Int J Oncology* 6:847-851.
- Hardell L, Eriksson M (1999). A case-control study of non-Hodgkin Lymphoma and exposure to pesticides. *Cancer* 85(6):1353-1360.
- Hertzman C, Teschke K, Dimich-Ward H, Ostry A (1988). Validity and reliability of a method for retrospective evaluation of chlorophenolate exposure in the lumber industry. *Am J Ind Med* 14:703-13.
- Hoppin JA, Tolbert PE, Herrick RF, Freedman DS, Ragsdale BD, Horvat KR, Brann EA (1998). Occupational chlorophenol exposure and soft tissue sarcoma risk among men aged 30-60 years. *Am J Epidemiol* 148:693-703.
- International Agency for Research on Cancer (IARC) Working Group. A Review of Human Carcinogens: Arsenic, metals, fibres, and dusts. Lyon: IARC, 2012. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans, vol 100C.



## References (continued)



International Agency for Research on Cancer (IARC) Working Group. A Review of Human Carcinogens: Chemical agents and related occupations. Lyon: IARC, 2012. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans, vol 100F.

Partanen T, Kauppinen T, Luukkonen R, Hakulinen T, Pukkala E (1993). Malignant lymphomas and leukemias, and exposures in the wood industry: an industry-based case-referent study. *Int Arch Occup Environ Health* 64:593-596.

Pearce NE, Smith AH, Howard JK, Sheppard RA, Giles HJ, Teague CA (1986). Non-Hodgkin's lymphoma and exposure to phenoxyherbicides, chlorophenols, fencing work, and meat works employment: a case-control study. *Br J Ind Med* 43:75-83.

Robinson CF, Fowler D, Brown DP, Lemen RA. Plywood mill workers' mortality patterns 1945-1977. Cincinnati, OH: National Institute for Occupational Safety and Health, 1987. NTIS Publication PB86-221694.

Smith AH, Pearce NE, Fisher DO, Giles HJ, Teague CA, Howard JK (1984). Soft tissue sarcoma and exposure to phenoxyherbicides and chlorophenols in New Zealand. *J Natl Cancer Inst* 73:1111-1117.

Teschke K, Hertzman C, Dimich-Ward H, Ostry A, Blair J, Hershler R (1989). A comparison of exposure estimates by worker raters and industrial hygienists. *Scand J Work Environ Health* 15:424-9.

Woods JS, Polissar L, Severson RK, Heuser LS, Kulander BG (1987). Soft tissue sarcoma and non-Hodgkin's lymphoma in relation to phenoxyherbicide and chlorinated phenol exposure in western Washington. *J Natl Cancer Inst* 78:899-910.



Occupational  
Cancer  
Research  
Centre

# **Towards** a cancer free workplace

<http://occupationalcancer.ca>