

Two-boundary centralizer algebras

Zajj Daugherty

University of Wisconsin, Madison

November 8, 2009

Background

Let \mathfrak{g} be a finite dimensional complex reductive Lie algebra.

e.g. $\mathfrak{gl}_n(\mathbb{C})$, $\mathfrak{sl}_n(\mathbb{C})$, $\mathfrak{so}_n(\mathbb{C})$, $\mathfrak{sp}_{2n}(\mathbb{C})$.

Let M , N , and V be finite dimensional simple \mathfrak{g} -modules.

Goal:

Understand $\text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k})$.

(the set of endomorphisms which commute with the action of \mathfrak{g})

Examples of $\text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k})$

Fix $k < n$ integers.

Let $L(\lambda)$ be the f.d. irreducible \mathfrak{g} -module of highest weight λ .

Let $V = L(\omega_1)$.

① If $\mathfrak{g} = \mathfrak{sl}_n$ and

- $M = N = L(0)$, this gives $\mathbb{C}S_k$;
- $M = L(0)$ and $N = L(\lambda)$, this gives is a quotient of the graded Hecke algebra of type A;

② If $\mathfrak{g} = \mathfrak{so}_n$ or \mathfrak{sp}_{2n} and

- $M = N = L(0)$, this gives the Brauer algebra;
- $M = L(0)$ and $N = L(\lambda)$, this gives a quotient of the degenerate affine Wenzl algebra.

Quantized versions yield standard and affine type A Hecke and Birman-Murakami-Wenzl algebras.

Big question:

Is there an algebra which has centralizers $\text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k})$ as quotients?

Definition

The *two-boundary graded braid group* \mathcal{G}_k is the \mathbb{C} -algebra generated by

$$\mathbb{C}S_k = \mathbb{C} \left\langle s_i \mid \begin{array}{l} i = 1, \dots, k \\ s_i^2 = 1 \\ s_i s_j = s_j s_i \quad |i - j| > 1 \\ s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1} \end{array} \right\rangle$$

$$\mathbb{C}[z_0, z_1, \dots, z_k], \mathbb{C}[y_1, \dots, y_k], \mathbb{C}[x_1, \dots, x_k]$$

and relations...

Representations of \mathcal{G}_k

We'll define an action of \mathcal{G}_k on $M \otimes N \otimes V^{\otimes k}$:

$\mathbb{C}S_k$ permutes factors of $V^{\otimes k}$,

$\mathbb{C}[x_1, \dots, x_k]$ acts on M and $V^{\otimes k}$,

$\mathbb{C}[y_1, \dots, y_k]$ acts on N and $V^{\otimes k}$,

$\mathbb{C}[z_1, \dots, z_k]$ acts on $M \otimes N$ together and $V^{\otimes k}$,

z_0 acts on $M \otimes N$ alone,

by nested central elements of $\mathcal{U}\mathfrak{g}$.

Let $\langle, \rangle : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathbb{C}$ be the trace form:

$$\langle x, y \rangle = \text{Tr}(xy), \quad \text{where } x \text{ and } y \text{ are viewed in a defining rep of } \mathfrak{g}.$$

Let $\{b\}$ be a basis of \mathfrak{g} and $\{b^*\}$ the dual basis wrt \langle, \rangle .

$$\text{Let } \kappa = \sum_b bb^*.$$

κ is the *Casimir invariant* and is central in $\mathcal{U}\mathfrak{g}$.

Theorem (D.)

Define $\Phi: \mathcal{G}_k \rightarrow \text{End}(M \otimes N \otimes V^{\otimes k})$

$$\Phi(s_j) = \text{id}_M \otimes \text{id}_N \otimes \text{id}_V^{\otimes(j-1)} \otimes s_1 \otimes \text{id}_V^{\otimes(k-j-1)},$$

$$\Phi(x_j) = \frac{1}{2}(\kappa|_{M \otimes V^{\otimes j}} - \kappa|_{M \otimes V^{\otimes j-1}}),$$

$$\Phi(y_j) = \frac{1}{2}(\kappa|_{N \otimes V^{\otimes j}} - \kappa|_{N \otimes V^{\otimes j-1}}),$$

$$\Phi(z_j) = \frac{1}{2}(\kappa|_{M \otimes N \otimes V^{\otimes j}} - \kappa|_{M \otimes N \otimes V^{\otimes j-1}} + \kappa|_V),$$

$$\Phi(z_0) = \frac{1}{2}(\kappa|_{M \otimes N} - \kappa|_M - \kappa|_N),$$

where $s_1 \cdot (v_{i_1} \otimes v_{i_2}) = v_{i_2} \otimes v_{i_1}$.

Then Φ is a representation of \mathcal{G}_k which commutes with the action of \mathfrak{g} .

An Example:

Is there an algebra which has centralizers
 $\text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k})$ as quotients
when \mathfrak{g} is of type A?

Definition

Fix $a, b, p, q \in \mathbb{Z}_{>0}$.

The *extended two-boundary graded Hecke algebra* $\mathcal{H}_k^{\text{ext}}$ is the quotient of the two-boundary graded braid group by the relations

$$\begin{aligned}t_{s_i} x_i &= x_{i+1} t_{s_i} - 1, \\t_{s_i} y_i &= y_{i+1} t_{s_i} - 1, \quad i = 1, \dots, k-1. \\t_{s_i} z_i &= z_{i+1} t_{s_i} - 1,\end{aligned}$$

$$(x_1 - a)(x_1 + p) = 0 \quad (y_1 - b)(y_1 + q) = 0.$$

A *partition* is a collections of boxes:

$$\lambda = \begin{array}{|c|c|c|c|} \hline 0 & 1 & 2 & 3 \\ \hline -1 & 0 & 1 & \\ \hline -2 & & & \\ \hline \end{array}$$

If a box B is in row i and column j , then the *content* of B is

$$c(B) = j - i.$$

If $\lambda = (a^p)$ is rectangular, there are exactly two “addable” boxes:

$$(a^p) = \begin{array}{|c|c|c|c|} \hline & & & a \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline -p & & & \\ \hline \end{array}$$

(recall relations $(x_1 - a)(x_1 + p) = 0$ and $(y_1 - b)(y_1 + q) = 0$)

Theorem (D.)

Fix $k < n$ non-neg. integers.

Let $\mathfrak{g} = \mathfrak{gl}_n$, $M = L((a^p))$, $N = L((b^q))$, and $V = L((1^1))$.

(1) Φ is a rep. of $\mathcal{H}_k^{\text{ext}}$ which commutes with the \mathfrak{g} -action, so

$$\Phi(\mathcal{H}_k^{\text{ext}}) \subseteq \text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k}).$$

(2) For suitable choices of a, b, p, q ,

$$\Phi(\mathcal{H}_k^{\text{ext}}) = \text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k}).$$

Remark


- (1) When Φ is not surjective, the image differs by a portion of the action of the center of $\mathcal{U}\mathfrak{g}$ on $M \otimes N$.
- (2) Same theorem for $\mathfrak{g} = \mathfrak{sl}_n$ and a shift of Φ .


Let $M = L((a^p))$ and $N = L((b^q))$. Then

$$M \otimes N = \bigoplus_{\lambda \in \Lambda} L(\lambda) \quad (\text{multiplicity one!})$$


where Λ is the set of partitions:...

(Okata)





A two-dimensional $\mathcal{H}_1^{\text{ext}}$ -module:



$$z_0 = \begin{pmatrix} 4a & 0 \\ 0 & 3a - p \end{pmatrix}$$


$$z_1 = \begin{pmatrix} -p & 0 \\ 0 & a \end{pmatrix}$$

$$x_1 \sim \begin{pmatrix} -p & 0 \\ 0 & a \end{pmatrix}$$

$$y_1 \sim \begin{pmatrix} -2 & 0 \\ 0 & 2 \end{pmatrix}$$

(formulas x_1, y_1, z_1, z_0 all given in terms of contents of added boxes)

An eight-dimensional $\mathcal{H}_2^{\text{ext}}$ -module:



(Labeling edges by action of $z_1 - \frac{1}{2}(a - p + b - q)$)

More examples of $\text{End}_{\mathfrak{g}}(M \otimes N \otimes V^{\otimes k})$

Fix $k < n$ integers.

Let $L(\lambda)$ be the f.d. irreducible \mathfrak{g} -module of highest weight λ .

Let $V = L(\omega_1)$.

- 1 When $\mathfrak{g} = \mathfrak{sl}_n$ or \mathfrak{gl}_n , and M and N are rectangular, we get the (extended) two-boundary graded Hecke algebra. (explored in thesis)
- 2 When $\mathfrak{g} = \mathfrak{so}_n$ or \mathfrak{sp}_{2n} , and M and N are rectangular, we get the *two-boundary graded Brauer algebra*. (future work)

Quantized versions should yield two-boundary affine Hecke and BMW algebras.

Striking: The two-boundary affine Hecke algebra is isomorphic to the type C affine Hecke algebra. Similarities also appear suggestively in graded versions.

References

- [OR] R. Orellana and A. Ram, *Affine braids, Markov traces and the category \mathcal{O}* , Proceedings of the International Colloquium on Algebraic Groups and Homogeneous Spaces Mumbai 2004, V.B. Mehta ed., Tata Institute of Fundamental Research, Narosa Publishing House, Amer. Math. Soc. (2007) 423-473.
- [GN] J. de Gier and A. Nichols, *The two-boundary Temperley-Lieb algebra*, J. Algebra **321** (2009) 11321167.

In preparation:

- [Dau] Z. Daugherty, *Two-boundary graded centralizer algebras*
- [DRV] Z. Daugherty, A. Ram, R. Virk, *Affine and graded BMW algebras*

find me at...

<http://www.math.wisc.edu/~daughert/>