

Innate Immunity (part 1)

BIOS 486A/586A

Kenneth J. Goodrum, Ph.D.
Dept. Biomedical Sciences
Ohio University
2005

Topic Outline

- Define innate vs. adaptive immunity
- Identify innate immune cells & molecules
- Describe functions of phagocytic cells
 - Detection of microbes (pattern recognition receptors)
 - Phagocytosis & microbicidal mechanisms
 - Secretion of pro-inflammatory proteins (Cytokines)
 - directed migration (Chemotaxis)
- Describe inflammation/leukocyte adhesion
- Describe other innate cells/factors

Immune Mechanisms

• INNATE

- Present at birth
- Self-discrimination
- Limited diversity
- Nonspecific defense
- No memory
- Examples: skin, cough, gastric pH, fever, inflammation (phagocytic cells)

• ACQUIRED

- Appears after contact with antigen
- Self-discrimination
- Vast diversity
- Specific defense
- Memory responses
- Examples: antibody, cytotoxic lymphocytes

Innate Immunity

- **Present at birth**
- **Nonspecific**
 - Response does not target one specific immunogen
- **Limited diversity**
 - Fixed, repeating, broad responses to a limited number of foreign substances
- **No memory**
 - primary and secondary responses are identical


Components of Innate Immunity

- **Physical/mechanical barriers**
 - Intact skin, epithelial layers, cough, fever
- **Nonspecific chemical factors**
 - Antimicrobial peptides & fatty acids, gastric pH, lysozyme
- **Inflammation**
 - Phagocytes (engulf and digest microbes)
 - **Proinflammatory factors** (cytokines, complement proteins)
- **Natural killer cells** (nonspecific cytotoxic cells)
- **Interferon** (produced by virus-infected cells and induces antiviral state in neighboring uninfected cells)

Intrinsic epithelial barriers to infection

Mechanical	Epithelial cells joined by tight junctions Longitudinal flow of air or fluid across epithelium Movement of mucus by cilia
Chemical	Fatty acids (skin) Enzymes: lysozyme (saliva, sweat, tears), pepsin (gut) Low pH (stomach) Antibacterial peptides; defensins (skin, gut), cryptidins (intestine)
Microbiological	Normal flora compete for nutrients and attachment to epithelium and can produce antibacterial substances


Fig 2.4 © 2001 Garland Science



Neutrophil (neutrophilic polymorphonuclear leukocyte) in stained blood smear. Neutrophils (PMN) are professional phagocytes

2. 3

Copyright © 2004, 2001 Elsevier Inc. All rights reserved.



(From Fawcett DW. Bloom and Fawcett Textbook of Histology, 12th ed. WB Saunders, Philadelphia, 1994.)
Copyright © 2004, 2001 Elsevier Inc. All rights reserved.

2. 4



Mononuclear phagocytes (monocytes & macrophages)

Phagocyte Functions

- Recognition of microbes (pattern recognition)
- Synthesis and secretion of cytokines/chemokines
- Phagocytosis (binding and engulfment of particles)
- Intralysosomal digestion and killing of ingested material.
 - Antigen presentation to lymphocytes
- Chemotaxis (migration toward a chemoattractive signal)

Innate recognition of Microbes by Phagocytes

- Phagocytes recognize **pathogen-associated molecular patterns** (PAMP) via pattern recognition receptors such as **CD14/Toll receptors** and produce proinflammatory cytokines and chemokines.
- Microbial substances (LPS,LTA,PPG) may directly activate plasma **complement** proteins stimulating inflammation




Innate immune recognition by Toll-like receptors	
Toll-like receptor	Ligand
TLR-1 dimer	Peptidoglycan Lipoproteins Lipoarabinomannan (mycobacteria) GPI (<i>T. cruzi</i>) Zymosan (yeast)
TLR-2/TLR-6 dimer	
TLR-3	dsRNA
TLR-4 dimer (plus CD14)	LPS (Gram-negative bacteria)
TLR-5	Flagellin
TLR-9	Unmethylated CpG DNA

Figure 2-12 Immunobiology, 6/e. © Garland Science 2005

Toll-like receptors (TLR) are pattern recognition receptors that bind molecular patterns common to multiple infectious microorganisms. Phagocytes use these receptors to distinguish self from inflammatory non-self material.

Phagocytosis (cellular ingestion of particles)

- **Non-opsonic**
 - Direct engulfment via innate pattern recognition receptors. Slow, limited, inefficient
- **Opsonic**
 - Engulfment of complement-coated or antibody-coated microbes via complement receptors (CR) or antibody receptors (FcR). Rapid, very efficient



Phagocyte Killing Mechanisms

- **OXYGEN INDEPENDENT**
 - lysosomal hydrolases, lysozyme, lactoferrin, defensins, acid pH
- **OXYGEN-DEPENDENT (Reactive oxygen and reactive nitrogen intermediates)**
 - Hydrogen peroxide, superoxide anion, hydroxyl radical, hypochlorite
 - Reactive Nitrogen Intermediates (RNI)
 - Nitric oxide, peroxynitrite



Inflammation



- Definition
 - local accumulation of fluid, plasma proteins, and WBC that is initiated by physical injury, infection, or a local immune response.

Proinflammatory cytokines

- Cellular sources: **macrophages, infected cells, injured cells**
- Factors: [interleukin = IL]
 - **CYTOKINES**: IL-1, IL-6, TNF- α , IL-12
 - Induce fever, acute phase protein synthesis, leukocyte adhesion
 - **CHEMOKINES**: IL-8
 - Attracts and recruits neutrophils

Proinflammatory cytokines secreted by macrophages stimulated by bacterial products via TLR





CHEMOTAXIS (directed migration)

- **Dependence on:**
 - **ADHESION MOLECULES (induced by cytokines)**
 - β 2-integrins on leukocyte
 - E-selectins on vascular endothelium
 - **RECEPTORS for Chemoattractants**
 - **CHEMOATTRACTANTS**
 - Leukotriene B4, Complement C5a, bacterial formyl-methionyl peptides, chemokines (interleukin-8, etc.)

Factors which activate and recruit inflammatory phagocytes

- **Cytokines (released locally) stimulate phagocyte adhesion to vascular endothelium and extravasation (diapedesis)**
- **Chemoattractant compounds direct phagocytes to the site of infection.**
- **“Acute phase” and complement proteins enhance phagocytosis and killing of microbes by recruited “inflammatory” phagocytes**

Leukocyte adhesion and diapedesis. Cytokine-induced adhesion molecules and locally-released chemokines promote binding of circulating phagocytes to the vascular endothelium and cause a directed migration to sites of tissue injury or infection. (slide 1)




Figure 2-44 part 2 of 3 Immunobiology, 6/e. (© Garland Science 2005)

Leukocyte adhesion and diapedesis (slide 2)




Figure 2-44 part 3 of 3 Immunobiology, 6/e. (© Garland Science 2005)




Figure 2-48 Immunobiology, 6/e. (© Garland Science 2005)

Innate immune functions of interferons




Figure 2-50 Immunobiology, 6/e. (© Garland Science 2005)

Natural killer cells exhibit cytotoxicity toward cells that downregulate MHC molecules that are present on normal cells.

Innate-like lymphocytes		
B-1 cells	Epithelial $\gamma\delta$ cells	NK T cells
Make natural antibody, protect against infection with <i>Streptococcus</i>	Produce cytokines rapidly	Produce cytokines rapidly
Ligands not MHC associated	Ligands are MHC class IB associated	Ligands are lipids bound to CD1d
Cannot be boosted	Cannot be boosted	Cannot be boosted

Figure 2-52 Immunobiology, 6/e. (© Garland Science 2005)