



*The Role of the  
Respiratory Therapist  
at End-of-Life Care*



# Kubler-Ross Model

(Five Stages of Grief)

Denial

Anger

Bargaining

Depression

Acceptance



# Denial

**“But I feel fine, this can’t be happening to me”**

- Denial is usually a temporary defense mechanism for the individual**
- Denial is the FIRST defense mechanism and some people can be locked in this stage**



# Anger

**‘Why me?..It’s not fair..How could this happen to me?’**

**-In this stage you recognize that denial cannot continue.**

**-Patients are difficult to care for**

**-Feelings of rage and envy**

**-Important to remain non-judgmental and don’t too many things personally**



# Bargaining

**“I’ll do anything for a few more years” .....**

- This stage involves hope that can delay or postpone**
- Negotiations are made with a higher power**
  - Promises are made**
- Bargaining never works with life and death**



# Depression

**“I’m so sad, why bother with anything, I’m going to die soon , what’s the point?”**

- Begins to understand certainly of death**
- Can become silent , withdrawn, refuse visitors**
- Don’t try to cheer up someone in this stage**
- These emotions show that a person has begun to accept the situation**



# Acceptance

**“It’s going to be ok, I can’t fight it anymore”**

**-Coming to terms with your mortality**

**-The individuals who are dying come to terms with this stage long before some others, who unfortunately must pass through their own individual stages when dealing with death**



# Quotes

**“Dying is nothing to fear, it can be the most wonderful experience of your life. It all depends on how you have lived”**

**“Those who learned to know death, rather than fight it, become our teachers about life”**

**“Watching a peaceful death of a human being reminds us of a falling star; one of a million lights in a vast sky that flares up for a brief moment only to disappear into the endless night forever”**





# Caregivers and Coping

- All have to face death in our lives
- Death usually comes at the end of a long struggle on the patients part, also for the caregiver.
- You may be worn to a frazzle
- Keeping your emotions on hold
- Not easy for anyone, especially caregivers



- Two and a half million people die annually in the United States and almost 80% occur in healthcare institutions**
  - RT's , other direct caregivers often times look subconsciously at death as failure, therefore shy away from those dying patients**
- Researchers have indicated caregivers often lack the basic knowledge for EOL care**
- There is evidence that communication with the dying and their families is less than optimal and that few healthcare providers receive adequate training in appropriate communication skills**



# Ethical vs Moral vs History

- RT's are trained to work in all aspects of healthcare
  - We manage diseases of the pulmonary system
- Highest skill set involves care of patients on mechanical ventilation and the operation of the ventilator itself
- Requires hands-on patient care and deal with life and death on a daily basis
  - One unique difference in this position
- RT's are frequently called upon to remove assistive ventilation which ultimately results in their death



# Terminal Weaning

- Removal of ventilator support is known as terminal weaning
- Patients that are terminally weaned , unable to maintain their own ventilation; no expectation of ever being able to live off a ventilator
- Actually requires the respiratory therapist to remove the ventilator, knowing that at completion of this task the patient expire
- This can sometimes lead to ethical dilemmas for some RT's, even though or she is doing their job and breaking no laws, they are playing a role in ending the life of another individual



# Weaning or Extubation

- Two common methods of assisted ventilation removal
  - Terminal Weaning; ETT left in place as assisted ventilation is withdrawn
  - Terminal Extubation; ETT is removed as ventilatory support is removed simultaneously
- Important to remember is not a form of euthanasia; there is no assistance with death in this procedure, as it merely allows death to occur naturally
- Patients can survive when taken off life support



# Patient Preference/Right

- It is very hard for some RT's and caregivers to accept that a great number of patients choose death over staying alive on the ventilator**
- Respiratory Therapists are trained to safeguard and protect life, so terminal weaning always creates ethical dilemmas**
- Patients often come to the realization, when their quality of life is not worthy of continuing and elect to refuse to continue life sustaining measures**





# Removal and Brain Death

- Patients that are brain dead are often much easier to terminal wean for respiratory therapists from both a moral and ethical standpoint than other patients
  - In this situation the RT is really not removing life support because the patient is already considered dead ; most religious organizations recognize brain death as true death
  - Often are best candidates for organ donation



# Conscious Terminal Wean

- Withdrawing ventilation from a conscious apneic patient is especially difficult for a respiratory therapist**
- There is often a bond of trust that develops with long term ventilator patients**
- The weaning process usually begins with the patient alert throughout the process until death occurs**
- This can be a very taxing experience for the attending respiratory therapist as you go from interacting to removing life support**





# Family Conflicts

- Conflicts about EOL care is common place in health care
  - Family members often have a difficult time remaining objective
- Important to remember that the patient, family members, and even the clinical team may all be going through the different stages of DABDA
  - Luckily clinicians make it to the acceptance phase simply because of their constant occupational exposure to death
- If the patient is capable, they are the best resource to defend the terminal weaning



# Support Systems

- In order to handle a terminal wean, respiratory therapists need the full support of the hospital to properly handle the whole process, from a professional, emotional and psychological stand point
- After a terminal wean there should be a debriefing with the entire clinical team
  - This provides and opportunity or outlet to discuss feelings with other members who can relate
    - Provides opportunity for improvement
- Not recommended for same RT to perform back to back terminal weans



# To Conclude

- Despite progress, practicing RT's and other direct caregivers face many dilemmas and emotional barriers in providing EOL care**
- Not in curricular content**
- RT's have an easier time terminally weaning brain dead patients**
- Terminal weaning IS one of the greatest ethical dilemmas that RT's face in their profession**
- Although terminal weaning is a difficult part of the job duties, it can be less stressful when a strong support system is in place**