

Project: Ghana Emergency Medicine Collaborative

Document Title: COPD in the Emergency Department

Author(s): Frank Madore (Hennepin County Medical Center), MD 2012

License: Unless otherwise noted, this material is made available under the terms of the **Creative Commons Attribution Share Alike-3.0 License:**
<http://creativecommons.org/licenses/by-sa/3.0/>

We have reviewed this material in accordance with U.S. Copyright Law **and have tried to maximize your ability to use, share, and adapt it.** These lectures have been modified in the process of making a publicly shareable version. The citation key on the following slide provides information about how you may share and adapt this material.

Copyright holders of content included in this material should contact open.michigan@umich.edu with any questions, corrections, or clarification regarding the use of content.

For more information about **how to cite** these materials visit <http://open.umich.edu/privacy-and-terms-use>.

Any **medical information** in this material is intended to inform and educate and is **not a tool for self-diagnosis** or a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional. Please speak to your physician if you have questions about your medical condition.

Viewer discretion is advised: Some medical content is graphic and may not be suitable for all viewers.

Use + Share + Adapt

{ Content the copyright holder, author, or law permits you to use, share and adapt. }



Public Domain – Government: Works that are produced by the U.S. Government. (17 USC § 105)



Public Domain – Expired: Works that are no longer protected due to an expired copyright term.



Public Domain – Self Dedicated: Works that a copyright holder has dedicated to the public domain.



Creative Commons – Zero Waiver



Creative Commons – Attribution License



Creative Commons – Attribution Share Alike License



Creative Commons – Attribution Noncommercial License



Creative Commons – Attribution Noncommercial Share Alike License



GNU – Free Documentation License

Make Your Own Assessment

{ Content Open.Michigan believes can be used, shared, and adapted because it is ineligible for copyright. }



Public Domain – Ineligible: Works that are ineligible for copyright protection in the U.S. (17 USC § 102(b)) *laws in your jurisdiction may differ

{ Content Open.Michigan has used under a Fair Use determination. }



Fair Use: Use of works that is determined to be Fair consistent with the U.S. Copyright Act. (17 USC § 107) *laws in your jurisdiction may differ
Our determination **DOES NOT** mean that all uses of this 3rd-party content are Fair Uses and we **DO NOT** guarantee that your use of the content is Fair.

To use this content you should **do your own independent analysis** to determine whether or not your use will be Fair.

COPD

in the emergency department

Frank Madore, MD
Hennepin County Medical Center
Minneapolis, MN, USA

BACKGROUND

epidemiology

- one of the top causes of death worldwide
- 7th most common cause of disability by 2030
- billions of dollars per year in treatment and lost productivity
- under-reported (only 50% see a physician for an exacerbation)
- 2% of all hospitalizations, 20% >65 years

pathophysiology

- chronic airway inflammation as in asthma
 - asthma: eosinophils
 - COPD: neutrophils, CD8+ lymphocytes, & macrophages
- lung parenchyma damaged by TNF, leukotriene B4, & interleukin 8
 - hence, poorer response to anti-inflammatory treatments than asthma
- COPD = chronic bronchitis + emphysema

chronic bronchitis

- progressive scarring and narrowing of airways
→ obstruction
- increase in goblet cells → mucus plugs
- epithelial damage → mucociliary impairment →
decreased clearing of bacteria and mucus

emphysema

- destruction of alveoli
- loss of elasticity
- collapse of small airways
- chronic air trapping/hyperinflation
- prolonged expiratory phase
 - decreased FEV1 (forced expiratory volume in 1 second)

natural course

- chronic bronchitis + emphysema → decreased size of pulmonary vascular bed → capacity for gas exchange
- chronic hypercapnia and hypoxemia result
- lung vessels thicken, hemoglobin increases → increased pulmonary vascular resistance → pulmonary hypertension → R-sided heart failure

etiology

- smoking, pollution (incl indoor cooking), occupational exposures, genetic factors
- negatively affect body's natural oxidant/antioxidant and protease/antiprotease balances
- secondary effects:
 - weight loss, muscular wasting, metabolic derangement, depression etc.

staging

- stage I – mild
 - FEV1 > 80% predicted, little to no symptoms
- stage II – moderate
 - FEV1 50-80%, shortness of breath on exertion, occasional exacerbations
- stage III – severe
 - FEV1 30-50%, shortness of breath at rest, frequent exacerbations
- stage IV – very severe
 - FEV1 < 30%, ↓pO₂, ↑pCO₂, cor pulmonale

CLINICAL

physical exam

- chronic bronchitis - “blue bloater”
 - chronic respiratory failure, cor pulmonale, polycythemia, hypoxia
 - cyanosis, facial plethora, edema, JVD, frequent cough
- emphysema - “pink puffer”
 - thin, anxious, dyspneic, barrel chested, uses accessory muscles, pursed lip exhalation (auto-PEEP)
- most patients exhibit elements of both

exacerbation

- acute and unusual change in baseline dyspnea, cough, or sputum production
- warrants change in baseline medication
- more common in winter
 - suggests relation to seasonal viruses (RSV, coronavirus, influenza, rhinovirus)
- bacteria isolated in 50% of exacerbations
 - also isolated in a similar proportion of COPD patients without exacerbation

comorbidities

- patients with COPD are also at much higher risk for:
 - CAD/acute coronary syndrome (smoking)
 - pulmonary embolism (sedentary)
 - pneumonia (decreased ciliary function)
 - congestive heart failure (R-sided)
 - pneumothorax (ruptured bullae)
 - arrhythmias (atrial tachy)
- be careful to avoid premature closure
 - “just a COPD exacerbation”

diagnostic approach

- pulse oximetry – compare to baseline
- ABG – limited utility, mgt guided by exam
 - acute resp acidosis, compensatory metabolic alkalosis – values don't predict outcome
- ECG – ACS, arrhythmia
- CBC – limited value (↑WBC, ↑Hb)
- BNP – may help rule out CHF
- d-dimer – may help rule out PE

imaging

- CXR – atelectasis, PNA, CHF, pneumothorax, bullae, pericardial/pleural effusions, pulmonary fibrosis, etc.
- US – CHF, PE, pericardial effusion, pneumothorax, pleural effusion
 - careful with interpretation of large RV
- CT – definitive diagnosis of PE, bullae vs. pneumothorax, atx vs. PNA, etc. in unclear clinical situations

MANAGEMENT

long term

- chronic inhaled steroids
- albuterol as needed
- influenza and pneumonia vaccines
- home oxygen when necessary
- smoking cessation

case

- 63 yo M with a history of COPD p/w shortness of breath, cough
- started out as a cold with worsening over 3 days
- increasingly productive cough
- more frequent albuterol use
- VS: P104, BP166/94, R26, T37.4, 89% RA
 - appears anxious, dyspneic
 - poor air movement on auscultation

acute exacerbation

- **A – airway**
 - intubation if obtunded, severe respiratory distress, or clinically tiring out (RSI vs. nasal)
- **B – breathing**
 - inhaled albuterol/ipatropium
 - non-invasive positive pressure ventilation
 - oxygen
- **C – circulate medications**
 - antibiotics
 - corticosteroids

airway

- indications for intubation
 - severe respiratory distress
 - agitation, obtundation
 - fails NIPPV
 - shock
 - respiratory arrest
- methods of intubation
 - RSI – takes away respiratory drive, familiar
 - nasal – preserves ventilation, less familiar

breathing

- oxygen (as little as possible for sat ~90%)
 - monitor respiratory rate, intervene if slow
- albuterol 2.5 mg neb, titrate to effect
 - beta 2 agonist → bronchodilation
- ipatropium 0.5 mg neb, repeat x3
 - anticholinergic → inhibits smooth muscle contraction, decreases secretions
- non-invasive positive pressure ventilation
 - provides extrinsic PEEP to ↓work of breathing

circulate medications

- corticosteroids
 - 60 mg prednisone PO x1, 40 mg daily x4
 - 125 mg solumedrol IV x1
- antibiotics
 - objective evidence of pneumonia
 - increased sputum production
 - critically ill patients (ventilated)

disposition

- significant worsening from baseline
- poor response to ED treatment
- significant co-morbid diseases
- hypoxia
- unable to care for self
 - can they get around their house?
 - can they keep down fluids?
 - who will call for help if they get worse?

SUMMARY

in conclusion...

- it's not always “just” a COPD exacerbation
- treat with beta agonists, anticholinergics, and steroids
- try NIPPV before intubating
- re-evaluate frequently
- make them prove to you that they can go home

QUESTIONS