



The cell as the basic unit of life



Prokaryotic	Eukaryotic
Bacteria, most single-celled organisms	All multicellular organisms
Simple, smaller	Larger, more complex
Lack membrane-bound organelles such as nucleus	Contain membrane-bound organelles such as a nucleus
Circular DNA	Linear DNA (chromosomes)


Figure 1.8





Overview: Cell Structure and Function

- Video: <http://www.youtube.com/watch?v=o1GQyciJaTA>
- Questions:
 - What is the function of the *plasma membrane*?
 - What is the function of the *nucleus*?
 - What is the function of the *mitochondria*?
 - What is the function of the *ribosomes*?
 - What is the function of the *Golgi*?
 - What is the function of the *endoplasmic reticulum*?


Eukaryotic cell



The nucleus contains the DNA




Nucleotide {



(a) DNA double helix

(b) Single strand of DNA

Overview of Gene Expression



LECTURE PRESENTATIONS

For **CAMPBELL BIOLOGY, NINTH EDITION**

Jane B. Reece, Lisa A. Urry, Michael L. Cain, Steven A. Wasserman, Peter V. Minorsky, Robert B. Jackson

Chapter 5

The Structure and Function of Large Biological Molecules - DNA

Lectures modified by Garrett Dancik

Lectures by
Erin Barley
Kathleen Fitzpatrick


Overview: The Molecules of Life

- All living things are made up of four classes of large biological molecules: carbohydrates, lipids, proteins, and nucleic acids


The Synthesis and Breakdown of Polymers

- A monomer is a building block of a polymer
 - DNA: the nucleotides (characters) A,C,G, and T
 - RNA: the nucleotides (characters) A,C,G, and U
 - Proteins: twenty kinds of amino acids (characters)
- A **dehydration reaction** occurs when two monomers bond together through the loss of a water molecule
- Polymers are disassembled to monomers by **hydrolysis**, a reaction that is essentially the reverse of the dehydration reaction

(a) Dehydration reaction: synthesizing a polymer




(b) Hydrolysis: breaking down a polymer



Relationship between DNA, RNA, and protein

- Genes are made of DNA, a **nucleic acid** made of monomers called nucleotides
- A gene is a unit of inheritance that codes for the amino acid sequence of a polypeptide




Role of Nucleic Acids

- Nucleic acids store, transmit, and help express hereditary information
- There are two types of nucleic acids
 - **Deoxyribonucleic acid (DNA)**
 - **Ribonucleic acid (RNA)**
- DNA provides directions for its own replication
- DNA directs synthesis of messenger RNA (mRNA) and, through mRNA, controls protein synthesis


The Components of Nucleic Acids

- Nucleic acids are polymers called **polynucleotides**
- Each polynucleotide is made of monomers called **nucleotides**
- Each nucleotide consists of a nitrogenous base, a pentose sugar, and one or more phosphate groups
- The portion of a nucleotide without the phosphate group is called a nucleoside



Figure 5.26



(a) Polynucleotide, or nucleic acid



(b) Nucleotide



(c) Nucleoside components

- Nucleoside = nitrogenous base + sugar
- There are two families of nitrogenous bases
 - **Pyrimidines** (cytosine, thymine, and uracil) have a single six-membered ring
 - **Purines** (adenine and guanine) have a six-membered ring fused to a five-membered ring
- In DNA, the sugar is **deoxyribose**; in RNA, the sugar is **ribose**
- Nucleotide = nucleoside + phosphate group

Nucleotide Polymers


- Nucleotide polymers are linked together to build a polynucleotide
- Adjacent nucleotides are joined by covalent bonds that form between the —OH group on the 3' carbon of one nucleotide and the phosphate on the 5' carbon on the next
- These links create a backbone of sugar-phosphate units with nitrogenous bases as appendages
- The sequence of bases along a DNA or mRNA polymer is unique for each gene

The Structures of DNA and RNA Molecules

- RNA molecules usually exist as single polypeptide chains
- DNA molecules have two polynucleotides spiraling around an imaginary axis, forming a **double helix**
- In the DNA double helix, the two backbones run in opposite $5' \rightarrow 3'$ directions from each other, an arrangement referred to as **antiparallel**
- One DNA molecule includes many genes

- Complementary base pairing
 - The nitrogenous bases in DNA pair up and form hydrogen bonds: adenine (A) always with thymine (T), and guanine (G) always with cytosine (C)
 - Complementary pairing can also occur between two RNA molecules or between parts of the same molecule
- In RNA, thymine is replaced by uracil (U) so A and U pair

Figure 5.27



© 2011 Pearson Education, Inc.

Genome sequencing

- The human genome project took over 13 years to complete and cost >\$3 billion (>\$1 / base pair sequenced)
 - Sequence assembly was one of the first bioinformatics challenges

The genomic revolution

- The \$1000 genome has arrived (sorta)
 - <http://www.forbes.com/sites/matthewherper/2014/01/14/the-1000-genome-arrives-for-real-this-time/>
 - Sequencing machines cost \$10 million
 - Can sequence 18,000 genomes / year
- Implications of cheap genomic sequencing
 - http://www.ted.com/talks/richard_resnick_welcome_to_the_genomic_revolution.html
 - What are they????

De novo sequence assembly

Unknown Genome: **AGCTATAGCGCTATCGTAGCTAGCGCTAGCT**

↓ Next-generation sequencing machine

AGCTATAG	CTATAGCG
GCTAGCGC	CGCTAGCT
TCTAGCGC	CGCTATCG
AGCTAGCG	ATCGTAGG

↓ Genome assembly software


AGCTATAG	GCTAGCGC	
TCTAGCGC	AGCTAGCG	
CTATAGCG	ATCGTAGG	CGCTAGCT
CGCTATCG		

↓

Reconstructed genome : **AGCTATAGCGCTATCGTAGCTAGCGCTAGCT**

Figure 1. Workflow of discovering the genome of a species

Cost per Genome



NIH National Human Genome Research Institute
genome.gov/sequencingcosts

The number of DNA nucleotides sequenced has grown exponentially

Current (Dec 2013):
169 billion bases

